

Comparing different responses of the cough reflex

Submission date 10/09/2021	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 16/09/2021	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 20/06/2025	Condition category Respiratory	<input type="checkbox"/> Individual participant data <input checked="" type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Cough challenge testing is widely used to see how sensitive someone's cough reflex is. This can be very useful when studying treatments for cough or how cough affects people with different lung conditions. The challenge involves a person inhaling increasing doses of a substance which usually makes people want to cough by irritating the throat and activating nerves in the airways. There are different pieces of equipment used to deliver these substances. The aim of this study is to compare the performance of two pieces of equipment to see if they give the same cough responses in people.

Who can participate?

Healthy volunteers aged 18 years and over

What does the study involve?

The study involves two visits. The first visit will involve completing a consent form after talking through the study with a researcher. Participants will then be asked about their medical history and any medications they are taking. Their height and weight will be measured and their BMI calculated. They will then be asked to perform a blowing test (spirometry) so that the researcher can check that they meet the eligibility criteria for the study. Participants will then be asked to complete a cough challenge. They will be fitted with a recording device, a cough monitor, throughout the challenge to enable the researcher to listen back and count how many times they cough. Capsaicin or citric acid will be given to them to inhale using one of the pieces of equipment. The dose will increase and they will be asked to take four breaths of each dose. The test will end when either the participant has inhaled all of the doses or when they ask to stop because the feeling from inhaling the capsaicin or citric acid is too unpleasant. They will be asked to fill in a questionnaire about how they found the challenge and then the blowing test will be repeated to ensure that their airways have not tightened. Participants will be asked to return 3-7 days later to repeat the cough challenge with the other piece of equipment.

What are the potential benefits and risks of participating?

No benefits.

Spirometry can cause light-headedness in participants.

Capsaicin (naturally occurring extract of red chilli peppers) and citric acid inhalation can cause

discomfort, including a temporary burning sensation in the throat, eye-watering, runny nose and coughing.

Cough challenge can cause bronchoconstriction, although this is rare in healthy volunteers.

Where is the study run from?

The NIHR Clinical Trials Facility and MFT Wythenshawe Hospital (UK)

When is the study starting and how long is it expected to run for?:

August 2021 to December 2024

Who is funding the study?

Manchester NIHR BRC charitable funds (UK)

Who is the main contact?

Joanne Mitchell

cough.research@manchester.ac.uk

Contact information

Type(s)

Scientific

Contact name

Prof Jaclyn Smith

ORCID ID

<https://orcid.org/0000-0001-8837-4928>

Contact details

2nd Floor ERC

Wythenshawe Hospital

Southmoor Road

Manchester

United Kingdom

M23 9LT

+44 (0)161 291 5031

jacky.smith@manchester.ac.uk

Type(s)

Public

Contact name

Mrs Joanne Mitchell

Contact details

North West Lung Research Centre

Wythenshawe Hospital

Southmoor Road

Manchester

United Kingdom

M23 9LT

+44 (0)161 291 5921
joanne.sale@manchester.ac.uk

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

Integrated Research Application System (IRAS)

288804

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

20-CAS-001

Study information

Scientific Title

A comparison and validation study of cough challenge responses evoked using two different bronchial provocation systems

Acronym

CasCaDe

Study objectives

Current study hypothesis as of 21/06/2024:

The aim of this study is to compare maximum cough responses to capsaicin and citric acid in healthy volunteers between two bronchial challenge systems: the Koko dosimeter and an alternative bronchial provocation system; ProvoX.

Previous study hypothesis:

The aim of this study is to compare maximum cough responses to capsaicin in healthy volunteers between two bronchial challenge systems: the Koko dosimeter and an alternative bronchial provocation system; ProvoX.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 13/09/2021, University of Manchester Research Ethics Committee 1 (Research Governance, Ethics and Integrity, 2nd Floor Christie Building, The University of Manchester, Oxford Road, Manchester, M13 9PL, UK; +44 (0)161 306 6000; research.ethics@manchester.ac.uk), ref: 2021-11272

Study design

Single-centre randomized cross over trial

Primary study design

Interventional

Study type(s)

Other

Health condition(s) or problem(s) studied

Cough challenge responses

Interventions

Current interventions as of 21/06/2024:

Participants will be asked to attend two visits. At the first visit, informed consent will be taken, followed by a screening visit to ensure study eligibility. This will involve the recording of medical history and current medications; measurement of height and weight; and spirometry to measure the participant's FEV1 and FVC. Upon confirmation of eligibility, the participant will be asked to complete either a capsaicin cough challenge or a citric acid cough challenge. This is done by inhaling doubling doses of capsaicin or citric acid using one of the systems. The challenge will be continued until the participant reaches their maximum tolerated dose. The cough challenge will be recorded on a VitaloJAK audio cough monitoring device.

Spirometry will be performed following the completion of the challenge. The participant will be asked to return for a second visit where they will repeat the spirometry and cough challenge using the other piece of equipment. The order of the equipment will be allocated in a randomised order created using Graphpad, a piece of statistical software with randomisation capabilities.

Previous interventions:

Participants will be asked to attend two visits. At the first visit, informed consent will be taken, followed by a screening visit to ensure study eligibility. This will involve the recording of medical history and current medications; measurement of height and weight; and spirometry to measure the participant's FEV1 and FVC. Upon confirmation of eligibility, the participant will be asked to complete a capsaicin cough challenge by inhaling doubling doses of capsaicin using one of the systems. The challenge will be continued until the participant reached their maximum tolerated dose. The cough challenge will be recorded on a VitaloJAK audio cough monitoring device.

Spirometry will be performed following the completion of the challenge. The participant will be asked to return for a second visit where they will repeat the spirometry and cough challenge using the other piece of equipment. The order of the equipment will be allocated in a randomised order created using Graphpad, a piece of statistical software with randomisation capabilities.

Intervention Type

Device

Phase

Not Applicable

Drug/device/biological/vaccine name(s)

-

Primary outcome(s)

Maximum cough response (Emax) is measured at visit 1 and visit 2

Key secondary outcome(s)

Current secondary outcome measures as of 21/06/2024:

1. Capsaicin and citric acid concentrations evoking at least 50% of the Emax cough number (ED50) is measured at visit 1 and visit 2
2. Concentration of capsaicin evoking at least two coughs (C2) is measured at visit 1 and visit 2
3. Capsaicin and citric acid concentrations provoking at least five coughs (C5) is measured at visit 1 and visit 2

Previous secondary outcome measures:

1. Capsaicin concentrations evoking at least 50% of the Emax cough number (ED50) is measured at visit 1 and visit 2
2. Concentration of capsaicin evoking at least two coughs (C2) is measured at visit 1 and visit 2
3. Capsaicin concentrations provoking at least five coughs (C5) is measured at visit 1 and visit 2

Completion date

04/12/2024

Eligibility

Key inclusion criteria

Current inclusion criteria as of 21/06/2024:

1. Aged ≥ 18 years
2. Normal spirometry (FEV1 and FVC %predicted of $\geq 80\%$)
3. No current or past history of chronic cough or any other respiratory disease
4. No current or history of chronic pain, irritable bowel syndrome, psychiatric illness or chronic headaches
5. No previous experience of performing cough challenges (capsaicin study arm only)
6. Must cough at least 2 times on a single inhalation of any dose of capsaicin/citric acid during the visit 1 cough challenge

Previous inclusion criteria:

1. Aged ≥ 18 years
2. Normal spirometry (FEV1 and FVC %predicted of $\geq 80\%$)
3. No current or past history of chronic cough or any other respiratory disease
4. No current or history of chronic pain, irritable bowel syndrome, psychiatric illness or chronic headaches
5. No previous experience of performing cough challenges
6. Must cough at least 2 times on a single inhalation of any dose of capsaicin during the visit 1 cough challenge

Participant type(s)

Healthy volunteer

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Total final enrolment

26

Key exclusion criteria

Current exclusion criteria as of 21/06/2024:

1. Current smoker or ex-smoker with >20 pack-year history, and/or <6 months abstinence
2. Upper respiratory tract infection within the last 4 weeks
3. Use of ACE inhibitors
4. Use of anti-depressants
5. Use of centrally acting medications that may alter the cough reflex e.g. opiates
6. History of drug or alcohol abuse
7. Current pregnancy or breastfeeding
8. Concomitant conditions that may alter cough reflex sensitivity e.g. diabetes mellitus, Parkinson's disease, cerebrovascular disease
9. Previous cough challenge experience (only applicable for capsaicin study arm)

Previous exclusion criteria:

1. Current smoker or ex-smoker with >20 pack-year history, and/or <6 months abstinence
2. Upper respiratory tract infection within the last 4 weeks
3. Use of ACE inhibitors
4. Use of anti-depressants
5. Use of centrally acting medications that may alter the cough reflex e.g. opiates
6. History of drug or alcohol abuse
7. Current pregnancy or breastfeeding
8. Concomitant conditions that may alter cough reflex sensitivity e.g. diabetes mellitus, Parkinson's disease, cerebrovascular disease
9. Previous cough challenge experience

Date of first enrolment

12/09/2021

Date of final enrolment

27/11/2024

Locations

Countries of recruitment

United Kingdom

England

Study participating centre

NIHR Clinical Research Facility - MFT Wythenshawe Hospital
Southmoor Road

Wythenshawe
Manchester
United Kingdom
M23 9LT

Sponsor information

Organisation

University of Manchester

ROR

<https://ror.org/027m9bs27>

Funder(s)

Funder type

Charity

Funder Name

Manchester NIHR BRC charitable funds

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are not expected to be made available as the study is purely a validation study and we are doing this research because the current equipment being used for cough challenge testing is being withdrawn from use and it is important that we find an alternative which produces reliable results comparable with the previous research carried out.

IPD sharing plan summary

Not expected to be made available

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Participant information sheet	version 1.2	10/09/2021	16/09/2021	No	Yes
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes