# Clips versus Ligatures in Thyroid Surgery

Submission date [X] Prospectively registered Recruitment status 16/12/2003 No longer recruiting [X] Protocol [ ] Statistical analysis plan Registration date Overall study status 22/12/2003 Completed [X] Results [ ] Individual participant data Last Edited Condition category 02/10/2017 Surgery

### Plain English summary of protocol

Not provided at time of registration

## Contact information

## Type(s)

Scientific

### Contact name

Prof Markus W Büchler

#### Contact details

University of Heidelberg Medical School Im Neuenheimer Feld 110 Heidelberg Germany 69120 +49 6221 56 6200 Markus Buechler@med.uni-heidelberg.de

## Additional identifiers

### Protocol serial number

KSC 02/2003

## Study information

#### Scientific Title

Clips versus Ligatures in Thyroid Surgery: a randomised controlled trial

### Acronym

CLIVIT-Trial

### Study objectives

To date, no clear scientific recommendation exists in regard to the available blood vessel ligation techniques in thyroid surgery. A systematic review comparing different blood vessel ligation techniques has not yet been performed. The literature on this topic shows several deficits thus reducing its internal and external validity. Main problems are underpowered sample sizes, missing data in regard to the surgical technique, lack of standardization of the various surgical techniques and of homogeneity of the used study designs.

Therefore a randomized controlled multicenter, patient-blinded, two-group parallel relevance study under standardized conditions is required in order to achieve high internal as well as external validity. This would allow the results to be generalized for thyroid surgery and also may have implications for other surgical settings. This study is one of the first studies designed and organized by the Study Center of the German Surgical Society (SDGC). Besides investigating the primary hypothesis, this study was also started with the aim of developing a nationwide structure for randomized trials using a common and surgical relevant procedure.

### Ethics approval required

Old ethics approval format

### Ethics approval(s)

Not provided at time of registration

### Study design

Randomised controlled trial

### Primary study design

Interventional

### Study type(s)

Treatment

### Health condition(s) or problem(s) studied

Thyroid surgery

#### **Interventions**

The standard surgical approach in both treatment groups will be the Dunhill-Procedure which consists of a hemithyroidectomy and a subtotal thyroidectomy (at least 2/3 of the gland). The randomisation will proceed when the thyroid gland has been exposed and the decision in favour of a Dunhill procedure is made. Initially the vessels of the upper pole of the thyroid (hemithyroidectomy side) are localised and dissected from medial. Afterwards they are ligated with the conventional clamp-tie technique (Novafil 3-0 or equivalent monofilament suture type). Time is measured from ligating the vessels of the upper pole to the removal of the complete specimen. All vessels except the upper pole vessels of both sides are ligated according to the randomised technique: either using the clip vessel ligation technique or occluding manually by single ligatures (Vicryl 3-0).

### Intervention Type

Procedure/Surgery

#### Phase

**Not Specified** 

### Primary outcome(s)

The primary objective of this study is to test whether there is a relevant reduction in operating time (>15 min), using the clip vessel occluding technique versus the conventional clamp-tie.

### Key secondary outcome(s))

Secondary objectives are divided into surgical and non-surgical categories. In the surgical category there are the following points: total duration of operation (skin incision until suture), weight of specimen, amount of postoperative bleeding, frequency of reoperation due to bleeding, recurrent laryngeal nerve paralysis (temporary or irreversible pre- and postoperative assessment of vocal cord function by ENT-specialist), wound infection rate and rate of impaired parathyroid gland function. The non-surgical category consists of duration of postoperative hospital stay.

To assess the relevance of the chosen endpoints, the following ten aspects have to be ranked from "most important" to "least important" by patients and surgeons (only once): Intraoperative complications, postoperative complications, length of hospital stay, voice function, dysphagia, death, postoperative pain, postoperative fatigue, convalescence of complete physical maximum resilience and cosmetic result.

### Completion date

31/12/2008

## Eligibility

### Key inclusion criteria

The trial population consists of hospitalised patients of the four participating Departments of Surgery, who are planned for an elective thyroid operation. These operations can be performed with either the clip vessel ligation technique or the conventional clamp-tie technique.

At study enrollment:

Age equal or greater than 18 years
Expected survival time more than 12 months
Patients with benign diseases of the thyroid gland scheduled for elective surgery
Euthyroid metabolism (normal level of TSH or T3/T4)
Normal function of the vocal cords
Informed consent

At the end of surgical exploration: Suitable for at least 2/3 resection of the thyroid

## Participant type(s)

Patient

### Healthy volunteers allowed

No

### Age group

Adult

## Lower age limit

#### Sex

All

### Key exclusion criteria

Malignant disease or high suspicion for malignancy (clinical and imaging evidence)

Nerve palsy

Graves' Disease

Coagulopathy

Current immunosuppressive therapy

Chemotherapy within 2 weeks before operation

Radiotherapy completed no longer than 8 weeks before operation

Severe psychiatric or neurologic disease

Drug- and/or alcohol-abuse

Participation in another intervention-trial with interference of intervention and outcome Inability to follow the instructions given by the investigator or the telephone interviewer Lack of compliance

### Date of first enrolment

05/03/2004

### Date of final enrolment

31/12/2008

## Locations

#### Countries of recruitment

Germany

Study participating centre
University of Heidelberg Medical School
Heidelberg
Germany
69120

## Sponsor information

### Organisation

University of Heidelberg Medical School (Germany)

### **ROR**

https://ror.org/038t36y30

## Funder(s)

### Funder type

Other

### Funder Name

Grant from the German Ministry of Research and Education and supported by the BBD-Aesculap® Company, Tuttlingen, Germany

## **Results and Publications**

Individual participant data (IPD) sharing plan

## IPD sharing plan summary

Not provided at time of registration

## **Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/10/2012		Yes	No
Protocol article	protocol	01/09/2006		Yes	No