New surgical technique for facial dimple creation

Submission date	Recruitment status	Prospectively registered
08/10/2024	No longer recruiting	[_] Protocol
Registration date	Overall study status	[] Statistical analysis plan
09/10/2024	Completed	[_] Results
Last Edited	Condition category	Individual participant data
09/10/2024	Surgery	[X] Record updated in last year

Plain English summary of protocol

Background and study aims

Surgical creation of facial dimples (dimpleplasty) was first introduced by Boo-Chai in 1962. Despite advancements in surgical approaches made in recent years, there remains a need for a reliable, minimally invasive method that ensures consistent results and high patient satisfaction. This study aims to evaluate a newly proposed surgical technique for facial dimpleplasty, focusing on patient satisfaction, and early recovery symptoms and complications.

Who can participate?

Patients aged between 18 and 30 years who requested dimpleplasty

What does the study involve?

Patients were randomly assigned to either the control group, who received the original dimpleplasty technique, or the test group, who underwent a new surgical technique. Patient-reported outcomes, such as the patient's satisfaction with their decision, outcome and face appearance, and complications were assessed.

What are the possible benefits and risks of participating? This study will determine the effectiveness of a newly developed surgical procedure for dimpleplasty compared to the conventional method. There is a risk of not achieving optimal results in some cases but the study team can manage these cases with other methods.

Where is the study run from? Damascus University (Syria)

When is the study starting and how long is it expected to run for? May 2022 to May 2024

Who is funding the study? Damascus University (Syria)

Who is the main contact? Dr Omar Hamdan, omarhamdancedr@gmail.com

Contact information

Type(s) Public, Scientific, Principal Investigator

Contact name Dr Omar Hamdan

ORCID ID https://orcid.org/0009-0000-6108-9625

Contact details Damascus Damascus Syria 00000 +963 (0)991184253 omarhamdancedr@gmail.com

Additional identifiers

EudraCT/CTIS number Nil known

IRAS number

ClinicalTrials.gov number Nil known

Secondary identifying numbers 0000

Study information

Scientific Title A new proposed surgical technique for dimpleplasty

Study objectives

Does the newly proposed surgical technique for dimpleplasty perform better than the conventional technique in terms of patient satisfaction, early recovery symptoms, and complications.

Ethics approval required

Ethics approval required

Ethics approval(s)

Approved 10/06/2022, Damascus University ethics committee (Damascus, Damascus, 0000, Syria; +963 (0)992964458; manager@hcsr.gov.sy), ref: 22918

Study design Comparative interventional randomized controlled study

Primary study design Interventional

Secondary study design Randomised controlled trial

Study setting(s) University/medical school/dental school

Study type(s) Quality of life, Efficacy

Participant information sheet

Not available in web format, please use the contact details to request a participant information sheet

Health condition(s) or problem(s) studied

Artificial facial dimpleplasty

Interventions

The first group received dimpleplasty following the technique described by Bao et al. (2007) (control group), while the second group underwent dimpleplasty using a new surgical technique proposed by the authors of this study (test group).

Randomization method: sealed envelope method

Control group:

All surgical procedures were performed under local anesthesia. The ideal location of the facial dimple was determined. The location of the dimple placement was marked on the cheek skin. The buccal mucosa was incised using a no.15 blade, with the incision measuring 2 to 3 mm in length and positioned below the papilla of the Stensen's duct. A syringe needle is then punctured through the cheek from the marked skin and pulled through the buccal mucosa incision. A 3-0 monofilament nylon suture is threaded into the pinhole of the syringe needle and drawn through using a vacuum extractor. Once the suture is pulled through, the needle is gradually withdrawn to the dermis. The needle's direction is then adjusted, puncturing through the dermis and muscle, and pulled through the buccal mucosa incision again. The suture was drawn through the pinhole of the syringe needle, which was then removed from the skin. This process sutures the active facial muscles and dermis together. The knot is tied, forming the dimple.

Test group:

All surgical procedures were performed under local anesthesia. The ideal location of the facial dimple was determined. The location of the dimple placement was marked on the cheek skin. A syringe needle was punctured through the cheek from the marked skin and pulled through the buccal mucosa incision. The buccal mucosa was incised using a no.15 blade, with the incision measuring 4 to 6 mm in length. Dissection of the mucosa was performed to reveal the underlying buccinator muscle. A rectangular portion of the muscle (6 × 4 mm) was excised using fine tissue scissors. The remaining muscle parts and the dermis were then connected using an

absorbable 3-0 Vicryl thread. A knot was made between the muscle and the dermis at the upper part of the incision, and another knot was made at the lower part of the incision. The buccal mucosa was closed with a 5-0 monofilament nylon suture.

Follow-up duration: 2 weeks

Intervention Type

Procedure/Surgery

Primary outcome measure

1. Patient's satisfaction with decision assessed using the FACE-Q instrument 1 week after the procedure

2. Patient's satisfaction with outcome assessed using the FACE-Q instrument 4 weeks after the procedure

3. Patient's satisfaction with overall face appearance assessed using the FACE-Q instrument at baseline and 2 weeks after

Secondary outcome measures

Post-operative complications evaluated using a four-point scale questionnaire 1 week after the procedure

Overall study start date 09/05/2022

Completion date 19/05/2024

Eligibility

Key inclusion criteria

1. Aged between 18 and 30 years

- 2. Capable of understanding the study procedures and committed to follow-up
- 3. Maintaining good oral hygiene

Participant type(s) Healthy volunteer

Age group Adult

Lower age limit 18 Years

Upper age limit 30 Years

Sex Female

Target number of participants

20

Total final enrolment

20

Key exclusion criteria
1. Presence of any facial deformities
2. Previous dimpleplasty procedures
3. Suffering from systemic diseases
4. Undergoing radiation therapy in the head and neck region within the last 6 months
5. Smoking

Date of first enrolment

22/11/2022 Date of final enrolment

22/03/2023

Locations

Countries of recruitment Syria

Study participating centre Damascus University Damascus Syria 0000

Sponsor information

Organisation

Damascus University

Sponsor details

Damascus Damascus Syria 0000 +963 (0)113341864 manager@hcsr.gov.sy

Sponsor type University/education Website http://www.damascusuniversity.edu.sy

ROR https://ror.org/03m098d13

Funder(s)

Funder type University/education

Funder Name Damascus University

Alternative Name(s) University of Damascus, , DU

Funding Body Type Government organisation

Funding Body Subtype Universities (academic only)

Location Syria

Results and Publications

Publication and dissemination plan Planned publication in a high-impact peer-reviewed journal

Intention to publish date 19/05/2025

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study will be available on request from Dr Omar Hamdan (omarhamdancedr@gmail.com) and in the publication related to it after the end of the research. Type of data that will be shared: demographic information (age, gender), location of the dimpleplasty, clinical measurements, and the photos of the clinical procedure.

IPD sharing plan summary

Available on request