A brief and early psychological intervention for individuals exposed to psychological trauma

Submission date	Recruitment status No longer recruiting	Prospectively registered	
27/08/2008		∐ Protocol	
Registration date	Overall study status	Statistical analysis plan	
06/10/2008	Completed	[X] Results	
Last Edited	Condition category	Individual participant data	
28/06/2019	Mental and Behavioural Disorders		

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Prof Alain Brunet

Contact details

Douglas Hospital Research Center Perry Building (E 4129) 6875 LaSalle Boulevard Montréal Canada H4H 1R3 +1 514 761 6131 x 4348 alain.brunet@douglas.mcgill.ca

Additional identifiers

Protocol serial number SR-4370

Study information

Scientific Title

A brief and early psychological intervention for individuals exposed to psychological trauma

Study objectives

A time by group interaction is expected whereby self-reported symptoms of posttraumatic stress disorder (PTSD) will be less severe in the intervention group than in the no intervention control group at the post-test, but not at the pre-test or at mid-treatment.

Ethics approval required

Old ethics approval format

Ethics approval(s)

McGill's Research and Ethic Board. Date of approval: 09/09/2002 (protocol number: A01-B03-02A)

Study design

Single-blind randomised controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Posttraumatic stress disorder

Interventions

This is a single-blind, randomised controlled trial with a treated group and another group on a waiting list completing the periodic assessments only.

This two-session dyadic intervention includes elements of psychoeducation and motivational interviewing, and targets communication between the patient and significant other, aiming to facilitate support, promote bi-directional disclosure, reduce disclosure-constraining behaviours, and improve coping. It promotes the disclosure of thoughts and emotions about the trauma in the natural environment of the dyad while attempting to reduce social constraints on disclosure and negative social support interactions. The intervention is led by a social worker or a nurse trained and supervised every second week by a clinical psychologist. The initial 90-minute session takes place within the first few weeks after the traumatic event. A 75-minute follow-up meeting is scheduled two weeks later.

Intervention Type

Other

Phase

Not Specified

Primary outcome(s)

PTSD self-reported symptoms as measured by the Impact of Event Scale Revised. We measured PTSD symptoms on average 26 days after trauma exposure (baseline, before treatment), then on average 40 days post-trauma at mid-treatment (after one treatment session), and 90 days after trauma exposure (a time when one can diagnose chronic PTSD), and then again 2 years after trauma exposure.

Key secondary outcome(s))

Negative social support. (It was hypothesised that the treatment would decrease the provision of negative social support from the significant other toward the trauma victim.). This was assessed using the Social Constraints Scale (SCS). Timepoints: 26 days (on average) after trauma exposure (baseline, before treatment), 40 days post-trauma at mid-treatment (after one treatment session), 90 days and 2 years after trauma exposure.

Completion date

14/03/2008

Eligibility

Key inclusion criteria

- 1. Both males and females, adults aged 18-65
- 2. Patients who present to the emergency room because of trauma exposure (in the last 10 days). Trauma is defined as an event eliciting a peritraumatic reaction of fear, helplessness, or horror.
- 3. Patients who speak either French or English
- 4. Medication free at trial onset
- 5. Patients who are in a medically stable condition
- 6. Patients who live in the Montreal metropolitan area
- 7. Patients who have a significant other whom they can bring to the therapy session
- 8. Patients who succeed at making an appointment with the therapist in a timely manner

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Upper age limit

65 years

Sex

Αll

Total final enrolment

74

Key exclusion criteria

- 1. Patients who have a suspected or confirmed traumatic brain injury
- 2. Patients who have a lifetime diagnosis of psychosis, substance/alcohol dependence, bipolar disorder, or mental retardation
- 3. Patients who have been clinically depressed in the last 2 years

Date of first enrolment 09/09/2002

Date of final enrolment 14/03/2008

Locations

Countries of recruitment
Canada

Study participating centre

Douglas Hospital Research Center

Montréal

Canada

H4H 1R3

Sponsor information

Organisation

Québec Fund for Research on Society and Culture (FQRSC) (Canada)

ROR

https://ror.org/00shpc021

Funder(s)

Funder type

Government

Funder Name

Québec Fund for Research on Society and Culture (Fonds Québécois de la Recherche sur la Société et la Culture) (FQRSC), formerly known as Québec Council for Social Research (Conseil Québécois de la Recherche Sociale) (CQRS) (Canada) (ref: SR-4370)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summaryNot provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	26/08/2013	28/06/2019	Yes	No