# Tactile acuity training for patients with chronic low back pain

Submission date	Recruitment status  No longer recruiting	Prospectively registered		
24/01/2012		☐ Protocol		
Registration date 06/02/2012	Overall study status Completed	Statistical analysis plan		
		[X] Results		
<b>Last Edited</b> 28/02/2014	Condition category  Musculoskeletal Diseases	[] Individual participant data		

## Plain English summary of protocol

Background and study aims

Tactile acuity (the keenness of the sense of touch) can be measured using the two-point discrimination test. This is the minimal distance between two points that can be detected on the skin surface. Tactile acuity is poor in patients with chronic pain conditions such as chronic regional pain syndrome and phantom limb pain (the sensation of pain in a limb that is missing /amputated). Tactile acuity training in these conditions can improve tactile acuity and, more importantly, it can reduce pain. The reduction in pain is strongly related to improvement in tactile acuity. Patients with chronic low back pain have poor tactile acuity. However, it is not known if tactile acuity training will improve pain and functioning in this patient group.

The main aim of this study was to investigate the how well tactile acuity training worked on pain and function for patients with persistent low back pain. The secondary aim of this study was to investigate how well tactile acuity training works on tactile acuity performance for patients with persistent low back pain.

Tactile acuity training has previously required considerable amounts of patient-therapist contact e.g. 30 minutes daily for two to three weeks. This level of contact may not be practical within the current NHS structure. As such, the intervention within this study had a large home exercise component delivered by an informal carer (friend/relative).

Another aim of this study was to find out what patients and informal carers thought of the home exercise part of the intervention.

#### Who can participate?

Participants were individuals ≥18 years of age with chronic low back pain and their informal carer.

#### What does the study involve?

Participants were randomly divided into two groups: the intervention group who received usual care physiotherapy and tactile acuity training or the placebo group who will receive usual care physiotherapy and placebo training.

Tactile acuity training involved two types of training: five point discrimination training and graphaesthesia training. In the former, five areas of the back are stimulated using either a pen top or a wine bottle cork. The participant provides feedback as to which area was stimulated and whether the pen top or bottle cork was used. The carer will then say if the patient was correct or not. In the graphaesthesia test, 60 letters would be traced onto the patients back using the carers finger and the patient asked to guess which letter was drawn. The carer then says if the patient was correct or not. The placebo training involves skin stimulation but no effort by the patient to describe what is happening e.g. they provide no feedback to the carer and vice versa.

Tactile acuity/placebo training involved three training sessions with a physiotherapist in the first two weeks of usual physiotherapy care. The remaining sessions were delivered as part of a home training package delivered by the informal carer (friend/relative).

Participation in the study lasted as long as usual care physiotherapy which was decided by the treating therapist (usually four to six weeks). Additionally, participants along with their informal carers were asked to take part in a focus group once all participants had completed their treatment. In this focus group we explored the attitudes of participants and their carers towards the intervention.

What are the possible benefits and risks of participating?

The risks associated with this study were minor. The likelihood of harm was low and the consequences would be small. Previous research using a very similar intervention and placebo group reported no adverse effects or risks and direct email correspondence with previous authors have confirmed this.

It was possible that the informal carers (friend/relative) would undergo some degree of psychological stress as a result of having to deliver the home training package. Arguably, an individual who is likely to be stressed by this was unlikely to volunteer. However, the informal carer received a high standard of training and support and the fact that no adverse effects or side effects were previously reported was emphasised. It was also emphasised that the informal carer could withdraw from the study at any time, without giving a reason, with no consequences. These measures should have helped to reduce any undue stress.

Where is the study run from?

The study was run from the Physiotherapy Department at the Friarage Hospital, Northallerton, UK.

When is study starting and how long is it expected to run for? The study ran from February 2012 to January 2013.

Who is funding the study?

The study was funded by the Teesside University Research Fund administered by Teesside University (UK).

Who is the main contact?
Dr Nicholas Harland PhD
nick.harland@nyypct.nhs.uk
Dr Cormac Ryan PhD
C.Ryan@tees.ac.uk

# Contact information

## Type(s)

Scientific

#### Contact name

Dr Nicholas Harland

#### Contact details

Team Lead of Extended Services
Friarage Hospital
Northallerton
United Kingdom
DL6 1JG
+44 (0)1609 763786
nick.harland@nyypct.nhs.uk

# Additional identifiers

#### Protocol serial number

11/NE/0328

# Study information

#### Scientific Title

Tactile acuity training for patients with chronic low back pain: A pilot randomised controlled trial

# Study objectives

There will be a statistically significant difference in pain and function between the tactile actuity training group and the placebo training group.

# Ethics approval required

Old ethics approval format

# Ethics approval(s)

NRES Committee North East C Northern & Yorkshire, UK, 06 January 2012, ref: 11/NE/0328

# Study design

Pilot randomised controlled trial

# Primary study design

Interventional

# Study type(s)

Treatment

# Health condition(s) or problem(s) studied

Chronic low back pain

#### **Interventions**

Tactile acuity training in the lower back region versus Sham Tactile Acuity (palcebo/control)

### Placebo (control group)

The placebo intervention will be delivered on three occasions in the physiotherapy department by the Physiotherapist and daily at home by an informal carer (friend/relative) as part of the home training package over the course of treatment. The training takes 30 minutes. There are two components to the placebo intervention.

#### 1. Tactile stimulation:

Will involve marking 8 number dots on the back area in most pain with a hypoallergenic pen specifically designed for skin use. These sites will be separated by the distance of the predetermined two point discrimination ability. The sites are then stimulated in a random order using either a big (wine bottle cork) or small (pen top) probe, again randomly applied. The interstimulus interval is 15 seconds. Three 6 minute blocks of 24 stimuli are performed with 3 minutes rest between blocks in accordance with previous protocols. The participant does not interact with the stimulus. Instead they engage in some other activity e.g. read a book or watch TV. The tactile stimulation session is performed daily and each session should provide 72 stimuli over 24 minutes.

#### 2. Back writing:

The area of most pain will have letters, approximately 1 inch high traced onto it. The researcher /informal carer will trace a series of 60 letters with the tip of their finger. Again no interaction with the patient will occur. This should take approximately five minutes.

## Intervention group

The intervention will be delivered on three occasions in the physiotherapy department by the Physiotherapist and daily at home by the informal carer as part of the home training package over the course of treatment. The training takes 30 minutes. There are two components to the intervention.

#### 1. Tactile acuity training:

Will involve marking 5-8 numbered dots on the back area in most pain. These sites will be separated by the distance of the predetermined two point discrimination ability. The sites are then stimulated in a random order using either a big (wine bottle cork) or small (pen top) probe, again randomly applied. The interstimulus interval is 15 seconds. Three 6 minute blocks of 24 stimuli are performed with 3 minutes rest between blocks in accordance with previous protocols. The participant will concentrate on the stimulus saying which of the five points has been stimulated and which probe was used. If after a number of sessions the participant begins to give the correct answers >90% of the time the marks should be moved 10% closer to one another making the task more difficult. The tactile stimulation session is performed daily and each session should provide 72 stimuli over 24 minutes.

#### 2. Back writing training:

The area of most pain will have letters, approximately 1 inch high traced onto it. The researcher will trace a series of 60 letters in a random order. The patient will be asked to guess which letter has been traced on their back. The first session is attended by the researcher, the participant and a friend/relative of the participant. At the training progresses patients may become very good at guessing the letter and the tester can make the game harder by writing the letters quicker or smaller. The training can be further progressed by writing words instead of individual letters. Numbers can also be written and added together to make the training more difficult and more varied.

# Intervention Type

#### Other

#### **Phase**

Not Applicable

# Primary outcome(s)

- 1. Pain (Numerical Rating Scale)
- 2. Function (Roland Morris Disability Questionnaire)

### Key secondary outcome(s))

Qualitative feedback about the intervention via interviews & focus groups

### Completion date

01/01/2013

# Eligibility

## Key inclusion criteria

- 1. More than 18 years of age
- 2. Pain duration of more than 6 months
- 3. Non-specific low back pain with or without leg pain
- 4. No red flags indicating potential serious pathology
- 5. No clinical signs of peripheral neuropathy such as:
- 5.1. Reduced muscle power
- 5.2. Loss of sensation or loss of reflex

### Participant type(s)

**Patient** 

# Healthy volunteers allowed

No

# Age group

Adult

## Lower age limit

18 years

#### Sex

All

#### Key exclusion criteria

- 1. Not having an informal carer (friend/relative) willing to facilitate with the home program
- 2. Being unable to write/speak English
- 3. Have learning difficulties or are not capable of giving consent

#### Date of first enrolment

01/02/2012

#### Date of final enrolment

# Locations

#### Countries of recruitment

United Kingdom

England

Study participating centre
Team Lead of Extended Services
Northallerton
United Kingdom
DL6 1JG

# Sponsor information

# Organisation

James Cook University Hospital (UK)

#### **ROR**

https://ror.org/02vqh3346

# Funder(s)

# Funder type

University/education

#### **Funder Name**

University Research Fund (URF), Teesside University (UK)

# **Results and Publications**

Individual participant data (IPD) sharing plan

# IPD sharing plan summary

Not provided at time of registration

# Study outputs

Output type	Details	Date created Date added	Peer reviewed?	Patient-facing?
Results article	results	26/02/2014	Yes	No
Participant information sheet	Participant information sheet	11/11/2025 11/11/2025	No	Yes