

# A double-blind, placebo controlled study to assess the effects of early intervention and/or treatment with epoetin alfa on anaemia in cancer patients receiving non-platinum containing chemotherapy

<b>Submission date</b> 19/08/2002	<b>Recruitment status</b> No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
<b>Registration date</b> 19/08/2002	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
<b>Last Edited</b> 19/02/2014	<b>Condition category</b> Haematological Disorders	<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

## Plain English summary of protocol

Not provided at time of registration

## Contact information

### Type(s)

Scientific

### Contact name

Dr - -

### Contact details

UKCCCR Register Co-ordinator  
MRC Clinical Trials Unit  
222 Euston Road  
London  
United Kingdom  
NW1 2DA

## Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

## Secondary identifying numbers

GN308

# Study information

## Scientific Title

## Study objectives

Not provided at time of registration

## Ethics approval required

Old ethics approval format

## Ethics approval(s)

Not provided at time of registration

## Study design

Randomised controlled trial

## Primary study design

Interventional

## Secondary study design

Randomised controlled trial

## Study setting(s)

Not specified

## Study type(s)

Treatment

## Participant information sheet

## Health condition(s) or problem(s) studied

Anaemia

## Interventions

Patients are randomised to one of two treatment arms:

1. Arm A: Patients receive 150 IU/Kg of Epoetin alpha
2. Arm B: Patients receive matching placebo

Epoetin alpha or placebo is given subcutaneously three times per week for the first four weeks (or the first on-study chemotherapy cycle, respectively). Haemoglobin level and/or reticulocyte count will then be used to determine whether the same dose (volume) or a doubled dose (volume) will be used for the remainder of the treatment.

## Intervention Type

Other

**Phase**

Not Specified

**Primary outcome measure**

Not provided at time of registration

**Secondary outcome measures**

Not provided at time of registration

**Overall study start date**

01/01/2003

**Completion date**

01/01/2004

## Eligibility

**Key inclusion criteria**

1. Diagnosis of non-myeloid malignancy, which requires non-platinum containing chemotherapy
2. Predicted chemotherapy of 12 to 24 weeks (three to six cycles) duration
3. Performance score zero to three
4. Life expectancy of more than six months
5. Aged over 18 years
6. Patients with acute leukaemia are excluded
7. No myeloablative chemotherapy
8. No uncontrolled hypertension
9. No evidence of untreated iron, folate or vitamin B12 deficiency

**Participant type(s)**

Patient

**Age group**

Adult

**Lower age limit**

18 Years

**Sex**

Both

**Target number of participants**

Not provided at time of registration

**Key exclusion criteria**

Does not comply with the above inclusion criteria

**Date of first enrolment**

01/01/2003

**Date of final enrolment**

01/01/2004

## Locations

### Countries of recruitment

England

United Kingdom

### Study participating centre

UKCCCR Register Co-ordinator

London

United Kingdom

NW1 2DA

## Sponsor information

### Organisation

Janssen-Cilag Ltd (UK)

### Sponsor details

Saunderton

High Wycombe

United Kingdom

HP14 4HJ

+44 (0)1494 567567

### Sponsor type

Industry

### Website

<http://www.janssen-cilag.co.uk>

### ROR

<https://ror.org/03qwpn290>

## Funder(s)

### Funder type

Industry

### Funder Name

## Results and Publications

### Publication and dissemination plan

Not provided at time of registration

### Intention to publish date

### Individual participant data (IPD) sharing plan

### IPD sharing plan summary

Not provided at time of registration