

Effects of reducing the parameters on dental x-ray machine on image quality and cell toxicity

Submission date 30/04/2020	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 06/05/2020	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 24/06/2021	Condition category Oral Health	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

A dental x-ray is essential to aid diagnosis and treatment planning. It is important to give clinicians information that cannot be obtained through physical examination. The increased use of x-rays increases radiation exposure. This has raised concern as exposure to radiation can cause cell damage and alterations which can increase the risk of developing oral cancer. One of the ways to reduce radiation exposure is by reducing the parameters available on the x-ray machine which are tube potential (kV) and tube current (mA). Although it is ideal and easy to reduce the parameters, the value and quality of the images must not be compromised. Thus, the aim of this study is to study the effect of reducing x-ray parameters on the diagnostic value and quality of the images. This study also looks at the presence of nuclear alterations after exposure to dental x-rays when the radiation dose is reduced.

Who can participate?

Patients aged 18 - 35 who attend the orthodontic clinic and were indicated for DPT and LC as part of routine assessment before starting orthodontic treatment

What does the study involve?

Patients are randomly allocated into two groups, control and intervention. The control group receive a standard radiation dose by setting x-rays machine in auto-mode. The intervention group receive a reduced radiation dose by decreasing the parameters on the x-ray machine. A buccal mucosa (the inner lining of the lips and cheeks) smear is taken using a brush and participants have a routine orthodontic assessment. Participants are then sent for a dental x-ray. They are called back to the clinic 10 days after the x-rays for a second buccal mucosa smear.

What are the possible benefits and risks of participating?

The participants in the intervention group may benefit from receiving a lower radiation dose. The risks include slight discomfort when the buccal mucosa smear is taken.

Where is the study run from?

Universiti Teknologi MARA (Malaysia)

When is the study starting and how long is it expected to run for?
March 2017 to August 2019

Who is funding the study?
Universiti Teknologi MARA (Malaysia)

Who is the main contact?
Dr Noraina Hafizan Norman
norainanorman@gmail.com

Contact information

Type(s)
Public

Contact name
Dr Liyana Ghazali

Contact details
Faculty of Dentistry
UiTM Sungai Buloh
Jalan Hospital
Sungai Buloh
Malaysia
47000
+60 (0)139288100
liyaghazali@yahoo.com

Additional identifiers

Clinical Trials Information System (CTIS)
Nil known

ClinicalTrials.gov (NCT)
Nil known

Protocol serial number
600-IRMI (5/1/6)

Study information

Scientific Title
Effects of scanning parameters reduction in dental radiographs on quality and diagnostics performance of the images and cytotoxicity of nuclei

Study objectives
1. There is no difference in image quality and diagnostic performance between standard and reduced scanning parameters.
2. There is a reduction in nuclei cytotoxicity when scanning parameters is reduced.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 05/07/2017, Uitm Research Ethics Committee (Universiti Teknologi MARA, Aras 3, Bangunan Wawasan, 40450 Shah Alam, Selangor, Malaysia; +60 (0)355448069; recsecretariat@uitm.edu.my), ref: REC/232/17

Study design

Single-centre double-blind parallel-group trial randomized controlled trial

Primary study design

Interventional

Study type(s)

Diagnostic

Health condition(s) or problem(s) studied

Digital dental panoramic tomogram and lateral cephalogram radiation exposure

Interventions

Participants were randomized 1:1 into control and intervention groups. The control group received standard scanning parameters as prescribed by the manufacturer by setting the radiographic machine in auto mode. Participants in the intervention group received reduced scanning parameters where the dental panoramic tomogram (DPT) was reduced by 60% and lateral cephalogram (LC) by 30%. All the participants were exposed to radiation once for each modality. Buccal mucosa smears were taken before radiation exposure and 10 days after the radiation exposure.

Intervention Type

Device

Phase

Not Applicable

Primary outcome(s)

1. Diagnostic performance of the images assessed using detection of anatomical landmarks (25 landmarks on dental panoramic tomogram and 35 landmarks on lateral cephalogram) after the x-rays were taken
2. The presence of nuclear alterations (pyknosis, karyolysis, karyorrhexis and nuclear budding) of buccal mucosa smear was noted (yes/no), taken before the exposure to x-rays and 10 days after the x-rays

Key secondary outcome(s)

Quality of the images assessed using a five-point rating scale after the x-rays taken (1= excellent, 2= more than adequately presented, 3= adequately presented 4= barely adequately presented, 5= inadequately presented)

Completion date

31/08/2019

Eligibility

Key inclusion criteria

1. Adults 18-35 years old
2. Indicated for dental panoramic tomogram and lateral cephalogram
3. Fit and healthy
4. Normal body mass index (BMI) 18.5 - 25

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Upper age limit

35 years

Sex

All

Total final enrolment

38

Key exclusion criteria

1. Patients contraindicated for radiographic examination
2. Pregnant
3. Occupational health exposure
4. Facial and skeletal abnormalities
5. Bone disease
6. Smoking or has stopped less than 6 months
7. Oral diseases including ulcers in the past 2 weeks

Date of first enrolment

01/02/2018

Date of final enrolment

30/06/2019

Locations

Countries of recruitment

Malaysia

Study participating centre
Universiti Teknologi MARA
Orthodontic Clinic
Faculty of Dentistry
Kampus Sungai Buloh
Jalan Hospital
Sungai Buloh
Malaysia
47000

Sponsor information

Organisation
Universiti Teknologi MARA

ROR
<https://ror.org/05n8tts92>

Funder(s)

Funder type
University/education

Funder Name
Universiti Teknologi MARA

Results and Publications

Individual participant data (IPD) sharing plan

For this study, the participants' level data are the x-ray images (DPT and LC) and slides containing buccal mucosa smears. All images are stored online on UiTM's Integrated Dental Records Management System (iDERMS) (<https://dentistry.uitm.edu.my/v2/index.php/en/>) and can only be assessed by clinician in-charged using UiTM's intranet system. X-rays and any patients' records are medico-legal documents thus the access is restricted. Prior to the commencement of the study, informed consent was obtained from patients.

IPD sharing plan summary

Stored in repository

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		01/03/2021	24/06/2021	Yes	No

