

A randomised study to compare the outcomes after realignment knee surgery versus non-surgical treatment with bespoke knee physiotherapy, for patients under 60 years of age with osteoarthritis of the knee

Submission date 17/01/2023	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 25/01/2023	Overall study status Ongoing	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 17/04/2025	Condition category Musculoskeletal Diseases	<input type="checkbox"/> Individual participant data <input checked="" type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Osteoarthritis (OA) is the leading cause of disability worldwide and most commonly affects the knee joint causing issues with mobility, quality of life, and ability to work. Treatment options include non-surgical management and surgery may also be considered. Knee replacement is successful in patients over 60 years but less so if under 60 as the implant wears out sooner, therefore alternative options are sought to delay knee replacement surgery as long as possible. High tibial osteotomy (HTO), is a surgical procedure where the bone is cut just below the knee joint and a small wedge is opened, to shift the person's weight away from the damaged part of the knee to a healthy part of the knee. This can decrease pain, improve function and delay or avoid the need for knee replacement. Personalised knee therapy (PKT) is a physiotherapist-delivered non-surgical focussed intervention consisting of a programme of exercise alongside exercise enabling pain relief (external braces if required) which aims to improve muscle control in the lower limb and knee joint, to shift weight away from the 'worn' part of the knee, help reduce pain and avoid the need for further surgery. In this study, the team want to find out if HTO is better at delaying or avoiding knee replacement surgery than PKT alone in patients under 60 years old by comparing these two interventions.

Who can participate?

Patients aged >18 and <60 years with old with symptomatic medial compartment knee OA who the treating orthopaedic surgeon considers a suitable candidate for medial opening wedge HTO

What does the study involve?

Patients will be enrolled at around 20 NHS hospitals in the UK to determine the clinical (pain relief, improvement in function, quality of life, return to work) and cost-effectiveness of both treatments at 24 months. A mixed method process evaluation in a subset of staff and patients will explore trial eligibility; recruitment and retention; acceptability of intervention

implementation including trial processes and collection of routine monitoring data; patient experience of taking part and the contextual factors that influence this.

For participants randomised to the surgical group:

1. They will be placed on a routine NHS waiting list to have the HTO surgery
2. They will have their surgery at their local hospital and follow the local process for having the surgery and recovering from the operation. After their operation, they will receive standard postoperative rehabilitation from their hospital

For participants randomised to the non-surgical group:

1. They will be referred to the local NHS Physiotherapy department at the hospital and receive the specialised PKT physiotherapy programme of rehabilitation for knee OA
2. PKT will be delivered at their local NHS physiotherapy department over six sessions within a period of 3-4 months

Participants in both arms of the study will get the same questionnaires at 12 and 24 months post-randomisation (either via post or email) to assess whether these treatments have worked.

What are the possible benefits and risks of participating?

Both treatment options (non-surgical and surgical) have been proven in previous studies to improve knee pain, reduce disability and delay or avoid the need for a knee replacement altogether.

Non-surgical treatment has the obvious advantage that it does not require an operation and all the risks that go with an operation. The non-surgical treatment takes around 3-4 months to deliver and requires participants to attend at least six sessions with the physiotherapist and commit to the tailored exercise programme over this time. Participants may also be offered bracing and steroid injections. The treatment is directed at improving symptoms from osteoarthritis and does not alter the alignment of the lower leg.

Where is the study run from?

The lead study site is The Royal Infirmary of Edinburgh - NHS Lothian with the trial management team in Edinburgh Clinical Trials Unit - University of Edinburgh.

When is the study starting and how long is it expected to run for?

August 2022 to July 2027

Who is funding the study?

National Institute for Health and Care Research (NIHR) (UK)

Who is the main contact for the study?

Mr Anish Amin (Chief Investigator), Anish.K.Amin@ed.ac.uk (UK)

Contact information

Type(s)

Principal investigator

Contact name

Mr Anish Amin

ORCID ID

<https://orcid.org/0000-0002-9768-7595>

Contact details

Department of trauma and orthopaedic surgery
Room S4334 (c/o Medical Secretary)
Royal Infirmary of Edinburgh
51 Little France crescent
Edinburgh
United Kingdom
EH16 4SA
+44 (0)131 242 6881
Anish.K.Amin@ed.ac.uk

Type(s)

Public

Contact name

Miss Rachel Penman

ORCID ID

<https://orcid.org/0000-0003-0297-6851>

Contact details

University of Edinburgh
level 2, NINE Edinburgh BioQuarter
9 Little France Road
Edinburgh
United Kingdom
EH16 4UX
+44 (0)131 651 9970
r.penman@ed.ac.uk

Additional identifiers**Clinical Trials Information System (CTIS)**

Nil known

Integrated Research Application System (IRAS)

306571

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

NIHR129820, CPMS 55238, IRAS 306571

Study information

Scientific Title

What is the clinical-effectiveness and cost-effectiveness of surgery with medial opening wedge high tibial osteotomy (HTO) compared with non-surgical treatment in the management of osteoarthritis (OA) of the knee in patients younger than 60 years?

Acronym

MOTION Trial

Study objectives

Determine whether the intervention is superior to the comparator by answering the following two research questions:

1. For patients aged <60 years old with medial compartment knee OA, what is the relative clinical effectiveness (pain relief, improvement in function, quality of life, return to work) of HTO compared with non-surgical management at 24 months?
2. For patients aged <60 years old with medial compartment knee OA, what is the relative cost-effectiveness of HTO compared to non-surgical management at 24 months and as modelled over a lifetime horizon?

Ethics approval required

Ethics approval required

Ethics approval(s)

approved 16/01/2023, South Central - Hampshire B Research Ethics Committee (2 The Square, Temple Quay, Bristol, BS1 6PN, United Kingdom; +44 (0)207 104 8088; hampshireb.rec@hra.nhs.uk), ref: 22/SC/0446

Study design

Multi-centre prospective randomized open blinded endpoint (PROBE) parallel-group controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Medial compartment knee osteoarthritis

Interventions

In this study, patients under 60 years old with knee OA who the treating orthopaedic surgeon considers a suitable candidate for medial opening wedge high tibial osteotomy (HTO) surgery will be selected to compare the two possible options (non-surgical versus HTO). Half of the participants will be randomised for surgery (HTO) and half will be randomised to undertake a personalised knee therapy (PKT) package. PKT consists of a core package of a progressive structured exercise programme and advice on pain management, lifestyle and if appropriate, weight management. If it falls within usual care, optional inclusion of additional manual therapy, steroid injection, insoles or bracing and treatment of any co-existing symptoms can be introduced. The package is delivered over a 12 – 16 week period and involves 6 face-to-face sessions of physiotherapy and a patient program of physiotherapy to complete at home. Patients will be asked to complete a diary to record their compliance with the programme and bring it along to each visit for the details to be recorded. Which of the two treatments patients

will get will be decided by chance, using a computer - half the patients will get one treatment, and half will get the other.

To compare the interventions, we will ask patients about their pain, knee function, quality of life and ability to work before and after the treatment. The research will be carried out at the local hospital or institution where the patient would normally be seen and treated.

Intervention Type

Mixed

Primary outcome(s)

Patients' opinions about their knee and associated problems measured using the patient-reported Knee Injury and Osteoarthritis Outcome Score (KOOS) at 24 months

Key secondary outcome(s)

1. Fidelity of interventions measured using the minimum requirements/acceptable variation (MRAV) proforma completed between 6 -12 months post-intervention (note that this interval may be different from the same period post-randomisation due to NHS waiting lists)
2. Patients' opinions about their knee and associated problems measured using the patient-reported Knee Injury and Osteoarthritis Outcome Score (KOOS) at 12 months post-randomisation.
3. Symptoms, pain, ADLs, sports/recreation, and quality of life measured using five separate KOOS subscales at 12- and 24-months post-randomisation
4. Function and pain measured using the Oxford Knee score (OKS) at 12 and 24- months post-randomisation
5. Artificial prosthesis awareness during daily activities measured using the Forgotten Joint Score-12 (FJS-12) at 12- and 24- months post-randomisation
6. Health-related quality of life measured using the EuroQol EQ-5D-5L score (EQ-5D-5L – EQ-5D) at 12- and 24- months post-randomisation
7. Sleep quality measure using the Pittsburgh Sleep Quality Index (PSQI) at 12- and 24-months post-randomisation
8. Employment status measured using a bespoke Return to Work/Employment Questionnaire developed at the lead centre at 12 and 24 months post-randomisation
9. Additional study-knee-related operative intervention measured using patient medical records at 12- and 24-months post-randomisation
10. Intraoperative and postoperative complications measured using patient medical records at 12 and 24 months post-randomisation
11. Health Economic Evaluation Outcomes derived from the EQ-5D score at 12- and 24- months post-randomisation
 - 11.1. Health and social care resource utilisation and associated NHS and personal social services (PSS) cost and Quality Adjusted life years (QALY) at 24 months
 - 11.2. Incremental Cost per QALY at 24 months
 - 11.3. NHS and PSS cost, and QALYs as modelled over a lifetime horizon to account for future impacts on the need for TKR and associated revision surgery and their timing relative to retirement
 - 11.4. Incremental Cost per QALY as modelled over a lifetime horizon to account for future impacts on the need for TKR and associated revision surgery and their timing relative to retirement

Completion date

31/07/2027

Eligibility

Key inclusion criteria

1. Patient aged <60 years old with symptomatic medial compartment knee OA who the treating orthopaedic surgeon considers a suitable candidate for medial opening wedge HTO

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Upper age limit

60 years

Sex

All

Total final enrolment

91

Key exclusion criteria

1. Aged <18 or >60 years old
2. Body mass index (BMI) >40
3. Patients considered for HTO but who DO NOT have any knee OA including:
 - 3.1. Offloading HTO for concomitant cartilage repair (No OA)
 - 3.2. Offloading HTO solely to treat ligamentous instability (ACL/PCL)
 - 3.3. Symptomatic avascular necrosis/osteonecrosis
 - 3.4. Correction of intraarticular or extraarticular post-traumatic knee deformity
4. Patients requiring double-level knee osteotomy for correction of deformity
5. History of inflammatory arthropathy including rheumatoid arthritis, gout, psoriasis
6. Previous high tibial or distal femoral osteotomy in the same or contralateral knee
7. Previous knee replacement (partial or total) in the same or contralateral knee
8. Cognitive impairment or inability to consent.
9. Inability to comply with study procedures.
10. Previous history of septic arthritis in the knee

Date of first enrolment

01/02/2023

Date of final enrolment

01/04/2025

Locations

Countries of recruitment

United Kingdom

England

Scotland

Wales

Study participating centre

NHS Lothian

Waverley Gate
2-4 Waterloo Place
Edinburgh
United Kingdom
EH1 3EG

Study participating centre

NHS Greater Glasgow and Clyde

J B Russell House
Gartnavel Royal Hospital
1055 Great Western Road Glasgow
Glasgow
United Kingdom
G12 0XH

Study participating centre

NHS Fife

Hayfield House
Hayfield Road
Kirkcaldy
United Kingdom
KY2 5AH

Study participating centre

NHS Grampian

Summerfield House
2 Eday Road
Aberdeen
United Kingdom
AB15 6RE

Study participating centre
University Hospitals Coventry and Warwickshire NHS Trust
Walsgrave General Hospital
Clifford Bridge Road
Coventry
United Kingdom
CV2 2DX

Study participating centre
Oxford University Hospitals
John Radcliffe Hospital
Headley Way
Headington
Oxford
United Kingdom
OX3 9DU

Study participating centre
Lewisham and Greenwich NHS Trust
University Hospital Lewisham
Lewisham High Street
London
United Kingdom
SE13 6LH

Study participating centre
Warrington and Halton Teaching Hospitals NHS Foundation Trust
Warrington Hospital
Lovely Lane
Warrington
United Kingdom
WA5 1QG

Study participating centre
Guys and St Thomas' NHS Foundation Trust
St Homas' Hospital
Westminster Bridge
London
United Kingdom
SE1 7EH

Study participating centre
East Suffolk and North Essex NHS Foundation Trust
Colchester Dist General Hospital
Turner Road
Colchester
United Kingdom
CO4 5JL

Study participating centre
St George's University Hospital NHS Foundation Trust
St George's Hospital
Blackshaw Road
Tooting
London
United Kingdom
SW17 0QT

Study participating centre
Frimley Health NHS Foundation Trust
Portsmouth Road
Frimley
Camberley,
United Kingdom
GU16 7UJ

Study participating centre
Whiston Hospital (site)
Whiston Hospital
Warrington Road
Prescot
United Kingdom
L35 5DR

Study participating centre
University Hospitals Birmingham NHS Foundation Trust
Queen Elizabeth Hospital
Mindelsohn Way
Edgbaston
Birmingham
United Kingdom
B15 2GW

Study participating centre

NHS Lanarkshire

14 Beckford Street

Hamilton

United Kingdom

ML3 0TA

Study participating centre

Kingston Hospital NHS Foundation Trust

Galsworthy Road

Kingston upon Thames

United Kingdom

KT2 7QB

Study participating centre

Salisbury District Hospital

Salisbury District Hospital

Odstock Road

Salisbury

United Kingdom

SP2 8BJ

Study participating centre

The Maidstone Hospital

Hermitage Lane

Maidstone

United Kingdom

ME16 9QQ

Study participating centre

Sunderland Royal Hospital

Kayll Road

Sunderland

United Kingdom

SR4 7TP

Sponsor information

Organisation

University of Edinburgh

ROR

<https://ror.org/01nrxf90>

Organisation

NHS Lothian

ROR

<https://ror.org/03q82t418>

Funder(s)**Funder type**

Government

Funder Name

National Institute for Health and Care Research

Alternative Name(s)

National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Results and Publications**Individual participant data (IPD) sharing plan**

The data sharing plans for the current study are unknown and will be made available at a later date.

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
HRA research summary			20/09/2023	No	No
Participant information sheet	version 3.0	10/01/2023	25/01/2023	No	Yes
Protocol file	version 2.0	06/12/2022	25/01/2023	No	No
Study website	Study website	11/11/2025	11/11/2025	No	Yes