Internet Smoking Cessation Intervention trial

Submission date	Recruitment status No longer recruiting	Prospectively registered		
22/12/2011		∐ Protocol		
Registration date	Overall study status Completed	Statistical analysis plan		
02/02/2012		[X] Results		
Last Edited 30/09/2014	Condition category Mental and Behavioural Disorders	Individual participant data		

Plain English summary of protocol

Background and study aims

Internet-based interventions can help smokers to stop compared with brief written materials or no intervention. However, they are not widely used in most countries, particularly by more disadvantaged smokers, and there is significant variation in their effectiveness. Therefore, there is a need for a generally available internet-based smoking cessation intervention that will be attractive across the social spectrum and of proven effectiveness. To address this need, a new website has been developed based on principles generated from theory, evidence, and the results from engaging disadvantaged smokers in user-testing and piloting. The aim of the current study is to evaluate the effectiveness of this new, interactive website and whether any identified efficacy translates across the social spectrum.

Who can participate?

You need to be:

- 1. A regular smoker
- 2. Over 18 years old
- 3. Prepared to make a serious attempt to quit smoking within the next month.

What does the study involve?

If you decide to take part you will be randomly assigned to receive support from one of two websites designed to help smokers to stop. We ask that you use your website to help you make a serious attempt to stop smoking within the next month.

Before quitting - During the time leading up to your quit attempt, we ask that you:

- 1. Answer some questions about your background, smoking history, current smoking, mood and craving levels.
- 2. Receive email reminders to visit the website.
- 3. Use the website and read the advice it provides to prepare for an attempt to quit smoking.

After quitting - During the month following your attempt to quit, we ask that you:

- 1. Receive email reminders to visit the website.
- 2. Use the website and read the advice it provides.
- 3. Answer questions about your smoking and the website.

We will email to ask questions about the website and your quit attempt, including whether you are still not smoking. If you are not smoking after 7 months we may need to confirm that. If selected, we ask that you provide a small sample of saliva to be analysed for a nicotine by-product. The sample will be destroyed after testing. We would send the kit through the post and you would just return it using a pre-paid envelope. In return for your trouble, you would receive a £20 gift voucher.

What are the possible benefits and risks of participating?

Possible risks or discomfort:

You may experience withdrawal symptoms such as irritability when you abstain from smoking, which you will be familiar with if you have previously attempted to stop.

Possible benefits:

Stopping smoking is the single most beneficial thing that smokers can do to improve their health. We hope that the results of this study will be of benefit to many thousands of other smokers who want to stop.

Financial considerations:

There are no financial costs to you for taking part. If you provide a saliva sample we will compensate you with a £20 gift voucher.

Where is the study run from?

The study is run online and is managed by researchers at University College London.

When is study starting and how long is it expected to run for?

The study started on 06/12/2011 and it is anticipated that recruitment will continue until 01/11/2012.

Who is funding the study?

The UKs National Prevention Research Initiative (NPRI). The NPRI is a national initiative made up of government departments, research councils and major medical charities that are working together to encourage and support research into chronic disease prevention.

Who is the main contact? Dr Jamie Brown stopsmoking@ucl.ac.uk

Contact information

Type(s)

Scientific

Contact name

Prof Robert West

Contact details

University College London Health Behaviour Research Centre Department of Epidemiology and Public Health 1-19 Torrington Place London United Kingdom WC1E 6BT

Additional identifiers

Protocol serial number

N/A

Study information

Scientific Title

A randomised controlled trial of a theory-based, interactive, internet-based smoking cessation intervention Stop Advisor

Acronym

ISCI

Study objectives

1. There will be an effect size of 3% difference between intervention and control conditions (i.e. 8% vs 5%) on the primary outcome of 6 months sustained and biochemically verified abstinence. 2. This efficacy will translate across the social spectrum.

A new smoking cessation website has been systematically developed based on principles generated from theory, evidence, and the results from engaging disadvantaged smokers in usertesting and piloting. The aim of the current study is to evaluate the effectiveness of this new, interactive website and whether any identified efficacy translates across the social spectrum. The proposed study is a two-arm randomised controlled trial, with participants randomised to the offer of the interactive website (intervention condition) or a non-interactive, static website (control condition) and followed up for up to 7 months post enrolment.

Ethics approval required

Old ethics approval format

Ethics approval(s)

University College London Research Ethics Committee, 14 February 2011 ref: 2808/001

Study design

Two-arm randomised controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Smoking cessation

Interventions

Participants randomised to the intervention or control condition and followed up for up to 7 months post-enrolment.

Content will consist of a set of behaviour change techniques (BCTs) linked together under the umbrella of an overarching theme. The theme will be that of an expert Stop Smoking Advisor who is both a ready source of useful information and a guide to help the smoker through the process of stopping, using a structured quit plan. Based on the success of the National Health Service (NHS) Stop Smoking Services, the website will follow as far as possible the guit plan that would be in operation in a well-run service, including the tailoring of information. The major difference will be that BCTs that are well suited to delivery by internet will be emphasized and those that require personal contact will be simulated as far as possible. Format will consist of a combination of menu items and tunnelled dialogue sessions presented according to a standard template. The information will be presented in the form of text, photographs, graphical images, video and audio clips. Users will follow a programme that takes them from preparation for the target quit date, to the quit date itself and then encourages them to report important information that the programme will use to help them overcome difficulties that they encounter. Prior to their quit date users will have access to a menu appropriate to their pre-quit status and up to five unique tunnelled dialogue sessions. Users can opt to take up to two weeks to get medication and two weeks to set a quit date. The pre-quit dialogue sessions will be tailored on assessing how soon they intend to quit, their intended use of medication, their success in obtaining and using medication, and reasons for guitting. The preparatory guidance they receive will focus on: ensuring that they acquire the appropriate medication and use it optimally, making necessary changes in their routines to minimise difficulties and urges to smoke after the target guit date, developing specific coping strategies for when difficulties arise, and having clear expectations about the natures of those difficulties. After their guit-date users will have access to menus appropriate to their post-quit status, and up to thirteen unique tunnelled dialogue sessions, which will be made available with decreasing frequency over the course of the first month following their guit date. The post guit dialogue sessions will be tailored on participants reports of:

- 1. Abstinence
- 2. Urges to smoke
- 3. Confidence in their ability to remain abstinent
- 4. Use of medication
- 5. Anticipated frequency of stressful or social events

The guidance they receive in response to this will involve specific advice on how to address these problems and plan for the future to minimise their occurrence.

The control will be a static website that presents brief and standard advice, with none of the presumed active ingredients of the interactive website. The advice offered on the control site about the timescale for setting a quit-date is equivalent to that on the interactive site: smokers will be encouraged to use medication; obtain medication within a fortnight; and set a quit date within two weeks of obtaining medication or deciding not to use it. By offering equivalent advice on quit dates, the average time since quit date at follow-up will be comparable between smokers from both conditions.

All participants will be followed up for up to 7 months post-enrolment.

Intervention Type

Other

Phase

Primary outcome(s)

Current primary outcome measures as of 10/01/2013:

Russell Standard 6 months sustained abstinence (RS6), defined as self-report of not smoking more than 5 cigarettes in the previous 6 months and no smoking in the previous week, verified by a saliva cotinine level of <15ng/ml at 7 months follow-up or by a saliva anabasine level of <1ng/ml in those recording a saliva cotinine level >14ng/ml and who also report using nicotine replacement therapy.

The final follow-up at 7 months post-enrolment should ensure that participants are at least 6 months post-quit: both control and intervention websites advise upon quit dates within one month of enrolment. Six months of abstinence is endorsed by both the Cochrane review group and National Institute for Health and Clinical Excellenc (NICE) as a sufficient basis for estimating long-term abstinence. Inclusion will be by intent to treat and participants whose smoking status cannot be determined will be counted as continuing smokers.

Previous primary outcome measures until 10/01/2013:

Russell Standard 6 months sustained abstinence (RS6), defined as self-report of not smoking more than 5 cigarettes in the previous 6 months and no smoking in the previous week, verified by a saliva nicotinine level of <15ng/ml at 7 months follow-up.

The final follow-up at 7 months post-enrolment should ensure that participants are at least 6 months post-quit: both control and intervention websites advise upon quit dates within one month of enrolment. Six months of abstinence is endorsed by both the Cochrane review group and National Institute for Health and Clinical Excellenc (NICE) as a sufficient basis for estimating long-term abstinence. Inclusion will be by intent to treat and participants whose smoking status cannot be determined will be counted as continuing smokers.

Key secondary outcome(s))

Current secondary outcome measures as of 10/01/2013:

The intervention and control groups will also be compared on a number of secondary measures including:

- 1. Point-prevalence abstinence, defined as a self-report of not smoking in the previous 7 days at 7 months follow-up, verified by a saliva cotinine level of <15ng/ml or by a saliva anabasine level of <1ng/ml in those recording a saliva cotinine level >14ng/ml and who also report using nicotine replacement therapy.
- 2. Self-report of abstinence in the previous 4 weeks at 2 months post-enrolment
- 3. Self-report of abstinence in the previous 3 months at 4 months post-enrolment
- 4. Self-report of a serious quit attempt in the previous 7 months at 7 months post-enrolment
- 5. Satisfaction ratings of the website at 2 and 7 months post-enrolment
- 6. Quantitative indices of website interaction, such as number of log-ins and page views Note: by definition, those classified as successes according to RS6 will also be classified as successes according to the secondary outcome measures 1 3.

Previous secondary outcome measures until 10/01/2013:

The intervention and control groups will also be compared on a number of secondary measures including:

- 1. Point-prevalence abstinence, defined as a self-report of not smoking in the previous 7 days at 7 months follow-up, verified by a saliva cotinine level of <15ng/ml
- 2. Self-report of abstinence in the previous 4 weeks at 2 months post-enrolment
- 3. Self-report of abstinence in the previous 3 months at 4 months post-enrolment
- 4. Self-report of a serious quit attempt in the previous 7 months at 7 months post-enrolment

- 5. Satisfaction ratings of the website at 2 and 7 months post-enrolment
- 6. Quantitative indices of website interaction, such as number of log-ins and page views Note: by definition, those classified as successes according to RS6 will also be classified as successes according to the secondary outcome measures 1 3.

Completion date

01/11/2012

Eligibility

Key inclusion criteria

- 1. Adult smokers from the UK
- 2. Smoke every day
- 3. Willing to make a serious quit attempt
- 4. Willing to use a stop-smoking website which sends email reminders
- 5. Willing to be followed up at 2, 4 and 7 months
- 6. Able to provide informed consent
- 7. Able to be contacted by email and telephone

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Key exclusion criteria

Does not meet inclusion criteria

Date of first enrolment

06/12/2011

Date of final enrolment

01/11/2012

Locations

Countries of recruitment

United Kingdom

England

Study participating centre

University College London

London United Kingdom WC1E 6BT

Sponsor information

Organisation

University College London (UK)

ROR

https://ror.org/02jx3x895

Funder(s)

Funder type

Research council

Funder Name

Medical Research Council (UK), ref: G0802035

Alternative Name(s)

Medical Research Council (United Kingdom), UK Medical Research Council, MRC

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Funder Name

National Prevention Research Initiative (NPRI) Phase 3 (UK), ref: 913685

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summaryNot provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/12/2014		Yes	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes
Study website	Study website	11/11/2025	11/11/2025	No	Yes