

# Pre-operative volume replacement versus usual care in diabetic patients having coronary artery bypass graft (CABG) surgery: a randomised controlled trial

<b>Submission date</b> 28/08/2008	<b>Recruitment status</b> No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
<b>Registration date</b> 29/10/2008	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
<b>Last Edited</b> 23/09/2019	<b>Condition category</b> Circulatory System	<input type="checkbox"/> Individual participant data

**Plain English summary of protocol**  
Not provided at time of registration

## Contact information

**Type(s)**  
Scientific

**Contact name**  
Prof Raimondo Ascione

**Contact details**  
Bristol Heart Institute  
Level 7 Queens Building  
Bristol Royal Infirmary (BRI)  
Marlborough Street  
Bristol  
United Kingdom  
BS2 8HW  
+44 (0)117 34 23286  
r.ascione@bristol.ac.uk

## Additional identifiers

## Study information

**Scientific Title**

Pre-operative volume replacement versus usual care in diabetic patients having coronary artery bypass graft (CABG) surgery: a randomised controlled trial

## Acronym

VeRDICT

## Study objectives

Post-operative incidence of renal insufficiency is lower and post-operative recovery faster, when diabetic patients are treated with volume replacement therapy (VR) prior to surgery.

## Ethics approval required

Old ethics approval format

## Ethics approval(s)

North Somerset & South Bristol REC, 25/02/2010, ref: 10/H0106/1

## Study design

Single-centre randomised controlled trial

## Primary study design

Interventional

## Study type(s)

Treatment

## Health condition(s) or problem(s) studied

Coronary heart disease and diabetes

## Interventions

Current interventions as of 01/05/2012

1. Volume replacement: CABG with or without cardiopulmonary bypass (CPB), with preoperative volume replacement therapy (1 ml/kg/hr of Hartmann's solution for 12 consecutive hours prior to surgery).
2. Usual care: CABG with or without CPB with conventional preoperative management (no preoperative fluids).

Previous interventions

The participants will be randomly allocated to the following two treatment groups in equal numbers:

1. Volume replacement: CABG with or without CPB, with pre-operative volume replacement therapy (1 ml/kg/hr of normal saline [N/saline] for 12 consecutive hours prior to surgery)
2. Usual care: CABG with or without CPB with conventional pre-operative management (no pre-operative fluids)

## Intervention Type

Procedure/Surgery

## Primary outcome(s)

Current primary outcome measures as of 15/05/2009:

Time until patients are classified as 'fit for discharge' since renal impairment is expected to impact on the risk of many post-operative complications. A patient must have normal

temperature, pulse and respiration, normal oxygen saturation on air, normal bowel function and be physically mobile in order to be classified as fit for discharge.

Previous primary outcome measures:

Time until patients are classified as 'fit-for-discharge' since renal impairment is expected to impact on the risk of many post-operative complications. In order to be classified as fit for discharge, a patient must have a chest X-ray with no evidence of pleural effusion requiring drainage, lung collapse/consolidation or pneumothorax, no suspected infection, normal routine blood tests and temperature and be physically mobile.

### **Key secondary outcome(s)**

Current secondary outcome measures as of 08/04/2013 (changes implemented as of 16/01/2013):

1. A participant's judgement about his or her readiness for discharge when the above criteria are met (too soon, about right, could have been discharged earlier);
2. Estimated GFR (eGFR) from serum creatinine measured from blood samples collected preoperatively (baseline, pre-trial intervention), and at 0, 12, 24, 36, 48, 72, 96 and 120 hours after the operation and the % of participants with GFR<60 mL/min on 2 of the 8 post-operative times;
3. Microalbumin/creatinine ratio measured in urine samples collected preoperatively (baseline, pre-trial intervention) and at 0, 24, 48 and 120 hours to assess microvascular disease and renal glomerular injury.
4. Tubular injury as expressed by N-acetyl glucosaminidase (NAG) release measured in urine samples collected preoperatively (baseline, pre-trial intervention) and at 0, 24, 48 and 120 hours in a consecutive sub-sample of 50 patients.
5. Acute Kidney Injury (AKI, doubling of baseline serum creatinine at any time); serum creatinine will be measured from blood samples collected preoperatively (baseline, pre-trial intervention) and at 0, 24, 48, 72, 96, and 120 hours; the peak of postoperative serum creatinine level will be used in relation to the preoperative value to calculate the incidence of AKI;
6. In-hospital mortality and other standard measures of morbidity as used in previous RCTs, e.g. post-operative myocardial infarction (MI), stroke, arrhythmia, need for haemodynamic support, renal failure and wound infection (including 6-8 week telephone ASEPSIS assessment);
7. Use of health care resources and associated costs, e.g. duration of operation, intensive care unit (ICU)/high dependency unit (HDU) and ward stay, additional interventions to treat complications, readmissions;
8. Coronary Revascularisation Outcome Questionnaire (CROQ) preoperatively (preoperative version) and at 3 months.
9. The following outcomes will be measured in a consecutive sub-sample of 40 patients.
  - 9.1. Preoperative blood glucose control, as measured by fasting blood glucose and haemoglobin A1c (HbA1c) prior to chest opening but after the intervention.
  - 9.2. MicroRNA and other biochemical predictors of health outcome in serum and plasma taken preoperatively (baseline, pre-trial intervention) and at 0, 24 and 120 hours after the operation, and also in any leftover material/specimens collected during surgery (this may include, but is not limited to: portions of internal mammary arteries with surrounding tissues, pericardial fluid, pericardial fat/adipose tissue, pericardium, waste blood).
  - 9.3. C-reactive protein as a marker of inflammation, measured preoperatively (baseline, pre-trial intervention) and at 0, 12, 24, 48, 72 and 120 hours after the operation.
  - 9.4. Cardiac damage as measured with serial troponin T release measured pre-operatively (baseline, pre-trial intervention) and at 0, 12, 24, 48, 72 and 120 hours after the operation.

Previous secondary outcome measures as of 01/05/2012:

1. A participant's judgement about his or her readiness for discharge when the above criteria are

met (too soon, about right, could have been discharged earlier)

2. Estimated GFR from serum creatinine measured from blood samples collected preoperatively, and at 0, 12, 24, 36, 48, 72, 96 and 120 hours after the operation) and the % of participants with GFR <60 mL/min on 2 of the 8 post-operative times

3. Microalbumin/creatinine ratio measured in urine samples collected preoperatively and at 0, 24, and 48 hours to assess microvascular disease and renal glomerular injury

4. Tubular injury as expressed by N-acetyl glucosaminidase (NAG) release measured in urine samples collected preoperatively and at 0, 24, and 48 hours in a consecutive sub-sample of 80 patients.

5. Acute Kidney Injury (AKI, doubling of baseline serum creatinine at any time); serum creatinine will be measured from blood samples collected preoperatively and at 0, 24, 48, 72, 96, and 120 hours; the peak of postoperative serum creatinine level will be used in relation to the preoperative value to calculate the incidence of AKI

6. In-hospital mortality and other standard measures of morbidity as used in previous RCTs, e.g., post-operative MI, stroke, arrhythmia, need for haemodynamic support, renal failure and wound infection (including 4-6 week telephone ASEPSIS assessment)

7. Use of health care resources and associated costs, e.g., duration of operation, ICU/HDU and ward stay, additional interventions to treat complications, readmissions

8. Coronary Revascularisation Outcome Questionnaire (CROQ) preoperatively (preoperative version) and at 3 months

Previous secondary outcome measures as of 15/05/2009:

1. A participant's judgement about his or her readiness for discharge when the above criteria are met (too soon, about right, could have been discharged earlier)

2. Estimated GFR from serum creatinine measured from blood samples collected preoperatively, and at 0, 12, 24, 36, 48, 72, 96 and 120 hours after the operation) and the % of participants with GFR <60 mL/min on 2 of the 7 post-operative times

3. Microalbumin/creatinine ratio measured in urine samples collected preoperatively and at 0, 24, and 48 hours to assess microvascular disease and renal glomerular injury

4. Tubular injury as expressed by N-acetyl glucosaminidase (NAG) release measured in urine samples collected preoperatively and at 0, 24, and 48 hours in a consecutive sub-sample of 80 patients.

5. Acute Kidney Injury (AKI, doubling of baseline serum creatinine at any time); serum creatinine will be measured from blood samples collected preoperatively and at 0, 24, 48, 72, 96, and 120 hours; the peak of postoperative serum creatinine level will be used in relation to the preoperative value to calculate the incidence of AKI

6. In-hospital mortality and other standard measures of morbidity as used in previous RCTs, e.g., post-operative MI, stroke, arrhythmia, need for haemodynamic support, renal failure and wound infection (including 4-6 week telephone ASEPSIS assessment)

7. Use of health care resources and associated costs, e.g., duration of operation, ICU/HDU and ward stay, additional interventions to treat complications, readmissions

8. Coronary Revascularisation Outcome Questionnaire (CROQ) preoperatively (preoperative version) and at 3 months

Previous secondary outcome measures:

1. A participant's judgment about his or her readiness for discharge when the above criteria are met (too soon, about right, could have been discharged earlier)

2. Estimated glomerular filtration rate (GFR) from serum creatinine measured from blood samples collected pre-operatively, and at 0, 4, 12, 24 and 48 hours after the operation) and the percentage of participants with GFR less than 60 mL/min on two of the five post-operative times

3. Renal glomerular and tubular injury as expressed by microalbumin/creatinine ratio and by N-acetyl glucosaminidase (NAG) release respectively and measured in urine samples collected pre-

- operatively and at 0, 24, and 48 hours in a consecutive sub-sample of 80 patients
4. Acute renal failure (ARF) (doubling of baseline serum creatinine at any time). Serum creatinine will be measured from blood samples collected pre-operatively and at 0, 24, 48, 72, 96, and 120 hours; the peak of post-operative serum creatinine level will be used in relation to the pre-operative value to calculate the incidence of ARF
  5. In-hospital mortality and other standard measures of morbidity as used in previous randomised controlled trials, e.g., post-operative myocardial infarction (MI), stroke, arrhythmia, need for haemodynamic support, renal failure and wound infection (including 4 - 6 week telephone ASEPSIS assessment)
  6. Use of health care resources and associated costs, e.g. duration of operation, intensive care unit (ICU)/high dependency unit (HDU) and ward stay, additional interventions to treat complications, readmissions
  7. Coronary Revascularisation Outcome Questionnaire (CROQ) pre-operatively (pre-operative version) and at 3 months

**Completion date**

31/08/2014

## Eligibility

**Key inclusion criteria**

Current inclusion criteria as of 15/05/2009:

1. Patients with diagnosed type I or type II diabetes, being treated with oral medication and/or insulin (i.e. not diet controlled only)
2. Both males and females, age >16 and <80 years
3. Undergoing elective or urgent, isolated first time coronary artery bypass graft (CABG) with or without cardiopulmonary bypass (CPB)
4. Left ventricular ejection fraction  $\geq 30\%$

Previous inclusion criteria:

1. Patients with diagnosed type I or type II diabetes
2. Both males and females, aged greater than 16 and less than 80 years
3. Undergoing elective or urgent, isolated first time coronary artery bypass graft (CABG) with or without cardiopulmonary bypass (CPB)
4. Left ventricular ejection fraction greater than or equal to 25%

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Adult

**Sex**

All

**Total final enrolment**

169

## Key exclusion criteria

1. Patients who have had previous cardiac surgery
2. Emergency or salvage operation
3. Chronic renal failure requiring dialysis
4. Current congestive heart failure
5. Left ventricular ejection fraction <30% (i.e. poor LV function)

Please note that the 5th exclusion criterion was updated as of 15/05/2009. The previous criterion was as follows:

5. Left ventricular ejection fraction less than 25%

## Date of first enrolment

01/06/2009

## Date of final enrolment

31/08/2014

## Locations

### Countries of recruitment

United Kingdom

England

### Study participating centre

**Bristol Heart Institute**

Bristol

United Kingdom

BS2 8HW

## Sponsor information

### Organisation

University Hospitals Bristol NHS Foundation Trust (UK)

### ROR

<https://ror.org/04nm1cv11>

## Funder(s)

### Funder type

Charity

## Funder Name

Garfield Weston Foundation (UK)

# Results and Publications

## Individual participant data (IPD) sharing plan

### IPD sharing plan summary

Not provided at time of registration

### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>	results	01/01/2020	23/09/2019	Yes	No
<a href="#">Protocol article</a>	protocol	19/06/2017		Yes	No
<a href="#">HRA research summary</a>			28/06/2023	No	No