

Comparison of outcomes following aortic valve replacement with two different types of valve substitutes

Submission date 17/02/2010	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 25/03/2010	Overall study status Completed	<input type="checkbox"/> Protocol
Last Edited 09/11/2023	Condition category Circulatory System	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Prof Magdi Yacoub

Contact details
Harefield Heart Science Centre
Hill End Road
Harefield
United Kingdom
UB9 6JH
-
m.yacoub@imperial.ac.uk

Additional identifiers

Study information

Scientific Title
A prospective randomised trial comparing autograft versus homograft aortic root replacement

Study objectives

Homografts and autografts have been used for many years with good clinical and haemodynamic results. In contrast to homografts, autografts are the only valve substitutes which ensure long-term viability of the aortic valve. We believe that this translates into clinically relevant endpoints following aortic root replacement.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Hillingdon Health Authority approved on the 12th January 1994

Study design

Single-centre prospective randomized comparison trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Aortic valve disease

Interventions

Two interventions will be compared:

1. Homograft aortic root replacement with coronary reimplantation
2. Autograft aortic root replacement with coronary reimplantation and replacement of the pulmonary root with a homograft

The operations will be carried out by a single surgeon (Sir Magdi Yacoub). The surgical techniques are well established and have been previously published.

Patients will be followed at 1 month, 6 months, 1 year and yearly thereafter with outpatient clinic appointments and echocardiographic evaluation. In patients with normal and stable echocardiographic results and no functional limitation, the follow-up will be extended to every 2 years.

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

Survival at 10 years

Key secondary outcome(s)

1. Incidence of the need for reoperation
2. Quality of life (QOL): assessed using the 36-item Short Form health survey (SF-36) quality of life standardised questionnaire, which will be sent by mail with a return envelope to all patients

3. Incidence of valve-related complications: endocarditis, major bleeding, thrombosis or thromboembolism
4. Specific echocardiographic parameters:
 - 4.1. Progression of transaortic pressure gradient
 - 4.2. Progression in the degree of aortic regurgitation and incidence of aortic regurgitation grade 3+ and 4+
 - 4.3. Changes in aortic root diameter measured at the sinuses of Valsalva and incidence of aortic root dilatation greater than 45 mm
 - 4.4. Changes in left ventricular end-diastolic and end-systolic diameters
 - 4.5. Changes in ejection fraction
 - 4.6. Progression of transpulmonary gradient through the homograft in the autograft group

Completion date

15/11/2001

Eligibility

Key inclusion criteria

1. Patients with symptomatic aortic valve disease requiring aortic valve replacement
2. Patients with concomitant aortic root dilatation and/or ascending aortic dilatation and aortic valve dysfunction requiring surgery
3. Patients with bicuspid aortic valve disease requiring aortic valve replacement
4. Patients with aortic valve endocarditis
5. Patients who have undergone previous cardiac surgery
6. Aged less than 69 years, either sex
7. Written informed consent

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

5 years

Upper age limit

69 years

Sex

All

Total final enrolment

228

Key exclusion criteria

1. Marfan syndrome
2. Reither's syndrome
3. Rheumatoid arthritis
4. Aged less than 5 years or greater than 69 years
5. Inability to consent
6. Other known disease potentially shortening life expectancy to less than 15 years
7. When completeness of follow-up is judged unlikely by the investigators

Date of first enrolment

15/05/1994

Date of final enrolment

15/11/2001

Locations

Countries of recruitment

United Kingdom

England

Study participating centre

Harefield Heart Science Centre

Harefield

United Kingdom

UB9 6JH

Sponsor information

Organisation

Royal Brompton and Harefield NHS Foundation Trust (UK)

ROR

<https://ror.org/02218z997>

Funder(s)

Funder type

Charity

Funder Name

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		14/08/2010		Yes	No
Other publications	Post hoc analysis	08/11/2023	09/11/2023	Yes	No