ISRCTN03720485 https://doi.org/10.1186/ISRCTN03720485

Effectiveness of adenoidectomy in children with recurrent upper respiratory tract infections

Submission date 30/05/2007	**Recruitment status** No longer recruiting	[_] Prospectively registered [_] Protocol [_] Statistical analysis plan [X] Results [_] Individual participant data
Registration date 30/05/2007	**Overall study status** Completed	
Last Edited 21/01/2013	**Condition category** Surgery	

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr Anne G.M. Schilder

Contact details
University Medical Centre Utrecht (UMCU)
Wilhelmina Children's Hospital
Department of Otorhinolaryngology
P.O. Box 85090
Utrecht
Netherlands
3508 AB
+31 (0)30 250 4004
A.Schilder@umcutrecht.nl

Additional identifiers

Study information

Scientific Title

Acronym
NOA (Nederlands Onderzoek Adenotomie)

Study objectives
This is a superiority trial testing the hypothesis that adenoidectomy is more effective than a watchful waiting strategy in children with recurrent upper respiratory tract infections.

Ethics approval required
Old ethics approval format

Ethics approval(s)
Approved by the Medical Ethical Committee UMC Utrecht on the 13th October 2006 (ref: NL14149.041.06).

Study design
Randomised, multicentre, active controlled, parallel group trial

Primary study design
Interventional

Study type(s)
Treatment

Health condition(s) or problem(s) studied
Adenoidectomy in children, common cold, upper respiratory tract infection, rhinosinusitis

Interventions
Adenoidectomy within six weeks versus watchful waiting.

Intervention Type
Procedure/Surgery

Phase
Not Specified

Primary outcome(s)
Upper respiratory tract infection with or without fever (38°C or higher); this will be measured continuously by means of symptom diaries and daily temperature measurements.

Key secondary outcome(s)
1. Acute otitis media and otitis media with effusion episodes, measured continuously by means of symptom diaries
2. Health related quality of life, measured at inclusion (0 months), 3 months, 12 months and 24 months follow-up
3. Cost-effectiveness, measured continuously by means of symptom diaries
4. Nasopharyngeal flora, investigated at inclusion (0 months), 3 months and 12 months follow-up
5. Exhaled nitric oxide, measured at inclusion (0 months), 3 months, 12 months and 24 months follow-up

Completion date
01/02/2010

Eligibility

Key inclusion criteria
Children aged 1 to 6 years selected for adenoidectomy primarily because of recurrent upper respiratory tract infections (common colds and rhinosinusitis).

Participant type(s)
Patient

Healthy volunteers allowed
No

Age group
Child

Lower age limit
1 years

Upper age limit
6 years

Sex
Not Specified

Key exclusion criteria
1. Age younger than 1 year or older than 6 years
2. Previous adenoidectomy or adenotonsillectomy
3. Tympanostomy tubes present
4. Selected for adenoidectomy combined with tympanostomy tubes
5. Downs syndrome
6. Craniofacial malformations (e.g. cleft lip or palate)

Date of first enrolment
01/04/2007

Date of final enrolment
01/02/2010

Locations

Countries of recruitment
Netherlands

Study participating centre
University Medical Centre Utrecht (UMCU)
Utrecht
Netherlands
3508 AB

Sponsor information

Organisation
University Medical Centre Utrecht (UMCU) (The Netherlands)

ROR
https://ror.org/04pp8hn57

Funder(s)

Funder type
Research organisation

Funder Name
The Netherlands Organisation for Health Research and Development (ZonMw) (The Netherlands)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary
Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Other publications	economic evaluation	01/02/2013		Yes	No
Thesis results	results	01/12/2012		No	No