ISRCTN04084517 https://doi.org/10.1186/ISRCTN04084517

Effect of nutritional intervention with ready to use therapeutic food (RUTF) on blood lipid profiles of moderate to severely malnourished adults receiving antiretroviral therapy (ART)

Submission date
20/01/2009

Recruitment status
No longer recruiting

Registration date
24/03/2009

Overall study status
Completed

Last Edited
24/03/2009

Condition category
Infections and Infestations

[_] Prospectively registered
[_] Protocol
[_] Statistical analysis plan
[_] Results
[_] Individual participant data
[_] Record updated in last year

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr Sandra Mutuma

Contact details
Valid International/ Nutrition
Addis Abba Road/ Chaholi Road
Alex Masala Close
Plot 4126
Lusaka
Zambia
P.O. Box 50719

Additional identifiers

Study information

Scientific Title

A randomised controlled study assessing the effect of nutritional intervention with ready to use therapeutic food (RUTF) on blood lipid profiles of moderate to severely malnourished adults receiving antiretroviral therapy (ART) in Lusaka, Zambia

Acronym
LRS (Lipid Research Study)

Study objectives
The use of high fat energy dense ready to use therapeutic food in HIV positive malnourished adults receiving antiretroviral therapy (ART) does not produce deleterious changes in their blood lipid profiles.

Ethics approval required
Old ethics approval format

Ethics approval(s)
University of Zambia Biomedical Research Ethics Committee, approved on 20/11/2008 (ref: 009-10-08)

Study design
Randomised controlled trial

Primary study design
Interventional

Study type(s)
Treatment

Health condition(s) or problem(s) studied
HIV and malnutrition

Interventions
Participants will be randomly allocated to the following two arms:

Intervention group: Participants will receive 2 jars/day of RUTF for 3 months, each jar weighs 250g (for 100g RUTF: energy content = 530 kcal, of which 47-59% are from oil and 10% are protein calories) in addition to prescribed ART regimen, nutritional advice and counselling.

Control group: Participants will receive ART, nutritional advice and counselling only

Total duration of interventions: 3 months

This trial will initially be a single-centre trial but may be extended to be a two-centre trial depending on recruitment rates.

Intervention Type
Other

Phase
Not Applicable

Primary outcome(s)
Total cholesterol/high density lipoprotein cholesterol (TC/HDL-C) ratio (TC/HDL-c ratio >= 5), assessed at baseline, Month 3, 6, 9 and 12 after enrolment into the study.

Key secondary outcome(s))
The following will be assessed at baseline, Month 3, 6, 9 and 12 after enrolment into the study:

1. Lipid profile:
1.1. Mean change in total cholesterol (TC)
1.2. Mean change in high density lipoprotein cholesterol (HDL-c)
1.3. Mean change in low density lipoprotein cholesterol (LDL-c)
1.4. Mean change in total triglycerides (TG)

2. Body composition:
The endpoints will be the correlation between change in TC and in:
2.1. Body fat mass proportion
2.2. Fat free mass (FFM) proportion
2.3. Fat-free mass index (FFMI) which controls for height

3. Nutritional status:
The endpoints will be the correlation between change in TC and in:
3.1. Weight (kg)
3.2. MUAC (mm)
3.3. BMI (kg/m^2)
3.4. Waist circumference

Note: Some of the secondary endpoints (e.g., weight, MUAC, waist measurements) may be taken at two weekly intervals from baseline up to Month 3 as well as at the timepoints indicated above.

Completion date
30/06/2010

Eligibility

Key inclusion criteria
1. Eligible for ART or has started ART within last 2 weeks
2. Both males and females, age >=18 years to <=49 years
3. mid-upper arm circumference (MUAC) <22.0 cm
4. Body mass index (BMI) <17.0 kg/m^2
5. CD4 Count >50
6. Signed informed consent

Participant type(s)
Patient

Healthy volunteers allowed
No

Age group
Adult

Lower age limit
18 years

Sex
All

Key exclusion criteria
1. Pre-existing diabetes
2. History of cardiovascular disease (CVD) including hypercholestrolaemia, hypertension
3. On other medications that may have an effect on blood lipid profiles (besides routine HIV prophylaxis and tuberculosis [TB] treatment)
4. CD4 count <50
5. Unable to tolerate solid foods
6. Failed appetite test
7. Subjects with pacemakers or any implantable electronic devices
8. Pregnancy or lactation
9. Psychiatric illness
10. Nut allergies
11. Consent declined
12. Any other reason why the consenting investigator thinks it is not appropriate for them to take part

Date of first enrolment
26/01/2009

Date of final enrolment
30/06/2010

Locations

Countries of recruitment
Zambia

Study participating centre
Valid International/ Nutrition
Lusaka
Zambia
P.O. Box 50719

Sponsor information

Organisation
Valid International (UK)

ROR

https://ror.org/00sb6vz77

Funder(s)

Funder type
Research organisation

Funder Name
Valid international (UK)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary
Not provided at time of registration