ISRCTN05754020 https://doi.org/10.1186/ISRCTN05754020

Re-stitching of a broken down perineal wound compared to leaving it to heal naturally

Submission date	Recruitment status	[X] Prospectively registered
08/12/2008	No longer recruiting	[X] Protocol
Registration date	**Overall study status**	[_] Statistical analysis plan
16/01/2009	Completed	[X] Results
Last Edited	**Condition category**	[_] Individual participant data
20/10/2017	Pregnancy and Childbirth	

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Miss Lynn Dudley

Contact details
University Hospital of North Staffordshire NHS Trust
Treatment Centre, Children's Services and Obstetrics
Academic Unit of Obstetrics and Gynaecology
Maternity Building, City General Site
Newcastle Road
Stoke-on-Trent
Staffordshire
United Kingdom
ST4 6QG
+44 (0)1782 552 058/434
lynn.dudley@uhns.nhs.uk

Additional identifiers

Protocol serial number
9098

Study information

Scientific Title
Perineal REpair following Vaginal delivery complicated by an Infected Episiotomy Wound: a feasibility study for a randomised controlled trial

Acronym
PREVIEW

Study objectives
Many women will require suturing to facilitate healing of the trauma site. However, practice varies widely between care given and established professionally agreed standards. There is limited data on the prevalence and consequences of perineal wound infection. In addition, there is only a small amount of information relating to the impact that perineal wound infection has on women's well-being during the immediate and long-term post-natal period. Anecdotal evidence suggests the number of women reporting perineal infections and dehiscence in the community is increasing; however, systems to track these complications following hospital discharge are lacking. Given that postpartum management of perineal trauma is a core component of routine maternity care it is vital that a true estimate of the problem is established using standardised definitions of wound infection and at the same time determine best practice when treating dehisced perineal wounds.

Hypotheses:
1. What is the prevalence of perineal wound infection and dehiscence in the UK?
2. What factors at the time of primary repair are most likely to be associated with perineal wound infection and dehiscence prior to discharge to the community?
3. What factors following discharge home are most likely to be associated with perineal wound infection and dehiscence in the community?
4. What are women's experiences of perineal infection and dehiscence and what types of information and support are most likely to benefit their post-natal recovery?
5. What is the best management for perineal wound infection and wound dehiscence?
6. What is the feasibility of conducting a definitive randomised controlled trial (RCT) comparing re-suturing of dehisced perineal wounds versus healing by secondary intention and what are the implications in terms of health benefits and costs?

Ethics approval required
Old ethics approval format

Ethics approval(s)
North Wales Research Ethics Committee, 29/04/2010, ref: 10/WNO03/16

Primary study design
Interventional

Study design
Computer-randomised controlled feasibility study

Study type(s)
Treatment

Health condition(s) or problem(s) studied
Dehisced perineal wounds

Interventions
The participants will be computer randomised into either immediate resuturing of their dehisced perineal wound in comparison to healing by secondary intention. Both groups will have an independent assessment of their perineal wound at 2 weeks and 6 weeks after trial entry. Both groups of participants will be asked to complete a questionnaire at 6 weeks, 3 months and 6 months after trial entry.

Intervention Type
Other

Phase
Not Applicable

Primary outcome(s)
Time taken for the dehisced perineal wound to heal, assessed at 2 weeks, 6 weeks, 3 months and 6 months, respectively.

An independent assessment of the primary outcome will be conducted at 2 weeks and 6 weeks using a questionnaire format, the title of the questionnaire being:
'PREVIEW' Independent Perineal Assessment 2 weeks, and
'PREVIEW' Independent Assessment 6 weeks

Mothers will also complete questionnaires at 6 weeks, 3 months and 6 months, respectively, also assessing primary and secondary outcomes, the title of these questionnaires being:
'PREVIEW' Mothers questionnaire 6 weeks
'PREVIEW' Mothers questionnaire 3 months
'PREVIEW' Mothers questionnaire 6 months

Key secondary outcome(s)
1. Woman's satisfaction with aesthetic results of perineal wound at 6 months post-natal
2. Pain at 6 weeks, 3 and 6 months post-natal
3. Dyspareunia (painful intercourse) at 3 - 6 months
4. Rates of breastfeeding

Mothers will complete questionnaires at 6 weeks, 3 months and 6 months respectively, assessing primary and secondary outcomes, the title of these questionnaires being:
'PREVIEW' Mothers questionnaire 6 weeks
'PREVIEW' Mothers questionnaire 3 months
'PREVIEW' Mothers questionnaire 6 months

Completion date
01/04/2011

Eligibility

Key inclusion criteria
1. Women (aged 18 - 40 years) referred to the perineal care clinic at the University Hospital of North Staffordshire
2. Dehisced perineal wound (spontaneous second, third or fourth tear or episiotomy)
3. Occurs within two weeks following childbirth

Participant type(s)
Patient

Healthy volunteers allowed
No

Age group
Adult

Lower age limit
18 Years

Upper age limit
40 Years

Sex
Female

Key exclusion criteria
1. Women that have not given their written consent to participate in the study
2. Women who have delivered a stillborn infant
3. Women under the age of 16 years
4. Women who cannot speak English or cannot read or write

Date of first enrolment
01/04/2009

Date of final enrolment
01/04/2011

Locations

Countries of recruitment
United Kingdom

England

Study participating centre
University Hospital of North Staffordshire NHS Trust
Staffordshire
United Kingdom
ST4 6QG

Sponsor information

Organisation
University Hospital of North Staffordshire NHS Trust (UK)

Funder(s)

Funder type
Charity

Funder Name
Smith and Nephew Foundation

Alternative Name(s)

Funding Body Type
Private sector organisation

Funding Body Subtype
Trusts, charities, foundations (both public and private)

Location
United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary
Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	nested qualitative study results	10/02/2017		Yes	No
Results article	results	10/02/2017		Yes	No
Protocol article	protocol	24/07/2012		Yes	No