

Randomised double-blind prospective controlled trial of intercostal nerve block for post-operative pain after bilateral thoracoscopic sympathectomy

Submission date 30/09/2004	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
Registration date 30/09/2004	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
Last Edited 21/03/2017	Condition category Signs and Symptoms	<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr David Raitt

Contact details

University Hospitals of Leicester
c/o Research and Development Office
Leicester General Hospital NHS Trust
Leicester
United Kingdom
LE1 4PW
+44 (0)116 258 4109
nicola.turner@uhl-tr.nhs.uk

Additional identifiers

Protocol serial number

N0123138366

Study information

Scientific Title

Randomised double-blind prospective controlled trial of intercostal nerve block for post-operative pain after bilateral thoracoscopic sympathectomy

Study objectives

To assess the efficacy of thoracoscopic intercostal nerve block by laevobupivacaine in alleviating immediate postoperative pain in patients undergoing bilateral thoracoscopic sympathectomy

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Study design

Randomised double-blind prospective controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Post-operative pain

Interventions

Randomised double-blind prospective controlled trial

Intervention Type

Drug

Phase

Not Applicable

Drug/device/biological/vaccine name(s)

Laevobupivacaine

Primary outcome(s)

An assessment of the efficacy of thoracoscopic intercostal nerve block by laevobupivacaine in alleviating immediate postoperative pain in patients undergoing bilateral thoracoscopic sympathectomy

Key secondary outcome(s)

Not provided at time of registration

Completion date

31/07/2003

Eligibility

Key inclusion criteria

Patients having undergone bilateral thoracoscopic sympathectomy

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Key exclusion criteria

Not provided at time of registration

Date of first enrolment

01/08/2002

Date of final enrolment

31/07/2003

Locations**Countries of recruitment**

United Kingdom

England

Study participating centre

University Hospitals of Leicester

Leicester

United Kingdom

LE1 4PW

Sponsor information**Organisation**

Department of Health

Funder(s)

Funder type

Government

Funder Name

University Hospitals of Leicester NHS Trust (UK)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration