

# Evaluation of effectiveness of pain control in patients after total hip replacement using the anaesthetic continuous-infusion device with bupivacaine

<b>Submission date</b> 03/03/2007	<b>Recruitment status</b> No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered
<b>Registration date</b> 13/04/2007	<b>Overall study status</b> Completed	<input type="checkbox"/> Protocol
<b>Last Edited</b> 09/05/2019	<b>Condition category</b> Musculoskeletal Diseases	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data

**Plain English summary of protocol**  
Not provided at time of registration

## Contact information

**Type(s)**  
Scientific

**Contact name**  
Dr Dave Weichih Chen

**Contact details**  
Orthopaedic Department  
Chang Gung Memorial Hospital  
Taipei-Linkou Medical Center  
5 Fu-Shing St., Kweishan  
Taoyuan  
Taiwan  
333  
-  
achih121@ms75.hinet.net

## Additional identifiers

## Study information

Scientific Title

Evaluation of effectiveness of pain control in patients after total hip replacement using the anaesthetic continuous-infusion device with bupivacaine

### **Study objectives**

The Anaesthetic Continuous-Infusion Device (ACID) with bupivacaine is effective for post-Total Hip Replacement (THA) pain control.

### **Ethics approval required**

Old ethics approval format

### **Ethics approval(s)**

Approval received from the Institutional Review Board of the Chang Gung Memorial Hospital on the 25th December 2006 (ref: 95-1192B).

### **Study design**

Randomised double-blind placebo-controlled study

### **Primary study design**

Interventional

### **Study type(s)**

Treatment

### **Health condition(s) or problem(s) studied**

Osteoarthritic hip

### **Interventions**

Group I (25 patients): ACID intra-articular (above lesser trochanter) and normal saline 100 ml

Group II (25 patients): ACID inter-muscular (between tensor fascia lata and gluteal medius) and normal saline 100 ml

Group III (25 patients): ACID intra-articular (above lesser trochanter) and bupivacaine 0.5% 100 ml

Group IV (25 patients): ACID intra-articular (between tensor fascia lata and gluteal medius) and bupivacaine 0.5% 100 ml

### **Intervention Type**

Drug

### **Phase**

Not Specified

### **Drug/device/biological/vaccine name(s)**

Bupivacaine

### **Primary outcome(s)**

Pain control

### **Key secondary outcome(s)**

Improve post-operative hip function and rehabilitation programs

### **Completion date**

31/07/2009

## Eligibility

### Key inclusion criteria

Consecutive 100 patients with indications for total hip replacement.

### Participant type(s)

Patient

### Healthy volunteers allowed

No

### Age group

Not Specified

### Sex

Not Specified

### Total final enrolment

92

### Key exclusion criteria

Patients who refuse to join the clinical trial.

### Date of first enrolment

01/08/2007

### Date of final enrolment

31/07/2009

## Locations

### Countries of recruitment

Taiwan

### Study participating centre

Orthopaedic Department

Taoyuan

Taiwan

333

## Sponsor information

### Organisation

Republic of China National Science Council (Taiwan)

**ROR**

<https://ror.org/02kv4zf79>

## Funder(s)

**Funder type**

Government

**Funder Name**

Republic of China National Science Council (Taiwan)

## Results and Publications

**Individual participant data (IPD) sharing plan**

**IPD sharing plan summary**

Not provided at time of registration

### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>	results	01/05/2010	09/05/2019	Yes	No