

Comparison between laser and diathermy assisted posterior cordotomy for bilateral vocal cord abductor paralysis

Submission date 27/02/2013	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 27/03/2013	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 06/01/2014	Condition category Respiratory	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Bilateral vocal cord paralysis (BVCP) is a potentially life threatening condition frequently requires surgical intervention called cordotomy. The objectives of treatment of BVCP are to achieve adequate airway and to preserve voice quality and laryngeal competence. The study compares two types of cordotomy.

Who can participate?

All patients with BVCP for one year at least with a respiratory chink of a maximal width of 4 mm or less and who need cordotomy.

What does the study involve?

Over a period of three years participants will be randomly allocated to one of two groups; Group (A) will be treated with laser assisted posterior cordotomy and Group (B) will be treated with diathermy assisted posterior cordotomy. The choice of the group will be decided by a process called randomisation, which is like a coin toss. During and at the end of the study, dyspnea severity, voice quality and aspiration will be assessed and compared between both groups.

What are the possible benefits and risks of participating?

There will be mostly improvement of the airway. But there will be benefits to future best methods of treating BVCP.

The main risk of cordotomy is laryngeal edema, decrease quality of voice or aspiration. All patients will receive routine care, safety procedures to monitor airway and follow up for cordotomy.

Where is the study run from?

This study was performed in Zagazig University Hospitals (Egypt).

When is study starting and how long is it expected to run for?

The study lasted from February 2008 to February 2011 on 20 patients. Participants were for a period of two to three years.

Who is funding the study?
Zagazig University Hospitals (Egypt)

Who is the main contact?
Dr Mohammad Waheed El-Anwar
mwenteg@yahoo.com

Contact information

Type(s)
Scientific

Contact name
Dr Mohammad El-Anwar

Contact details
Otorhinolaryngology, Head and Neck Surgery Department
Faculty of Medicine
Zagazig University
Zagazig
Egypt
44519
+20 552309843
mwenteg@yahoo.com

Additional identifiers

Study information

Scientific Title
Comparison between laser and diathermy assisted posterior cordotomy for bilateral vocal cord abductor paralysis: a prospective randomized controlled trial

Study objectives
It is hypothesised that the results of laser assisted posterior cordotomy will be better than diathermy assisted posterior cordotomy for bilateral vocal cord paralysis (BVCP) in regard to dyspnea severity, voice quality and aspiration.

Ethics approval required
Old ethics approval format

Ethics approval(s)
Zagazig University Ethical Committee had approved this study and written formal consents had been signed by patients or their relatives

Study design
Prospective randomized controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Optimum method of cordotomy

Interventions

Participants will be randomly categorized into two groups:

Group (A) will be treated with laser assisted posterior cordotomy

Group (B) will be treated with diathermy assisted posterior cordotomy

During and at the end of the study, dyspnea severity, voice quality and aspiration will be assessed and compared between both groups.

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

1. Dyspnea severity scale
2. Voice assessment protocol
3. Pearson aspiration scale

Key secondary outcome(s)

No secondary outcome measures

Completion date

01/02/2011

Eligibility

Key inclusion criteria

Patients [male and female, any age (no age limitation)] who are diagnosed as having bilateral abductor vocal cord paralysis for one year at least and maximal width of respiratory chink is 4 mm or less.

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Key exclusion criteria

1. Patients have unilateral vocal cord paralysis.
2. Patients who are diagnosed as having bilateral abductor vocal cord paralysis for less than one year duration.
3. Maximal width of respiratory chink is more than 4 mm.

Date of first enrolment

01/02/2008

Date of final enrolment

01/02/2011

Locations

Countries of recruitment

Egypt

Study participating centre

Otorhinolaryngology, Head and Neck Surgery Department

Zagazig

Egypt

44519

Sponsor information

Organisation

Zagazig University (Egypt)

ROR

<https://ror.org/053g6we49>

Funder(s)

Funder type

University/education

Funder Name

Zagazig University (Egypt) - Otorhinolaryngology, Head and Neck Surgery Department

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/09/2013		Yes	No