

Incidence and outcome of perioperative renal failure

Submission date 22/05/2013	Recruitment status No longer recruiting	☐ Prospectively registered
		☐ Protocol
Registration date 14/06/2013	Overall study status Completed	☐ Statistical analysis plan
		☐ Results
Last Edited 14/06/2013	Condition category Urological and Genital Diseases	☐ Individual participant data
		☐ Record updated in last year

Plain English summary of protocol

Background and study aims

Though acute kidney injury (AKI) [kidneys suddenly (acute) become unable to do the work expected of them] is extremely common, the knowledge about this disease in the perioperative phase (at or around the time of operation) is very poor. This study aims to understand the interfering factors and predictors of renal function in the perioperative setting (within 48 h postoperatively) after moderate to high-risk surgery in both patients with normal or already preoperatively compromised renal function. Independent predictor for acute kidney injury (AKI) and incidence of AKI in the perioperative setting will be determined. Finally, different types of surgery will be assessed and the relationship between occurrence of AKI and the type of surgery will be assessed. The findings of this study will be used in the improvement and quality program in University Hospital Brussels.

Who can participate?

All patients who underwent a moderate to high-risk surgery in one year will be included in this study.

What does the study involve?

Preoperative and postoperative renal function data, mortality after 30 days after surgery and length of stay in the hospital will be assessed. The association between preoperative patient characteristics and AKI will be assessed. Independent risk factors to develop AKI and on mortality will be determined. Finally, those patients with chronic kidney disease will be analysed separately to determine independent predictors to develop worsening kidney function.

What are the possible benefits and risks of participating?

As this is an observational study, there is no risk.

Where is the study run from?

The study is run from the University Hospital, department of Anesthesiology and Perioperative Medicine in Brussels.

When is the study starting and how long is it expected to run for?

The study started in June 2013 and is expected to last till September 2013.

Who is funding the study?
University Hospital Brussels (Belgium) - Department of Anaesthesiology and Perioperative Medicine

Who is the main contact?
Jan Poelaert, MD, PhD
University Hospital Brussels

Contact information

Type(s)
Scientific

Contact name
Prof Jan Poelaert

Contact details
UZ Brussel
Laarbeeklaan 101
Brussels
Belgium
1090

Additional identifiers

Study information

Scientific Title
Incidence and outcome of perioperative renal failure after moderate and severe surgery

Study objectives
Retrospectively, incidence of perioperative renal failure in moderate or high risk surgical procedures will be assessed, in addition in those patients with diminished renal function, postoperative renal failure will noted and registered.

Ethics approval required
Old ethics approval format

Ethics approval(s)
Ethics Committee University Hospital Brussels, Approval date: 22/12/2011, ref: 2011/282

Primary study design
Observational

Study design
Retrospective observational study

Study type(s)
Other

Health condition(s) or problem(s) studied

Renal failure, postoperative outcome

Interventions

The observational study concerns the occurrence rate of renal failure preoperatively and postoperatively. The purpose is to come to a risk score in our hospital. Therefore we need a large population (we deliberately took one year of patients) and variables such as preoperative status (cardiac, respiratory, renal) as well as the same variables on postoperative day (POD) 1 and 30 day mortality and length of stay in the hospital.

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

30 day mortality

Key secondary outcome(s)

1. Renal function at discharge. The renal function will be assessed according to the RIFLE criteria:
Risk: increase of the serum creatinine x1.5GFR or drop down >25%
Injury: increase of the serum creatinine x2GFR or drop down >50%
Failure: increase of the serum creatinine x3 GFR or drop down >75%
Loss: total loss of kidney function for more than 4 weeks
ESDR: complete loss of kidney function, need for dialysis
2. Need for postoperative supportive measures

Completion date

01/09/2013

Eligibility**Key inclusion criteria**

All patients ≥ 18 years male/female, who have had in 2010/2011 a moderate to high risk surgery

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 Years

Sex

All

Key exclusion criteria

1. Absence of moderate or high risk surgery
2. < 18 years and kidney transplant recipients
3. Surgery under local anesthesia
4. Patients in which no serum creatinine determination pre-and postoperative was done
5. Patients who already have preoperative end-stage renal disease (ESRD) according to the RIFLE criteria

Date of first enrolment

01/06/2013

Date of final enrolment

01/09/2013

Locations**Countries of recruitment**

Belgium

Study participating centre

UZ Brussel

Brussels

Belgium

1090

Sponsor information**Organisation**

University Hospital Brussels (Universitair Ziekenhuis Brussel [UZ Brussel]) (Belgium)

ROR

<https://ror.org/038f7y939>

Funder(s)**Funder type**

Hospital/treatment centre

Funder Name

University Hospital Brussels (Belgium) - Department of Anaesthesiology and Perioperative Medicine

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration