

The possible beneficial role of the regular use of potent mouthwash solutions in the treatment of COVID-19

Submission date 03/10/2020	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 05/10/2020	Overall study status Completed	<input type="checkbox"/> Protocol
Last Edited 16/12/2022	Condition category Infections and Infestations	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

COVID-19 is a condition caused by the coronavirus (called SARS-CoV-2) that was first identified in late 2019. This virus can infect the respiratory (breathing) system. Some people do not have symptoms but can carry the virus and pass it on to others. People who have developed the condition may develop a fever and/or a continuous cough among other symptoms. This can develop into pneumonia. Pneumonia is a chest infection where the small air pockets of the lungs, called alveoli, fill with liquid and make it more difficult to breathe.

In 2020, the virus has spread to many countries around the world and neither a vaccine against the virus or specific treatment for COVID-19 has yet been developed. As of March 2020, it is advised that people minimize travel and social contact, and regularly wash their hands to reduce the spread of the virus.

Groups who are at a higher risk from infection with the virus, and therefore of developing COVID-19, include people aged over 70 years, people who have long-term health conditions (such as asthma or diabetes), people who have a weakened immune system and people who are pregnant. People in these groups, and people who might come into contact with them, can reduce this risk by following the up-to-date advice to reduce the spread of the virus.

Some over-the-counter mouthwash solutions have been shown to kill bacteria and viruses in laboratory studies. This study aims to test this effect in people with COVID-19.

Who can participate?

Patients aged 18 and older who have tested positive for COVID-19

What does the study involve?

Participants will be randomly allocated to receive a strong mouthwash solution, to be used 3 times a day for 2 weeks, or treatment as usual.

What are the possible benefits and risks of participating?

Possible benefits include faster recovery and fewer secondary bacterial infections in the upper

respiratory tract. Possible risks include mouth irritation, especially if there are mouth ulcers or sensitivity to the components, as well as digestive system irritation if hydrogen peroxide is ingested in quantities larger than the total daily recommended dose.

Where is the study run from?

1. Hazm Mebaireek General Hospital (Qatar)
2. Communicable Disease Center - Hamad Medical Corporation (Qatar)

When is the study starting and how long is it expected to run for?

April 2020 to November 2020

Who is funding the study?

Hamad Medical Corporation (Qatar)

Who is the main contact?

Dr Khalid Mukhtar
kmukhter@hamad.qa

Contact information

Type(s)

Scientific

Contact name

Dr Khalid Mukhtar

ORCID ID

<https://orcid.org/0000-0002-9261-731X>

Contact details

Medical Research Center
Hamad Medical City
Doha Industrial Area (west)
Doha
Qatar

-

+974 (0)77446394
kmukhter@hamad.qa

Additional identifiers

Protocol serial number

MRC 05-106

Study information

Scientific Title

Effect of regular use of hydrogen peroxide and chlorhexidine gluconate mixed solution for mouth wash and gargles on the COVID-19 recovery rate: a randomized controlled trial

Study objectives

The development and progression of the disease following exposure to SARS-CoV-2 is affected by the microbial communities in the oral cavity; proposing a "complicit agent" hypothesis, either a single microbial agent effect (presence of complicit or absence of a protective) or the state of dysbiosis in general. Meanwhile, game theory and evolutionary biology suggests that inhibiting cooperation - reciprocal altruism - between two organisms can negatively affect their survival. The researchers propose that regular use of potent mouthwash can improve the disease course, as the strain of the continual reduction in the microbial load is likely to inhibit their cooperation.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 20/08/2020, Hamad Medical Corporations' Medical Research Center IRB (Medical Research Center, Hamad Medical City, Doha, Qatar; +974 (0)5554 6316; IRB@hamad.qa), ref: MRC 05-106

Study design

Interventional randomized parallel trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

COVID-19 (SARS-CoV-2 infection)

Interventions

New admissions (24 hrs) will be screened for eligibility. Recruited subjects will be randomly allocated to either the study or control group.

Study group: Potent disinfectant solution (10 ml Chlorhexidine gluconate 0.2% + 5 ml of hydrogen peroxide 6%) will be provided three times daily for 2 weeks, to be used for mouth rinse and gargles (>30 sec) and to avoid eating or drinking for at least 5 more minutes

Control group: treatment as usual

Intervention Type

Other

Primary outcome(s)

1. Recovery assessed using clinical improvement along with a negative COVID RT-PCR test at 5 and 15 days of starting treatment
2. Changes in the CT-values of COVID-19 RT-PCR inconclusive test at 5 and 15 days of starting treatment

Key secondary outcome(s)

1. Clinical symptoms assessed using the Sore Throat Assessment Tool for the first 5 days
2. COVID progression (ICU intubation vs discharge or transfer to quarantine) recorded by

reviewing medical records within the study duration

3. Hospital stay measured using (days) at discharge; or at the end of study for those remained admitted

4. Disposition type (discharged/death) measured using hospital records entries at the time of discharge, or reported mortality during hospital stay

Completion date

20/11/2020

Eligibility

Key inclusion criteria

1. Age 18 years and older
2. COVID-19 reported via positive PCR within 2 weeks of admission
3. Conscious, oriented and can comprehend the study purpose and risks

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Total final enrolment

100

Key exclusion criteria

1. Pregnancy (females)
2. Intubated (at the time of recruitment) or had any reported cognitive impairment that can prevent proper comprehension of the study or communicating side effects
3. Contra-indication for mouth wash use, e.g. reported allergies to the solution constituents, recent facial/head injuries, maxillofacial conditions

Date of first enrolment

08/09/2020

Date of final enrolment

01/10/2020

Locations

Countries of recruitment

Qatar

Study participating centre

Hazm Mebaireek General Hospital

Hamad Medical Corporation

Doha Industrial Area (west)

Doha

Qatar

-

Study participating centre

Communicable Disease Center - Hamad Medical Corporation

Hamad Medical City

Doha

Qatar

-

Sponsor information

Organisation

Hamad Medical Corporation

ROR

<https://ror.org/02zwb6n98>

Funder(s)

Funder type

Industry

Funder Name

Hamad Medical Corporation

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

Kmukhter@gmail.com

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Basic results		04/05/2021	05/05/2021	No	No
Preprint results	non-peer-reviewed results in preprint	30/11/2020	16/03/2021	No	No