

Study on schizophrenia patients responsive to fish oil supplementation therapy

Submission date 19/12/2023	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
Registration date 29/12/2023	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
Last Edited 19/01/2026	Condition category Mental and Behavioural Disorders	<input type="checkbox"/> Individual participant data
		<input checked="" type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Schizophrenia is a complex disorder that presents long-term challenges such as a lack of objective diagnostic markers and complex and varying clinical symptoms. The blunted niacin skin flushing response is commonly observed in patients with schizophrenia and is considered one of the objective diagnostic markers for the disorder. It is closely related to the membrane lipid disorder hypothesis of schizophrenia, which suggests that schizophrenia subgroups selected based on blunted niacin response are believed to be caused by membrane lipid disorders. Factors leading to membrane lipid disorders in the body include lipid peroxidation caused by oxidative stress and inflammatory responses. Fish oil is a common antioxidant and anti-inflammatory supplement, composed mainly of n-3 polyunsaturated fatty acids, which are important components of membrane lipids. In this study, we randomly assigned participants to groups and added fish oil supplementation to the conventional treatment in the experimental group. The aim was to identify the schizophrenia population that responds well to this supplementary treatment of fish oil, with the expectation of providing guiding suggestions for clinical treatment.

Who can participate?

Patients aged between 18 and 65 years with schizophrenia who are registered at Renmin Hospital of Wuhan University, and eligible healthy volunteers

What does the study involve?

The patients were randomly recruited in the hospital outpatient department and were divided into into a control group and an experimental group. The control group received conventional treatment, while the experimental group was given additional fish oil supplementation. The supplemental dose of fish oil (Puritan's Pride OMEGA-3 Fish Oil, USA) is 1800-2700 mg per day. Niacin response were administered to all subjects before treatment (baseline) by trained psychiatrists, and all patients were assessed with the Positive and Negative Syndrome Scale (PANSS). At follow-up, niacin response and PANSS scales were assessed in fish oil supplementation patients.

What are the possible benefits and risks of participating?

All participants have access to a clinical evaluation by the professional psychiatrists, which is free

of charge. And participants who receive the PUFA may benefit from a reduction of their psychiatric symptoms. There are no known risks involved with participating.

Where is the study run from?

Renmin Hospital of Wuhan University (China)

When is the study starting and how long is it expected to run for?

January 2019 to December 2023

Who is funding the study?

National Natural Science Foundation of China

Who is the main contact?

1. Shijing Wang, wdrmiit@163.com

2. Zhiyan Gu, gzy981114@163.com

Contact information

Type(s)

Public, Scientific, Principal investigator

Contact name

Mr Shijing Wang

Contact details

238 Jiefang Road, Renmin Hospital of Wuhan University

Wuhan

China

430000

+86 (0)18077408842

wdrmiit@163.com

Additional identifiers

Study information

Scientific Title

Study on efficacy evaluation of fish oil supplementation therapy for schizophrenia based on niacin skin flushing responses

Study objectives

Current study objectives as of 02/12/2025:

Blunted niacin response is widely shown in patients with schizophrenia, which has become an objective auxiliary diagnostic marker for schizophrenia and is closely related to the pathological hypothesis of abnormal membrane phospholipids in schizophrenia. The main component of fish oil is n-3 polyunsaturated fatty acid, which plays an important physiological role in regulating membrane phospholipids. This study aims to identify schizophrenia patients with impaired

nicotinic acid response and provide fish oil supplementation as a targeted intervention, ultimately identifying the population that responds to fish oil supplementation, reducing trial-and-error costs, and improving treatment efficiency.

Previous study objectives:

Blunted niacin response is widely shown in patients with schizophrenia, which has become an objective auxiliary diagnostic marker for schizophrenia and is closely related to the pathological hypothesis of abnormal membrane phospholipids in schizophrenia. The main component of fish oil is n-3 polyunsaturated fatty acid, which plays an important physiological role in regulating membrane phospholipids. This study aims to identify schizophrenia patients with impaired niacin response and administer fish oil supplementation as a targeted intervention, ultimately shedding light on the potential of niacin flush response as a guide for clinical schizophrenia treatment and prognostic indicators.

Ethics approval required

Ethics approval required

Ethics approval(s)

approved 24/05/2019, Ethics Committee of Bio-X Institutes at Shanghai Jiao Tong University (Shanghai, Huashan Road, Shanghai, 200030, China; +86 (0)15921495069; wangdandan26@126.com), ref: ML2019041

Study design

Interventional open-label randomized controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Schizophrenia

Interventions

Current interventions as of 02/12/2025:

Patients with mental disorders were randomly allocated to the experimental group and the control group. The control group received routine treatment, whereas the experimental group received adjuvant fish oil supplementation in addition to routine treatment, with a total treatment course of 21 days. The method of randomisation is according to a computer-generated random sequence using block randomisation with random block sizes. The randomisation is performed by the independent statistician. Patients are enrolled and assigned sequentially to adjuvant interventions by the physician. The allocation sequence is not available to any member of the research team until the databases had been completed and locked.

The supplemental dose of fish oil (Puritan's Pride OMEGA-3 Fish Oil, USA) is 1800-2700 mg per day.

All patients were assessed with the Positive and Negative Syndrome Scale (PANSS). At follow-up, niacin response and PANSS scales were assessed in fish oil supplementation patients.

Previous interventions:

Recruited patients are tested for niacin skin flushing responses and divided into blunted and normal niacin response groups according to their niacin skin flushing responses. Subsequently, patients with blunted niacin responses are randomly assigned to receive fish oil supplementation for 3 months. Those not randomly assigned were not given fish oil supplementation. The method of randomisation is according to a computer-generated random sequence using block randomisation with random block sizes. The randomisation is performed by the independent statistician. Patients are enrolled and assigned sequentially to adjuvant interventions by the physician. The allocation sequence is not available to any member of the research team until the databases had been completed and locked.

The supplemental dose of fish oil (Puritan's Pride OMEGA-3 Fish Oil, USA) is 1800-2700 mg per day.

Niacin response and Brief Assessment of Cognition in Schizophrenia (BACS) were administered to all subjects before treatment (baseline) by trained psychiatrists, and all patients were assessed with the Positive and Negative Syndrome Scale (PANSS). At follow-up, niacin response, BACS and PANSS scales were assessed in fish oil supplementation patients.

Intervention Type

Supplement

Primary outcome(s)

Psychotic symptoms measured using the Positive and Negative Syndrome Scale were measured at baseline and month 3

Key secondary outcome(s)

Current key secondary outcome(s) as of 02/12/2025:

1. Niacin skin flushing response were measured at baseline and month 3
-

Previous key secondary outcome(s):

1. Niacin skin flushing response were measured at baseline and month 3
2. Cognition measured using the Brief Assessment of Cognition in Schizophrenia were measured at baseline and month 3

Completion date

31/12/2023

Eligibility

Key inclusion criteria

1. Patients with schizophrenia diagnosed according to Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5) criteria
2. Age between 18 and 65 years
3. Commitment to comply with the study procedures and cooperate with the implementation of the entire research process

Participant type(s)

Healthy volunteer, Patient

Healthy volunteers allowed

No

Age group

Mixed

Lower age limit

18 years

Upper age limit

65 years

Sex

All

Total final enrolment

129

Key exclusion criteria

1. Severe neurological diseases or traumatic brain injury
2. Substance dependence
3. Administration of nonsteroidal or steroidal anti-inflammatory drugs within 2 weeks
4. Having a history of severe allergies, currently suffering from skin diseases or immune system diseases
5. Pregnancy

Added 19/01/2026:

6. Allergic to adhesive tape

Date of first enrolment

01/06/2019

Date of final enrolment

31/12/2022

Locations**Countries of recruitment**

China

Study participating centre
Renmin Hospital of Wuhan University
238 Jiefang Road
Wuhan
China
430000

Sponsor information

Organisation
Renmin Hospital of Wuhan University

ROR
<https://ror.org/03ekhbz91>

Funder(s)

Funder type
Research organisation

Funder Name
National Natural Science Foundation of China

Alternative Name(s)
Chinese National Science Foundation, Natural Science Foundation of China, National Science Foundation of China, NNSF of China, NSF of China, National Nature Science Foundation of China, Guójiā Zìrán Kēxué Jījīn Wěiyuánhùi, , NSFC, NNSF, NNSFC

Funding Body Type
Government organisation

Funding Body Subtype
National government

Location
China

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study will be available upon request from Shijing Wang (wdrmiit@163.com). The data will be available beginning 3 months

and ending 5 years following article publication. It can be shared with researchers who provide a methodologically sound proposal to achieve the aims in the approved proposal. To gain access, data requesters will need to sign a data access agreement. Consent from participants was obtained. Data will be de-identified by removing personally identifiable information.

IPD sharing plan summary

Available on request