

# Is Hedrin® 4% lotion Dev 42 more effective than Hedrin® 4% lotion against head louse eggs?

<b>Submission date</b> 09/03/2009	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered
<b>Registration date</b> 19/03/2009	<b>Overall study status</b> Completed	<input type="checkbox"/> Protocol
<b>Last Edited</b> 30/03/2020	<b>Condition category</b> Infections and Infestations	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

### Background and study aims

We know from previous studies that Hedrin 4% lotion gets rid of a head louse infestation. The aim of this study is to compare Hedrin 4% lotion with a modified version of the product to see if the alternative is more effective to cure head louse infestation.

### Who can participate?

The study will be conducted in two schools but is open to anyone in the villages serving those schools who has head lice and is over the age of 6 months and who fits the other entry criteria.

### What does the study involve?

The study first involves checking for presence of lice. If you then wish to take part we shall ask you (or your carer) to sign a consent form after which we can apply the treatment. A second application of the treatment product is applied after 7 days to eliminate any young lice that emerge from eggs during that first week. We shall check the safety and acceptability of the treatment and the progress of the effectiveness when we comb everyone looking for lice twice between treatments, on days 1, 2 and 6 after first treatment, and then twice after the second treatment on days 9 and 14. Anyone who has lice after the second treatment will be provided with a further treatment (Hedrin 4% lotion) to eliminate the infestation.

### What are the possible benefits and risks of participating?

We hope everyone who takes part will have their head lice eliminated.

### Where is the study run from?

The study will be conducted in the schools of Maldan and Osmancali, two villages in Manisa province, Turkey. Our team of investigators will first visit the schools to identify students with lice and then visit them at home to run all the procedures.

### When is the study starting and how long is it expected to run for?

The study was carried out in April and May of 2009.

Who is funding the study?

It is being funded by Thornton & Ross Ltd, the makers of Hedrin 4% lotion.

Who is the main contact?

The main contact for the study is Dr Özgür Kurt of Celal Bayar University School of Medicine.

## Contact information

### Type(s)

Scientific

### Contact name

Mr Ian Burgess

### Contact details

Medical Entomology Centre  
Insect Research & Development Limited  
6 Quay Court  
Colliers Lane  
Stow-cum-Quy  
Cambridge  
United Kingdom  
CB25 9AU  
+44 (0)1223 810070  
ian@insectresearch.com

## Additional identifiers

### Protocol serial number

CTMK06

## Study information

### Scientific Title

A randomised, controlled, assessor-blind, clinical trial to compare Hedrin® 4% dimeticone lotion with Hedrin® 4% lotion Dev 42 in the treatment of head lice

### Study objectives

To compare efficacy of Hedrin® 4% lotion with Hedrin® 4% lotion Dev 42 in the eradication of head lice, to compare the products for ovicidal activity, and to compare the products for safety, ease of application and participant acceptability.

### Ethics approval required

Old ethics approval format

### Ethics approval(s)

Research Ethics Committee of the Medical Faculty, Celal Bayar University gave approval on the 25th February 2008 (ref: 25)

### Primary study design

Interventional

### **Study design**

Randomised controlled assessor blinded parallel group study

### **Study type(s)**

Treatment

### **Health condition(s) or problem(s) studied**

Head louse infestation

### **Interventions**

1. Dimeticone 4% lotion (Hedrin® 4% lotion)
2. Dimeticone 4% lotion, alternative mixture (Hedrin® 4% lotion Dev 42)

Both products will be applied directly to dry hair. Sufficient product will be applied to saturate the hair and scalp. The product will be left in place for eight hours (or overnight) before being shampooed and rinsed off with water. Treatments were applied on days 0 and 7 and post-treatment assessments were conducted on days 1, 2, 6, 9, 14.

### **Intervention Type**

Drug

### **Phase**

Phase III

### **Drug/device/biological/vaccine name(s)**

Hedrin® 4% lotion Dev 42, Hedrin® 4% lotion

### **Primary outcome(s)**

1. To compare Hedrin® 4% lotion with Hedrin® 4% lotion Dev 42 with regard to inhibition of hatching of louse eggs (ovicidal activity): no nymphal lice emerging from eggs between treatments (i.e. on day 1, 2, or 6) or after the second treatment
2. To compare the efficacy of Hedrin® 4% lotion with Hedrin® 4% lotion Dev 42 with regard to the eradication of head lice: assessed post-second treatment, i.e. no lice on days 9 or 14

### **Key secondary outcome(s)**

To compare Hedrin® 4% lotion with Hedrin® 4% lotion Dev 42 with regard to safety, ease of application and participant acceptability, assessments made on days 0 and 7 post-treatment application and at each of the efficacy assessments on days 1, 2, 6, 9, and 14

### **Completion date**

31/05/2008

## **Eligibility**

### **Key inclusion criteria**

1. Both males and females, aged 4 years and over with no upper age limit
2. Participants who upon examination, are confirmed to have live head lice
3. Participants who give written informed consent, or if the participant is under 16 years of age

whose parent/guardian gives written informed consent to participate in the study  
4. Participants who will be available for follow-up visits by study team members over the 14 days following first treatment

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Other

**Sex**

All

**Total final enrolment**

72

**Key exclusion criteria**

1. Participants with a known sensitivity to any of the ingredients in Hedrin® 4% lotion or Hedrin® 4% lotion Dev 42
2. Participants with a secondary bacterial infection of the scalp (e.g. impetigo) or who have an active long-term scalp condition (e.g. psoriasis of the scalp)
3. Participants who have been treated with other head lice products within the previous two weeks
4. Participants who have bleached hair, or hair that has been permanently waved within the previous four weeks
5. Participants who have been treated with the antibiotics co-trimoxazole or trimethoprim within the previous four weeks, or who are currently taking such a course
6. Pregnant or nursing mothers
7. Participants who have participated in another clinical study within 1 month before entry to this study
8. Participants who have already participated in this clinical study

**Date of first enrolment**

01/04/2008

**Date of final enrolment**

31/05/2008

**Locations****Countries of recruitment**

United Kingdom

England

Türkiye

**Study participating centre**  
**Medical Entomology Centre**  
Cambridge  
United Kingdom  
CB25 9AU

## Sponsor information

### Organisation

Thornton & Ross Ltd (UK)

### ROR

<https://ror.org/00frd0c49>

## Funder(s)

### Funder type

Industry

### Funder Name

Thornton & Ross Ltd (UK)

## Results and Publications

### Individual participant data (IPD) sharing plan

### IPD sharing plan summary

Not provided at time of registration

### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>	results	01/12/2009	30/03/2020	Yes	No