

Correcting non cephalic presentation with moxibustion: Study protocol for a multi-centre randomised controlled trial in general practice

Submission date 02/04/2008	Recruitment status No longer recruiting	☐ Prospectively registered
Registration date 30/04/2008	Overall study status Completed	☐ Protocol
Last Edited 14/08/2013	Condition category Pregnancy and Childbirth	☐ Statistical analysis plan
		☒ Results
		☐ Individual participant data

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Jorge Vas

Contact details

Centro de Salud
Unidad de Tratamiento del Dolor
C/ Segovia s/n
Dos Hermanas
Spain
41700
jorgef.vas.sspa@juntadeandalucia.es

Additional identifiers

Protocol serial number

0053/2007

Study information

Scientific Title

Study objectives

A sensorial stimulation with moxibustion in the outer corner of the little toenail (BL67) produce a greater proportion of version in non-cephalic presentations than the usual care.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Clinical Trials Ethics Committee of the Andalusian Regional Government. Date of approval: 26/02/2008 (ref: acta 02/08)

Primary study design

Interventional

Study design

Multi-centre randomised controlled trial with three parallel arms.

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Non cephalic presentation of fetus

Interventions

All the study subjects will be given various postural recommendations.

The participants will be randomly allocated to the following three arms:

Arm A: True moxibustion (TM): The first intervention will be made at the Healthcare Clinic. This session (overseen by the responsible midwife) will serve as a training session for the friend or relative who is to apply the remaining sessions at home. The mother should lie down, face up, wearing comfortable clothing and with no pressure at the waist from buttons, belt, etc. and with her legs slightly bent at the knee. Heat will be applied for 20 minutes, by means of a moxa cone (a herbal preparation with *Artemisia vulgaris*) at point BL67 (Zhiyin), close to the outer angle of the little toenail. The heat will be applied from a distance of 1.5-3 cm, to ensure that the subject notices a feeling of warmth in the zone, but without any pain or burning. The application site (right or left foot) will be alternated every 2 minutes, or more often if any untoward sensation of heat is felt. True moxibustion will be carried out once a day for two weeks, or until cephalic version occurs.

Arm B: False/sham moxibustion (SM): This procedure will be identical to that applied to the TM group except that the point stimulated will be a non-active one according to the principles of Traditional Chinese Medicine, point SP1 (Yinbai), close to the inner angle of the big toenail. False moxibustion will be carried out once a day for two weeks, or until cephalic version occurs.

Arm C: Control group, under observation (CG): The subjects in this group will receive the standard treatment, in accordance with the Andalusian Healthcare System Treatment Guide.

Intervention Type

Drug

Phase

Not Specified

Drug/device/biological/vaccine name(s)

Moxibustion

Primary outcome(s)

Rate of cephalic presentation at term.

Key secondary outcome(s)

1. Rate of cephalic presentation at week 38 of gestation (CP38), determined by echography
 2. Number of days of treatment received until version occurs
 3. Rate of caesarean sections
 4. Security variables:
 - 4.1. Increased rates of foetal cardiac rhythm change, followed-up until birth
 - 4.2. Risk of premature birth
 - 4.3. Changes in Apgar scores at 1 minute and at 5 minutes
1. Rate of cephalic presentation at week 38 of gestation (CP38), determined by echography
 2. Number of days of treatment received until version occurs
 3. Rate of caesarean sections
 4. Security variables:
 - 4.1. Increased rates of foetal cardiac rhythm change
 - 4.2. Risk of premature birth
 - 4.3. Changes in Apgar scores at 1 minute and at 5 minutes

Completion date

01/12/2010

Eligibility**Key inclusion criteria**

1. Pregnant women
2. Aged at least 18 years
3. Foetus is in a non cephalic position (diagnosed by physical examination and by echography)
4. Gestational age of 33-35 weeks (estimated by echography)
5. Normal foetal biometry
6. Those who have signed their informed consent
7. Have not previously received moxibustion treatment to correct the position of their unborn baby

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 Years

Sex

Female

Key exclusion criteria

1. Twin pregnancy
2. Pelvic defect
3. Previous uterine surgery
4. Foetal malformation or chromosome disorder
5. Uterine malformation
6. Risk of premature birth (preterm uterine contractions and/or initial dilation or shortening of the cervix with a Bishop's score of 4)
7. Uterine fibromas >4 cm
8. Treatment with tocolytic drugs
9. Heart or kidney disease affecting the mother
10. Other pregnancy-related complications
11. Inability to complete the questionnaire or respond to the assessor's questions

Date of first enrolment

01/04/2008

Date of final enrolment

01/12/2010

Locations

Countries of recruitment

Spain

Study participating centre

Centro de Salud

Dos Hermanas

Spain

41700

Sponsor information

Organisation

Andalusian Regional Ministry of Health (Spain)

ROR

<https://ror.org/03q4c3e69>

Funder(s)

Funder type

Government

Funder Name

Andalusian Regional Ministry of Health (Project No. 0053/2007)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/03/2013		Yes	No