

Beneficial effects of thoracic epidural anaesthesia on mortality after surgery for colon cancer

Submission date 25/05/2019	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
Registration date 07/06/2019	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
Last Edited 29/08/2019	Condition category Surgery	<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Perioperative use of epidural analgesia reduces the 36-month postoperative mortality rate. This effect may be due to systemic effects of local anaesthetics or to a reduced stress response caused by the thoracic epidural analgesia itself.

Who can participate?

Patients scheduled to undergo elective open colon cancer surgery.

What does the study involve?

Patients records are followed up over a 4 year period following surgery.

What are the possible benefits and risks of participating?

None.

Where is the study run from?

Marienhospital Vechta gGmbH, Germany.

When is the study starting and how long is it expected to run for?

January 1997 to December 2015

Who is funding the study?

The study is funded by the investigator.

Who is the main contact?

Dr Christian Hönemann,
c.honemann@icloud.com

Contact information

Type(s)

Scientific

Contact name

Dr Christian Hönemann

ORCID ID

<https://orcid.org/0000-0002-9658-253X>

Contact details

arienstraße 6-8

Vechta

Germany

49377

+4915208961106

christian.hoenemann@kh-vec.de

Additional identifiers

Protocol serial number

MarienVEC_12

Study information

Scientific Title

Epidural anaesthesia reduces mortality after colon cancer surgery

Acronym

EARMAC

Study objectives

Epidural anaesthesia reduces mortality after colon cancer surgery

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved DATE, Ethikkommission der Aerztekammer Niedersachsen (Ethical Committee of the Aerztekammer Niedersachsen, ÄKN Ethikkommission, Berliner Allee 20, 30175 Hannover, Germany; 05113802208; ethikkommission@aekn.de), ref:

Study design

Observational study

Primary study design

Observational

Study type(s)

Other

Health condition(s) or problem(s) studied

Patients with elective colon cancer surgery

Interventions

Patient records were used to follow-up on outcomes over 4 years for patients who had undergone elective colon cancer surgery with or without thoracic epidural anesthesia/analgesia.

Intervention Type

Other

Primary outcome(s)

Mortality, measured by review of patient charts and a database of a local cancer database in Oldenburg, Germany

Key secondary outcome(s)

1. Perioperative complications
2. Stroke
3. Myocardial infarction
4. Blood transfusion
5. Duration of stay

All measured by review of patient charts and a database of a local cancer database in Oldenburg, Germany

Completion date

31/12/2015

Eligibility

Key inclusion criteria

Scheduled to undergo elective open colon cancer surgery

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Key exclusion criteria

1. Non elective surgery
2. Laparoscopic surgery

Date of first enrolment

01/01/1997

Date of final enrolment

31/12/2007

Locations

Countries of recruitment

Germany

Study participating centre

Marienhospital Vechta gGmbH

Marienstraße 6-8

Vechta

Germany

49377

Sponsor information

Organisation

Marienhospital Vechta gGmbH

Funder(s)

Funder type

Other

Funder Name

Investigator initiated and funded

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request

IPD sharing plan summary

Available on request