

# The effect of magnesium sulfate on recurrence of rebound pain in patients undergoing regional anesthesia in the arm

<b>Submission date</b> 25/11/2022	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 22/12/2022	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
<b>Last Edited</b> 20/12/2022	<b>Condition category</b> Surgery	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

Peripheral nerve blocks provide effective analgesia for long durations, and lower opioid intake, nausea and vomiting, resulting in faster hospital discharge. But some patients might experience intense pain after the effects of regional anesthesia, known as rebound pain. There are medicines when mixed with local anesthetics can provide better results, such as dexmedetomidine, dexamethasone, and magnesium sulfate. The study aims to evaluate the rebound pain with and without magnesium sulfate as an adjuvant with ropivacaine in supraclavicular brachial plexus nerve block in upper limb surgeries. Our hypothesis is that magnesium sulfate as a perineural adjuvant could reduce the incidence of rebound pain, improving patients postoperative pain control.

### Who can participate?

Adults over the age of 18, submitted to surgeries of the distal arm, forearm, and hand, eligible to supraclavicular brachial plexus nerve block and sedation, in Mario Covas State Hospital in Santo Andre, Sao Paulo.

### What does the study involve?

Participants are asked to join this study while they wait in the room for nursing triage before their surgery. Participants must understand the anesthesia process and the diary of pain they will complete in the following 24h after the surgery. Participants are randomly allocated to one of two groups. Those in the first group will receive an adjuvant mixed with the local anesthetic. Those in the second group will receive local anesthetic only. Both groups will fill in scores of pain on the diary of pain in the next 24h and note the need for opioids, in the search for better pain scores.

### What are the possible benefits and risks of participating?

The possible benefit to those taking part is lower pain scores after surgery and less opioid intake. The main risk is block failure, others like pneumothorax, blood bruises, and nerve damage are also in consideration, all of them diminished by the usage of ultrasonography-guided puncture.

Where is the study run from?

The study is being run by the ABC Foundation and takes place in Mario Covas State Hospital in Santo Andre, Sao Paulo (Brazil)

When is the study starting and how long is it expected to run for?

April 2021 to December 2023

Who is funding the study?

Investigator initiated and funded

Who is the main contact?

1. Marcelo Marcolla

marcelamarcolla@hotmail.com

2. Victor Peres de Melo Goulart

victor@roffor.com.br

## Contact information

### Type(s)

Public

### Contact name

Mr Victor Goulart

### ORCID ID

<https://orcid.org/0000-0002-8770-8014>

### Contact details

Rua cananeia, 138

apto 71

Santo Andre

Brazil

09060480

+55 92984151064

victor@roffor.com.br

## Additional identifiers

## Study information

### Scientific Title

The impact of perineural magnesium sulfate on rebound pain after supraclavicular brachial plexus block - randomized clinical trial

### Study objectives

The use of magnesium sulfate as a perineural adjuvant could reduce the incidence of rebound pain, improving postoperative pain control in patients undergoing upper limb surgery.

### Ethics approval required

Old ethics approval format

### **Ethics approval(s)**

Approved 15/08/2022, College of Medicine of ABC Research Ethics Committee (2000 Lauro Gomes avenue, Santo André, SP; +55(11)49935453; cep@fmabc.br), ref: 5.582.819.

### **Study design**

Interventional double-blinded randomized parallel trial

### **Primary study design**

Interventional

### **Study type(s)**

Prevention

### **Health condition(s) or problem(s) studied**

Prevention of rebound pain in patients undergoing supraclavicular brachial plexus block for upper limb surgery.

### **Interventions**

The sample size was calculated using STATA/IC v.17 (StataCorp, College Station, TX, USA) and also considering previous studies and potential exclusions from the protocols. The sample is 34, with 17 participants each group.

The participants are randomized to group A (saline) and group B (magnesium sulfate) using Research Electronic Data Capture, REDCap® v. 10.

The participants and researchers are blinded to the intervention.

The solution is prepared by an independent assistant according to the solution preparation guide:

Group A: One syringe of 20mL with 0.5% of ropivacaine.

Preparation: aspirate 10mL of 1% ropivacaine [100mg] and 10mL of 0.9% saline solution with a 20mL syringe.

Group B: One syringe of 20mL with 0.5% ropivacaine and 1% magnesium sulfate.

Preparation: aspirate 10mL of 1% ropivacaine [100mg] with a 20mL syringe. After, add 2mL of 10% magnesium sulfate [200mg] and 8mL of 0.9% saline solution in the same syringe.

The syringes and their contents are indistinguishable for both groups to mask solutions.

The location of the supraclavicular brachial plexus will be done in real time with ultrasound linear probe and with peripheral nerve stimulator connected to needle (Stimuplex® A 30° B Braun 50mm, 22 Gauge short bevel).

The bevel tip of the needle will be positioned towards the angle formed by the first rib and the subclavian artery, called the "corner pocket". Then, an electrical current with an intensity of 0.3 mA, frequency of 2Hz and pulse width of 0.1ms will be activated. After that, the stimulation of the structures innervated by the ulnar nerve is expected. Afterwards, 10mL of anesthetic solution will be administered in the absence of resistance.

Next, the needle will be repositioned until the main neural cluster formed by the superior and middle trunks. In this place, 10mL of the anesthetic solution will be added without resistance.

The sensory and motor block are checked for up to 45 minutes. In case of total or partial failure of anesthesia after that time, the block is considered not viable for the surgical procedure. If there are no contraindications, Ondansetron 8mg, Ketoprofen 100mg, Pantoprazole 40mg and

Dipyrone 2 grams will administered intravenously 40 minutes before the surgical incision for both groups.

At hospital discharge, the participants will responsible for completing the Visual Numeric Scale (VNS) in the pain diary by the first 24 hours after the block. All participants will receive a phone call 24 hours after the block as part of routine care and to collect clinical relevant data.

### **Intervention Type**

Drug

### **Phase**

Not Applicable

### **Drug/device/biological/vaccine name(s)**

Magnesium sulfate, ropivacaine hydrochloride.

### **Primary outcome(s)**

1. Pain measured by the Visual Numerical Scale (VNS) every 6 hours after the block for 24 hours
2. The time of block resolution, considered as the time when the first rescue analgesia was taken. However, in cases that didn't require rescue analgesia, the time of block resolution will be considered when the patient reports the end of numbness or the end of heaviness in the blocked region.

### **Key secondary outcome(s)**

Average cumulative opioid consumption in milligrams between the groups in the first 24 hours after the block, according to the doses taken and checked in the medical prescription.

### **Completion date**

31/12/2023

## **Eligibility**

### **Key inclusion criteria**

1. Patients with ASA Physical Status classification I-III;
2. Ability to understand and complete the pain diary;
3. Patients undergoing arm, forearm and hand surgery with indication for supraclavicular brachial plexus block without the association of another regional block or general anesthesia.

### **Participant type(s)**

Patient

### **Healthy volunteers allowed**

No

### **Age group**

Mixed

### **Sex**

All

### **Key exclusion criteria**

1. Severe heart disease;
2. Decompensated kidney disease;
3. Decompensated liver disease;
4. Severe respiratory failure;
5. Cognitive dysfunction that prevents cooperation with the study;
6. Calcium channel blocker treatment;
7. Confirmed coagulopathy;
8. Infection at the procedure location;
9. Illicit drug used 5 days ago;
10. Daily use of opioids more than 14 days before surgery;
11. Block failure;
12. Need to send to the intensive care unit (ICU) in the immediate postoperative period;
13. Neuropathy in the upper limb to be operated;
14. Allergy to local anesthetic, propofol or non-steroidal anti-inflammatory drugs;
15. Weight <50 Kilograms (kg);
16. Body mass index (BMI) > 35 kg/m<sup>2</sup>;
17. Chronic pain;
18. Pregnancy;
19. Need to change the anesthetic technique intraoperatively.

**Date of first enrolment**

01/10/2022

**Date of final enrolment**

31/10/2023

## Locations

**Countries of recruitment**

Brazil

**Study participating centre**

**Mario Covas State Hospital**

321 Dr Henrique Calderazzo St, Paradise

Santo André

Brazil

09190-615

## Sponsor information

**Organisation**

The Committee for Ethics in Research (CEP) in ABC's Foundation

## Funder(s)

**Funder type**

Other

**Funder Name**

Investigator initiated and funded

## Results and Publications

**Individual participant data (IPD) sharing plan**

The datasets generated and analysed during the current study will be available upon request from Marcela Marçolla Tavares by the e-mail address: marcelamarcolla@hotmail.com.

All the data will become available after the scientific article is published during 5 years.

Data will be stored for 5 years after publication of the results in a scientific journal. Datasets will be available to identified researchers upon request who are interested in scientific research. After shared, they can be analyzed by all types of statistical mechanisms according to the researcher's interest.

The recorded data are anonymous, there are no ethical restrictions and were/will be collected after the consent signed by the research participant.

**IPD sharing plan summary**

Available on request