

Combining exercise with a low-energy diet to minimise muscle losses during weight loss

Submission date 12/12/2023	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 19/12/2023	Overall study status Completed	<input checked="" type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 25/03/2026	Condition category Nutritional, Metabolic, Endocrine	<input type="checkbox"/> Individual participant data <input checked="" type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

The COMBINE Trial investigates the physiological and metabolic impact of adding structured exercise to a 12-week low-energy diet in South Asians living with type 2 diabetes and obesity. Secondary outcomes will report cardiometabolic (insulin sensitivity) and physical function (strength) measures captured through a range of assessment methods.

Who can participate?

South Asian adults aged 40-65 years old with type 2 diabetes and obesity

What does the study involve?

The COMBINE trial will recruit 36 participants to a 12-week low-energy diet with or without supervised structured aerobic and resistance exercise (3 days/week).

The low-energy diet (which is actively used to achieve diabetes remission) will make use of meal replacements and ongoing macronutrient/calorie targets. Meanwhile, the exercise component will involve a mixture of supervised and independent training (encompassing structured aerobic and resistance exercise).

There is no standard control group, all eligible participants will receive the diet intervention, with half of the participants being randomly assigned to also receive exercise support.

What are the possible benefits and risks of participating?

During the study participants will receive close monitoring of their diabetes. The study will also provide detailed results of their blood glucose control, body fat and exercise levels. The researchers will be happy to review the results with them after the study is completed. The diet used in this study can reverse diabetes, being that blood sugar levels may fall outside of diabetes ranges without the need for medication. This happens in just under half of patients who manage to stick to it and lose weight, and this diet is now actively being offered through GPs because of the likelihood of reversing diabetes. Patients often feel much healthier and this tends to occur quickly. This is usually within the first week. Medication is likely to be reduced or stopped altogether. However, not everyone reverses their diabetes. If they do reverse diabetes, whether they are able to maintain this after the study depends on how their body responds over

the longer term and how well they are able to continue the new eating habits. Exercise training improves fitness. It also helps to control blood sugar. Exercise training also lowers blood pressure and the amount of harmful fat in the blood and may improve the blood supply to the heart. These changes may have long-term benefits in preventing heart disease. The study will hopefully improve understanding of how to reduce muscle loss while attempting to reverse diabetes. It will also show how diabetes affects muscle quality. The results could lead to improved medical treatments and programmes in the future.

In past studies, the low-energy diet has led to symptoms like constipation, dizziness, fatigue, thirst, and/or headache in some people taking part. These tend to get better with fibre-based laxatives and time. It is also important to drink as much water as needed when on the diet. With exercise, there are risks of injury to joints, bones, and muscles. The supervisor will work with the participants to exercise as safely as possible. Exercise may also lead to fatigue and dizziness in some cases. Again, over time these should improve. Stopping diabetes and blood pressure medications at the beginning of the diet and exercise programme may lead to blood sugar and blood pressure going up. The researchers will monitor this to see if they need to restart medications. Diet and exercise may lead to low blood sugar and low blood pressure, but this is unlikely when not taking medication.

Where is the study run from?
University of Leicester (UK)

When is the study starting and how long is it expected to run for?
October 2022 to September 2025

Who is funding the study?
Wellcome Trust (UK)

Who is the main contact?
Mr Franciskos Arsenyadis, fa280@leicester.ac.uk (UK)

Contact information

Type(s)
Public, Scientific

Contact name
Mr Franciskos Arsenyadis

ORCID ID
<https://orcid.org/0000-0003-3885-0548>

Contact details
Diabetes Research Centre
Leicester General Hospital
Gwendolen Road
United Kingdom
LE5 4PW
+44 (0)1162525564
fa280@leicester.ac.uk

Type(s)

Principal investigator

Contact name

Prof Tom Yates

Contact details

Leicester Diabetes Centre
Leicester General Hospital
Gwendolen Road
United Kingdom
LE5 4PW
+44 (0)116 258 4312
ty20@leicester.ac.uk

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

Integrated Research Application System (IRAS)

326665

University of Leicester protocol reference number

0931

Central Portfolio Management System (CPMS)

57941

Study information

Scientific Title

Combining structured exercise with a low-energy diet to attenuate lean mass loss in South Asian adults living with type 2 diabetes: The COMBINE trial

Acronym

COMBINE

Study objectives

Current study objectives as of 16/02/2026:

It is hypothesised that the combination of exercise training and a Meal Replacement Product (MRP)-based Low Energy Diet (LED) will result in considerable amelioration of expected Lean Body Mass (LBM) losses.

Previous study objectives:

It is hypothesised that the combination of exercise training and a high-protein Meal Replacement Product (MRP)-based Low Energy Diet (LED) will result in considerable amelioration of expected Lean Body Mass (LBM) losses.

Ethics approval required

Ethics approval required

Ethics approval(s)

approved 05/12/2023, West Midlands - Black Country Research Ethics Committee (2 Redman Place, London, E20 1JQ, United Kingdom; +44 (0)2071048388; blackcountry.rec@hra.nhs.uk), ref: 23/WM/0201

Study design

Prospective 2-arm parallel-group open-label randomized blinded-endpoint (PROBE) single-site trial

Primary study design

Interventional

Study type(s)

Efficacy

Health condition(s) or problem(s) studied

Type 2 diabetes and obesity

Interventions

Current interventions as of 16/02/2026:

This study aims to investigate whether a structured exercise training programme combined with a Meal Replacement Product (MRP)-based Low Energy Diet (LED) minimises Lean Body Mass (LBM) losses compared to a MRP-based LED (without structured exercise training) in South Asian adults (≥ 40 and ≤ 65 years old) with type 2 diabetes and obesity over 12 weeks. LBM changes are to be assessed via Dual-energy X-ray absorptiometry (DXA).

Randomisation (at the individual level, stratified by sex) occurs once all baseline measures are completed and individuals are confirmed to be eligible. Individuals deemed ineligible following baseline evaluation will not be randomised. A permuted block randomisation will be performed and an independent statistician will develop a computer-assisted random sequence within the block. When the investigator is not blinded, and the block size is fixed, the allocation is predictable. To preserve allocation concealment, the block size will be varied (e.g., 2, 4).

Intervention 1: Low energy diet

This group will be prescribed a low-energy diet for 12 weeks.

Intervention 2: Low energy diet with structured exercise

This group will follow the same diet as Intervention 1 but will additionally undertake combined aerobic and resistance exercise in a supervised setting.

Both groups will receive the same level of clinical and dietetic management. Both intervention groups will be taken off their glucose-lowering medication and antihypertensive medication by a study physician at the onset of the intervention. However, ACE inhibitors or angiotensin receptor blockers will not be discontinued in the context of albuminuria.

Following the 12-week intervention, both groups will receive a further 4-week post-intervention period of support to safely progress food reintroduction and facilitate exercise self-management longer term.

Previous interventions:

This study aims to investigate whether a structured exercise training programme combined with a high protein Meal Replacement Product (MRP)-based Low Energy Diet (LED) minimises Lean

Body Mass (LBM) losses compared to a high protein MRP-based LED (without structured exercise training) in South Asian adults (≥ 40 and ≤ 65 years old) with type 2 diabetes and obesity over 12 weeks. LBM changes are to be assessed via Dual-energy X-ray absorptiometry (DXA).

Randomisation (at the individual level, stratified by sex) occurs once all baseline measures are completed and individuals are confirmed to be eligible. Individuals deemed ineligible following baseline evaluation will not be randomised. A permuted block randomisation will be performed and an independent statistician will develop a computer-assisted random sequence within the block. When the investigator is not blinded, and the block size is fixed, the allocation is predictable. To preserve allocation concealment, the block size will be varied (e.g., 2, 4).

Intervention 1: High protein low energy diet

This group will be prescribed a low-energy diet supplemented with protein (up to 1.3g/kg bodyweight/day) for 12 weeks.

Intervention 2: High protein low energy diet with structured exercise

This group will follow the same diet as Intervention 1 but will additionally undertake combined aerobic and resistance exercise in a supervised setting.

Both groups will receive the same level of clinical and dietetic management. Both intervention groups will be taken off their glucose-lowering medication and antihypertensive medication by a study physician at the onset of the intervention. However, ACE inhibitors or angiotensin receptor blockers will not be discontinued in the context of albuminuria.

Following the 12-week intervention, both groups will receive a further 4-week post-intervention period of support to safely progress food reintroduction and facilitate exercise self-management longer term.

Intervention Type

Mixed

Primary outcome(s)

Differences in the change of Lean Body Mass between groups measured using Dual-Energy X-ray Absorptiometry at baseline and 12 weeks

Key secondary outcome(s)

Physiological:

1. Anthropometry and body composition (including total body weight, Body Mass Index, waist circumference, fat mass, and appendicular lean mass) measured using Dual-Energy X-ray Absorptiometry at baseline and week 12
2. Cross-sectional area of muscle, echo intensity (indication of intermuscular adipose tissue), fibre pennation angle, muscle and subcutaneous fat thickness, and volume of lower limb muscles such as the rectus femoris (quadriceps) measured using ultrasound imaging probe at baseline and week 12
3. Aerobic capacity (peak oxygen uptake; VO_{2peak}) (to include absolute VO_{2peak} ($L \cdot min^{-1}$) and VO_{2peak} relative to lean body mass (LBM) (i.e., $ml \cdot kg \text{ LBM}^{-1} \cdot min^{-1}$) and relative to overall body mass ($ml \cdot kg \text{ BW}^{-1} \cdot min^{-1}$)) measured using graded treadmill maximal exercise test at baseline and week 12
4. Physical function measured using the Short Physical Performance Battery (SPPB) and sit-to-stand 60 at baseline and week 12

5. Hand grip and lower body strength measured using a hand dynamometer and BIODEX at baseline and week 12
6. Resting metabolic rate (RMR) measured using indirect calorimetry at baseline and week 12

Cardiometabolic:

7. Fasting and post-prandial markers of glucose, insulin and C-peptide in response to a Mixed Meal Tolerance Test at baseline and week 12
8. Mean glucose levels, glucose variability (standard deviation and coefficient of variation), time in ranges above and below range measured using continuous glucose monitoring
9. The proportion of participants achieving diabetes remission defined as HbA1c <6.5% (48mmol/mol) without prescribed glucose-lowering medications between baseline and 12 weeks of the study period
10. Cardiometabolic risk factors, such as blood lipid profile (including total cholesterol, HDL, LDL and triglycerides), urine albumin to creatinine ratio (ACR) and systolic and diastolic blood pressure at baseline and week 12

Behavioural:

11. Device-measured sleep, sedentary time and physical activity measured using a wrist-worn accelerometer at baseline and week 12
12. Habitual diet intake measured using a screening diet diary at baseline and week 12

Patient-reported outcome measures (PROMs):

13. Mental well-being measured using the Hospital Anxiety and Depression Scale and the Diabetes Distress Scale at baseline and week 12
14. Quality of life measured using the EQ-5D-5L instrument at baseline and week 12
15. Disability and functional impairment measured using the WHO Disability Assessment Schedule (WHODAS) at baseline and week 12

Process Measures:

Adherence and compliance to intervention measured using dietary coaching visits attended, prescribed exercise sessions completed, and percentage weight loss achieved through LED

Removed 16/02/2026:

Primary Sub-Study Endpoint:

Change in whole-body insulin sensitivity (M) between groups measured using hyperinsulinaemic euglycaemic clamp at 12 weeks

Completion date

10/09/2025

Eligibility

Key inclusion criteria

Current inclusion criteria as of 16/02/2026:

1. South Asian ethnicity (Self-declared ethnicity of self)
2. Men and women
3. Aged ≥ 40 and ≤ 65 years,
4. A clinically coded diagnosis of type 2 diabetes between 3 months and 10 years previously
5. HbA1c 6.5% (48 mmol/mol) to 10% (86 mmol/mol) if not taking glucose-lowering medication; 6% (42 mmol/mol) to 10% (86 mmol/mol) if taking glucose-lowering medication
6. Body Mass Index ≥ 25 and ≤ 45 kg/m²

7. Self-reported stable weight over the previous 6 months (<+/-5% of bodyweight)
8. Treatment stable; no significant change to glucose-lowering regimen in the preceding 3 months, as determined by a study investigator
9. Able to provide informed consent
10. Able to understand written and spoken English or willing to use the University Hospitals of Leicester professional interpreter service
11. Able to take part in structured exercise training requiring the lower limbs (e.g., able to walk without assists or impairment)
12. Willingness and availability to participate in the proposed interventions to which they may be assigned, including attendance of intervention visits such as exercise sessions and adoption of Low Energy Diet which requires abstinence from alcohol
13. Willingness to self-monitor glucose, blood pressure and weight

Previous inclusion criteria:

1. South Asian ethnicity (Self-declared ethnicity of self)
2. Men and women
3. Aged ≥ 40 and ≤ 65 years,
4. A clinically coded diagnosis of type 2 diabetes between 3 months and 10 years previously
5. HbA1c 6.5% (48 mmol/mol) to 10% (86 mmol/mol) if not taking glucose-lowering medication; 6% (42 mmol/mol) to 10% (86 mmol/mol) if taking glucose-lowering medication
6. Body Mass Index ≥ 27.5 and ≤ 45 kg/m²
7. Self-reported stable weight over the previous 6 months (<+/-5% of bodyweight)
8. Treatment stable; no significant change to glucose-lowering regimen in the preceding 3 months, as determined by a study investigator
9. Able to provide informed consent
10. Able to understand written and spoken English or willing to use the University Hospitals of Leicester professional interpreter service
11. Able to take part in structured exercise training requiring the lower limbs (e.g., able to walk without assists or impairment)
12. Willingness and availability to participate in the proposed interventions to which they may be assigned, including attendance of intervention visits such as exercise sessions and adoption of Low Energy Diet which requires abstinence from alcohol
13. Willingness to self-monitor glucose, blood pressure and weight

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Mixed

Lower age limit

40 years

Upper age limit

65 years

Sex

All

Total final enrolment

40

Key exclusion criteria

1. Individuals with type 1, gestational or monogenic diabetes mellitus
2. On insulin therapy (NOTE: No COMBINE participant will be on insulin at baseline as this is an exclusion criterion. An exception may be made for women who are on insulin because of its lack of teratogenic effects rather than because of the inability to control glycaemia on oral agents alone)
3. Drugs or conditions thought by the investigators to have a significant impact on the study protocol or outcomes
4. Serum alanine transaminase level >2.5-fold above the upper limit of the reference range
5. eGFR <45 ml.min⁻¹ per 1.73m²
6. Currently participating in a weight reduction program in addition to routine care
7. Previous bariatric surgery
8. Currently on oral or injected steroids
9. Currently on weight loss medications (not including glucose-lowering medication)
10. Conditions that could impact weight (i.e., active malignancy/treatment in the past year, pregnancy, lactation, planning to become pregnant in the next 8 months)
11. Individuals with a self-reported or diagnosed eating disorder
12. Self-reported milk protein allergy or other allergy or dietary practice that prohibits the use of meal replacement products
13. Previous myocardial infarction, stroke, amputation secondary to type 2 diabetes/peripheral vascular disease, or admission due to cardiovascular disease-related event within 12 months
14. Previous clinically diagnosed atrial fibrillation
15. Previous clinically diagnosed heart failure
16. Pacemaker or implantable cardioverter defibrillator (ICD)
17. Substance abuse. The requirement for alcohol abstinence during the initial 12 weeks will make it unlikely that individuals with alcohol dependence will enrol. Substance abuse will be queried.
18. Presenting with cardiac abnormalities during the exercise electrocardiogram test (inclusive of very high blood pressure)
19. Currently receiving or requiring active treatment for retinopathy.
20. Current participation in another research study with investigational medical product
21. Severe Intolerance or unwillingness/inability to undertake Mixed Meal Tolerance Test (Severity of intolerance to be assessed by a member of the research team during screening visit)

Date of first enrolment

05/02/2024

Date of final enrolment

29/05/2025

Locations

Countries of recruitment

United Kingdom

England

Study participating centre
Leicester Diabetes Centre
Leicester General Hospital NHS Trus
Gwendolen Road
Leicester
England
LE5 4PW

Sponsor information

Organisation
University of Leicester

ROR
<https://ror.org/04h699437>

Funder(s)

Funder type
Research council

Funder Name
Wellcome Trust

Alternative Name(s)
Wellcome, WT

Funding Body Type
Private sector organisation

Funding Body Subtype
Trusts, charities, foundations (both public and private)

Location
United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

Data request processes will be developed for investigators and their teams to conduct follow-up data analyses once the primary paper has been published using anonymised data.

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Protocol article		23/03/2026	25/03/2026	Yes	No
Statistical Analysis Plan			16/02/2026	No	No
Study website		11/11/2025	11/11/2025	No	Yes