

# Exploring a new approach to preventing self-harm in psychiatric inpatient hospitals

<b>Submission date</b> 13/05/2024	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered
<b>Registration date</b> 17/06/2024	<b>Overall study status</b> Completed	<input type="checkbox"/> Protocol
<b>Last Edited</b> 26/03/2026	<b>Condition category</b> Mental and Behavioural Disorders	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data
		<input checked="" type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

Suicide is a leading cause of death for all age groups and a growing public health concern. Literature suggests asking about suicidal ideation reduces it, additionally risk of suicide is dynamic and not linear, with fluctuations throughout a day, week month or even lifespan. Hence interventions must match this. Just In Time Adaptive Interventions (JITAI) use real time information to intervene just in time and are used to respond to varying needs over time. Hence this study will aim to evaluate the feasibility of using a just in time approach to aim to reduce suicidal self-harm in NHS inpatient psychiatric settings.

### Who can participate?

Staff and patients at participating wards will be eligible to participate. With exception to patients who lack capacity, whose first language is not English (the booklet is currently only available in English) or are in seclusion. Agency staff will be unable to participate as it is highly unlikely they will return to the same ward during the course of the trial.

### What does the study involve?

The study uses a booklet- based intervention. Staff on participating wards are invited to training to learn about the background of the study, how to receive consent from patients and how to use the study booklet with patients.

Staff then advertise and recruit eligible patients to the study, once recruited, patient participants start to use the booklet. This involves the patient completing daily questionnaires within the booklet in 1:1 sessions with participating staff, which asks questions pertaining to wellbeing, frequency of suicidal ideation and distress levels. Scores are plotted in charts within the booklet. If the score indicates high risk, staff will then initiate routine risk assessment and management procedures using prompts provided to them during training.

### What are the possible benefits and risks of participating?

The project is a new way of monitoring risks on the ward and can help staff to intervene early. Whilst there are no immediate benefits for people participating in the project, it is hoped that this work will help us to see if this type of intervention method can be used in inpatient settings in the NHS and will provide us with enough data to do a preliminary randomised control trial.

The questionnaires ask about topics that may be upsetting. If patients are distressed by the content of the questionnaires, they can speak to the staff member in their 1:1 Session. If any of the questions cause staff distress this can be discussed and dealt with in clinical supervision. Additionally, staff can attend weekly staff support sessions in addition to clinical supervision and workplace wellbeing. Staff can also contact the psychologist or ward manager for additional support and signposting if needed.

Where is the study run from?

Rotherham Doncaster and South Humber NHS Foundation Trust (UK)

When is the study starting and how long is it expected to run for?

December 2022 to August 2024

Who is funding the study?

National Institute for Health and Care Research (NIHR) (UK).

MindLife (UK)

Who is the main contact?

Prof Jaime Delgadoillo, [jaime.delgadoillo@kcl.ac.uk](mailto:jaime.delgadoillo@kcl.ac.uk)

## Contact information

### Type(s)

Public, Scientific

### Contact name

Miss Ibreeze Ahmed

### Contact details

University of Sheffield

Sheffield

United Kingdom

S10 2TN

-

[iahmed6@sheffield.ac.uk](mailto:iahmed6@sheffield.ac.uk)

### Type(s)

Principal investigator

### Contact name

Prof Jaime Delgadoillo

### ORCID ID

<https://orcid.org/0000-0001-5349-230X>

### Contact details

Professor of Clinical Psychology at the Institute of Psychiatry, Psychology & Neuroscience at

King's College London

London

United Kingdom

-

-  
jaime.delgadillo@kcl.ac.uk

## **Additional identifiers**

**Integrated Research Application System (IRAS)**  
324004

**Central Portfolio Management System (CPMS)**  
56194

## **Study information**

### **Scientific Title**

Feasibility trial of a Just-in-Time Adaptive Intervention (JITAI) to prevent self-harm events in an inpatient care setting

### **Acronym**

JITAI

### **Study objectives**

To evaluate the feasibility of a just in time adaptive intervention (JITAI) to prevent suicidal self-harm events in NHS psychiatric inpatient care settings.

### **Ethics approval required**

Ethics approval required

### **Ethics approval(s)**

approved 04/09/2023, West of Scotland Research Ethics Service (Ward11, Dykebar Hospital, Grahamston Road, Paisley, PA2 7DE, United Kingdom; +44 141 314 0212; WoSREC3@ggc.scot.nhs.uk), ref: 23/WS/0087

### **Study design**

Interventional randomized controlled trial

### **Primary study design**

Interventional

### **Study type(s)**

Treatment

### **Health condition(s) or problem(s) studied**

Suicidal self-harm

### **Interventions**

Participating inpatient care wards across two NHS Trusts will be randomly assigned to two groups. Participants will be randomised and then take part in the same 4-month intervention at a different time (phase one is up to month 4 and phase two from month 4 to month 8), depending on their group assignment.

During the first phase months 1-4 of the study, Group 1 will have immediate access to JITAI training and booklet. Group 2 will serve as a delayed intervention control during this phase. After 4 months, during the second phase, Group 2 will be trained and will have access to JITAI and a booklet. During this second phase, Group 1 will continue to use JITAI.

Participants will be of two types- staff participants and patient participants.

At the start of each phase staff participants within the assigned group will attend training which will provide information on the daily measures being used as well as the JITAI intervention and how they can view the data input by patients to respond. So at this time point, zero staff participants assigned to wards in group one will attend the training. Staff in group 2 will attend training before the start of phase 2- 4 months after the study starts.

The JITAI intervention works based on daily Routine Outcome Monitoring (ROM) and when these indicate things are not going well, this triggers an intervention. The patient participant completes daily ROM using a web-based secure survey, which asks questions about well-being, frequency of suicidal ideation and distress levels. Ward staff review this data daily to assess changes in the patient's well-being and risk factors. If the feedback provided by the JITAI algorithm indicates high risk, staff will then initiate routine risk assessment and management procedures. It's much the same as if you were continuously monitoring a cardiac patient's heart rate during a post-operative (emergency) hospital stay – the technology alerts clinicians to critical changes in cardiac rhythm leading to immediate emergency procedures.

This stepped wedge trial design assumes that significant differences in the primary clinical outcome (e.g., number of self-harm incidents during 4 months) would be found after phase 1 (primary hypothesis test A). No significant differences would be found after phase 2 (secondary hypothesis test B) or after 12 months (secondary hypothesis test C). The present feasibility study seeks to evaluate whether implementing such an intervention (JITAI) and collecting data is at all feasible within this stepped wedge design. The absolute number of self-harm incidents will be collected for each 4-month interval.

## **Intervention Type**

Behavioural

## **Primary outcome(s)**

Staff self-evaluation of the efficacy of risk assessment will be measured using the Risk Assessment and Management Self-Efficacy Scale (RAMSES) at the start of the trial, month 4 and month 8

## **Key secondary outcome(s)**

Measured at admission:

1. The Depression Anxiety Stress Scale, DASS-21
2. The Health of the Nation Outcome Scales (HoNOS)
3. Quality of life enjoyment and satisfaction questionnaire (Q-LES-Q)
4. Alcohol Use Disorders Identification Test AUDIT-10

Measured daily:

1. The Daily Index (DI-5)
2. World Health Organisation Wellbeing Index (WB-10)
3. Items from the Perceived Mastery Scale (PM)

Qualitative interviews will be conducted with some participating staff at the end of the study to gain views on experiences of using the JITAI method in clinical practice and the barriers and enablers to this.

**Completion date**

30/08/2024

## Eligibility

**Key inclusion criteria**

1. Between the ages of 18 - 65 years
2. Inpatient on one of the participating wards
3. Staff participants must be permanent members of staff
4. Staff participants must have completed all the relevant mandatory Trust risk management training
5. Staff member on one of the participating wards who have attended JITAI training

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Mixed

**Lower age limit**

18 years

**Upper age limit**

65 years

**Sex**

All

**Total final enrolment**

0

**Key exclusion criteria**

1. Not fluent in English
2. Patients who are in seclusion
3. Patients assessed not to have the capacity to consent to take part- only participants deemed to have the mental capacity to provide informed consent will be included, this may fluctuate, so we can include patients who may gain capacity during the active phase of data collection
4. Agency staff members

**Date of first enrolment**

22/11/2023

**Date of final enrolment**

30/08/2024

## Locations

### Countries of recruitment

United Kingdom

England

### Study participating centre

**Sheffield Health & Social Care NHS Foundation Trust**

Centre Court

Atlas Way

Sheffield

England

S4 7QQ

### Study participating centre

**Rotherham Doncaster and South Humber NHS Foundation Trust**

Woodfield House

Tickhill Road

Doncaster

England

DN4 8QN

## Sponsor information

### Organisation

Rotherham Doncaster and South Humber NHS Foundation Trust

## Funder(s)

### Funder type

Government

### Funder Name

National Institute for Health and Care Research

### Alternative Name(s)

National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

**Funding Body Type**

Government organisation

**Funding Body Subtype**

National government

**Location**

United Kingdom

**Funder Name**

Mindlife UK LTD

## **Results and Publications**

**Individual participant data (IPD) sharing plan**

The datasets generated during and/or analysed during the current study are/will be available upon request from the chief investigator, Professor Jaime Delgadillo (jaime.delgadillo@kcl.ac.uk), in a CSV file alongside a data dictionary. The data will be shared with qualified researchers who pre-register their planned secondary analysis plan, for a time-limited period dependent on the planned secondary analyses, in a fully anonymised format and in compliance with the conditions of ethical approval and informed consent provided by participants.

**IPD sharing plan summary**

Available on request