

Assessment of lung water in patients with low blood pressure after infection

Submission date 16/01/2015	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 04/02/2015	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 09/10/2018	Condition category Circulatory System	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Extravascular lung water refers to fluid within the lung. Increase in extravascular lung water is associated with death in very ill patients. The purpose of this study is to compare three methods of lung ultrasound with transpulmonary thermodilution technique.

Who can participate?

Individuals with septic shock, 18 years old or over

What does the study involve?

All patients will be assessed with three methods of lung ultrasound (total B-line scores, BLUE points and scanning eight regions) and the results will be compared with the transpulmonary thermodilution technique.

What are the possible benefits and risks of participating?

A benefit is improvement in the monitoring of the patients. The risks are bleeding, haematoma and pneumothorax.

Where is the study run from?

Phramongkutklao Hospital (Thailand)

When is study starting and how long is it expected to run for?

From March 2013 to July 2014

Who is funding the study?

Phramongkutklao Hospital Foundation Under Her Royal Highness Princess Maha Chakri Sirindhorn Patronage (Thailand)

Who is the main contact?

Dr. Pattarin Pirompnich

Contact information

Type(s)

Public

Contact name

Dr Pattarin Pirompnich

ORCID ID

<https://orcid.org/0000-0003-0181-5394>

Contact details

Division of Pulmonary and Critical Care Medicine
Department of Internal Medicine
Faculty of Medicine
Thammasat University
Klong 1
Klong Luang
Pathumthani
Thailand
12120

Additional identifiers

Study information

Scientific Title

Comparison of three methods of lung ultrasound with the transpulmonary thermodilution technique for assessment of extravascular lung water in patients with septic shock

Study objectives

Extravascular lung water index greater than 10 mL/kg in patients with cardiogenic pulmonary oedema correlates with pulmonary capillary wedge pressure of greater than 20 mm Hg. Measurement of extravascular lung water needs sophisticated tools and use of the invasive transpulmonary thermodilution technique. By contrast, multiple B-lines by transthoracic portable ultrasound have been recently proposed to correlate with increased extravascular lung water in patients with cardiogenic pulmonary oedema.

Lung ultrasound has a good correlation with transpulmonary thermodilution technique for the assessment of extravascular lung water in patients with septic shock.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Institutional review board of the Royal Thai Army Medical Department, 22/04/2009
Reference number: R032h/56

Primary study design

Observational

Study design

Cross sectional study

Study type(s)

Diagnostic

Health condition(s) or problem(s) studied

Septic shock

Interventions

1. Total B-line scores
2. BLUE points
3. Scanning eight regions

These three methods of lung ultrasound were used on each patient to assess extravascular lung water and the results were compared with those from obtained with the transpulmonary thermodilution technique.

Intervention Type

Device

Primary outcome(s)

Best method of lung ultrasound to diagnose pulmonary oedema in patients with septic shock: cutoff point of extravascular water index measured by transpulmonary thermodilution greater than

10 mL/kg predicted bodyweight, which is correlated with a pulmonary capillary wedge pressure of greater than 20 mmHg to indicate pulmonary oedema

Key secondary outcome(s)

Cutoff point of total B-line scores to predict extravascular lung water index of 10 mL/kg or more

Completion date

30/07/2014

Eligibility

Key inclusion criteria

1. Age 18 years old or over
2. Septic shock requiring central venous catheterisation and arterial catheterisation for continuous pressure monitoring

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 Years

Sex

All

Key exclusion criteria

1. Valvular heart disease
2. Pregnancy
3. Acute or history of pulmonary embolism
4. Lung resection surgery
5. Acute respiratory distress syndrome
6. Interstitial lung diseases
7. Multilobar pneumonia
8. Pleural diseases

Date of first enrolment

01/09/2013

Date of final enrolment

30/06/2014

Locations**Countries of recruitment**

Thailand

Study participating centre

Phramongkutklao Hospital

315 Phayathai Road

Ratchathewi

Bangkok

Thailand

10400

Sponsor information**Organisation**

Phramongkutklao Hospital Foundation Under Her Royal Highness Princess Maha Chakri Sirindhorn Patronage

Funder(s)**Funder type**

Charity

Funder Name

Phramongkutklao Hospital Foundation Under Her Royal Highness Princess Maha Chakri Sirindhorn Patronage

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Stored in repository

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/09/2018		Yes	No