

# Fish oil-based lipid emulsion decrease inflammation and bronchopulmonary dysplasia in extremely premature infants

<b>Submission date</b> 07/09/2015	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 15/09/2015	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
<b>Last Edited</b> 27/06/2018	<b>Condition category</b> Neonatal Diseases	<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

### Background and study aims

Preterm infants have significantly lower long chain polyunsaturated fatty acids (LC-PUFA) concentrations than full term infants because they miss the major period of docosahexaenoic acid (DHA) and eicosapentaenoic acid (EPA) development in the brain and other organs during the third trimester of pregnancy. LC-PUFA play a potentially significant role in the modulation (control) of developmental processes affecting short- and long-term health outcomes related to growth, body composition, immune and allergic responses, and nutrition-related chronic disease. The most widely used formulation for parenteral (intravenous) nutrition (PN) is a soybean oil-based lipid emulsions (LE), which is rich in n-6 LC-PUFA and phytosterols which may cause liver toxicity and inflammation. LC-PUFAs (EPA and DHA) from fish oil may prevent the inflammatory response and even reduce any inflammation that exists. Early fish oil-based lipid emulsion (LE) administration to very low birth weight premature infants may reduce the development of chronic lung disease (CLD) and liver dysfunction by decreasing inflammatory cytokines (small signalling proteins) with IL-1 $\beta$  and IL-6 levels (suggesting inflammation) in serum (a component of the blood) and bronchoalveolar lavage fluid (fluid from the lungs). The aim of this study is to investigate whether this is the case.

### Who can participate

Very low birth weight (VLBW) premature infants with a birth weight <1250gm or gestation age <32 weeks.

### What does the study involve

Participants are randomly allocated into one of two groups. Those in group 1 are given total parenteral nutrition (TPN) containing a fish oil-based lipid emulsion. Those in group 2 are given TPN containing conventional soybean lipid emulsion. Treatment starts within 48 hours of birth and lasts for at least seven days. The dose starts at 1gm/kg but then increases by 1gm/kg per day until the maximum dose of 3gm/kg.

What are the possible benefits and risks of participating  
The treatment may reduce the development of lung and liver problems by reducing inflammation

Where is the study run from  
Changhua Christian Hospital (Taiwan)

When is study starting and how long is it expected to run  
March 2012 to February 2014

Who is the main contact  
Chien-Chou Hsiao  
68206@cch.org.tw

## Contact information

**Type(s)**  
Scientific

**Contact name**  
Dr Chien-Chou Hsiao

**ORCID ID**  
<https://orcid.org/0000-0002-4523-5300>

**Contact details**  
135 Nanshiao Street  
Changhua city  
Taiwan  
500  
886-4-7238595 ext 1902  
68206@cch.org.tw

## Additional identifiers

**Protocol serial number**  
101-CCH-IRP-15

## Study information

**Scientific Title**  
Fish oil-based lipid emulsion attenuates inflammatory cytokines and development of bronchopulmonary dysplasia in extremely premature infants: a randomised controlled trial

**Study objectives**  
The aim of this study to is investigate whether giving preterm infants an early supply of long chain polyunsaturated fatty acids (LC-PUFA) and fish oil-based lipid emulsions (LE) containing  $\omega$ -3 LC-PUFAs helps prevent the development of bronchopulmonary dysplasia (BPD) by modulating inflammation and neonatal immune function.

## **Ethics approval required**

Old ethics approval format

## **Ethics approval(s)**

Changhua Christian Hospital Institutional Review Board Committee A, 22/12/2011, ref: CCH-IRB-110909

## **Study design**

Randomised controlled trial

## **Primary study design**

Interventional

## **Study type(s)**

Treatment

## **Health condition(s) or problem(s) studied**

Insufficient docosahexaenoic acid (DHA) and eicosapentaenoic acid (EPA) in premature babies

## **Interventions**

Sixty very low birth weight (VLBW) premature infants were randomized to 2 groups:

1. Study group received total parenteral nutrition (TPN) containing fish oil-based lipid emulsion (SMOF lipid contain 30% soybean-LCT, 30% MCT, 25% olive oil and 15% fish oil).
2. Control group received TPN containing conventional soybean lipid emulsion (Lipovenoes MCT 20%).

Treatment was started within 48 hours of birth and then for at least the next 7 days. The dose of both lipid emulsions were 1 gm/kg at first day of TPN, increasing to 1 gm/kg per day until 3 gm/kg. Macronutrients were provided using the same products for glucose and amino acid solutions in both groups. The premature infant was fed by NG tube or bottle with breast milk or premature formula milk if breast milk was not enough.

## **Intervention Type**

Supplement

## **Primary outcome(s)**

Comparison of immune effects of Interleukin (IL)-1 $\beta$  and IL-6, assessed on the first 48 hours and 7 days after TPN use, including serum IL-1 $\beta$  and IL-6 and bronchoalveolar lavage fluid from tracheal aspiration if intubated infants. The levels of cytokines in the serum and bronchoalveolar lavage fluid were estimated by enzyme-linked immunosorbent assay.

## **Key secondary outcome(s)**

1. Mortality
2. Ventilator use days
3. Oxygen dependent days
4. Length of hospital stay (LOS)
5. Growth rate
6. Liver function
7. Parenteral nutrition associated cholestasis (PNAC)
8. Bronchopulmonary dysplasia (BPD)
9. Retinopathy of prematurity (ROP)

- 10. Necrotizing enterocolitis (NEC)
- 11. Intraventricular hemorrhage (IVH)
- 12. Nosocomial infection

Follow up program will finish when the infant is discharged.

**Completion date**

28/02/2014

## Eligibility

**Key inclusion criteria**

PVLBW infants who weighed below 1250 gm and gestational age below 32 weeks

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Neonate

**Sex**

All

**Key exclusion criteria**

1. Chromosomal disorders or lethal congenital abnormalities
2. Congenital cyanotic heart disease
3. Anatomic obstructive gastrointestinal pathologies, such as intestinal malrotation with or without volvulus, stenosis, or atresia, gastroschisis, omphalocele and Hirschsprung's disease
4. Confirmed or family history of hereditary metabolic disorder

**Date of first enrolment**

01/03/2012

**Date of final enrolment**

25/02/2014

## Locations

**Countries of recruitment**

Taiwan

**Study participating centre**

Changhua Christian Hospital

Taiwan

500

# Sponsor information

## Organisation

Changhua Christian Hospital

## ROR

<https://ror.org/05d9dtr71>

# Funder(s)

## Funder type

Hospital/treatment centre

## Funder Name

Changhua Christian Hospital

## Alternative Name(s)

CCH

## Funding Body Type

Private sector organisation

## Funding Body Subtype

Other non-profit organizations

## Location

Taiwan

# Results and Publications

## Individual participant data (IPD) sharing plan

### IPD sharing plan summary

Available on request

### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>	results	01/06/2019		Yes	No