

The ADHD controversy project: Long-term effects of ADHD diagnosis

Submission date 08/05/2020	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
		<input checked="" type="checkbox"/> Protocol
Registration date 28/08/2020	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
Last Edited 12/02/2024	Condition category Mental and Behavioural Disorders	<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Attention-Deficit/Hyperactivity Disorder (ADHD) is among the most common mental disorders in children and adolescents, and it is a strong risk factor for several adverse psychosocial outcomes over the lifespan. Over recent decades there have been substantial increases in clinical diagnosis and medication for ADHD in many countries. Large between- and within-country variations in diagnosis and medication rates have been reported. Due to ethical and practical considerations, few studies have examined the effects of receiving a diagnosis, and there is a lack of research on the effects of medication on long-term outcomes.

This study is a part of a research project titled “the ADHD controversy project” which has four aims organized in four work packages: (WP1) To examine the prognosis of ADHD (with and without medication) compared to patients with other psychiatric diagnoses, patients in contact with public sector child and adolescent psychiatric outpatient clinics (without a diagnosis), and the general population; (WP2) Examine within-country variation in ADHD diagnoses and medication rates by clinics’ catchment area; (WP3) Identify causal effects of being diagnosed with ADHD and (WP4) ADHD medication on long-term outcomes, including emergency visits, comorbidity, education, contact with child welfare services, contact with adult psychiatric health services, employment and income, welfare dependency, crime, self-harm, suicide, and mortality. This trial registration covers WP3 of this project which aim is to identify causal effects of being diagnosed with ADHD on outcomes, using variation in diagnosis as an instrumental variable (IV).

Who can participate?

The patient sample is all persons in Norway between 5 and 18 years of age that were in contact with child and adolescent psychiatric outpatient clinics in 2009-2011 (i.e. birth cohorts 1991-2006). We will also include a control sample that is a random sample from the general population of equal size to the patient sample. The control sample is defined as individuals in Norway between 5-18 years not in contact with child and adolescent psychiatric outpatient clinics (regardless of reason) during the calendar years 2009-2011.

What does the study involve?

Our comparative analysis of prognosis will be based on survival analysis and mixed-effects models. Our analysis of variation will apply mixed-effects models and generalized linear models.

We have two identification strategies to retrieve the effect of being diagnosed with ADHD and of receiving medication on long-term outcomes. Both strategies rely on using preference-based instrumental variables, which in our project are based on provider preferences for ADHD diagnosis and medication.

What are the possible benefits and risks of participating?

None

Where is the study run from?

All public sector child and adolescent psychiatric outpatient clinics in Norway (BUP centres in Norway).

When is the study starting and how long is it expected to run for?

October 2015 to December 2021

Who is funding the study?

1. The Research Council of Norway
2. The Western Norway Regional Health Authority

Who is the main contact?

Prof. Arnstein Mykletun, arnstein.mykletun@fhi.no

Mr Tarjei Widding-Havneraas, tarjei.widding-havneraas@helse-bergen.no

Ms Ashmita Chaulagain, ashmita.chaulagain@helse-bergen.no

Ms Ingvild Lyhmann, ingvild.lyhmann@helse-bergen.no

Contact information

Type(s)

Scientific

Contact name

Prof Arnstein Mykletun

ORCID ID

<https://orcid.org/0000-0002-3878-0079>

Contact details

Center for Research and Education in Forensic Psychiatry

Haukeland University Hospital

Sandviksleitet 1

Bergen

Norway

5036

+47 91689600

arnstein.mykletun@uit.no

Type(s)

Scientific

Contact name

Mr Tarjei Widding-Havneraas

ORCID ID

<https://orcid.org/0000-0002-5823-7508>

Contact details

Center for Research and Education in Forensic Psychiatry
Haukeland University Hospital
Sandviksleitet 1
Bergen
Norway
5036
+47 92451487
tarjei.widding-havneraas@helse-bergen.no

Type(s)

Scientific

Contact name

Ms Ashmita Chaulagain

ORCID ID

<https://orcid.org/0000-0001-5467-181X>

Contact details

Center for Research and Education in Forensic Psychiatry
Haukeland University Hospital
Sandviksleitet 1
Bergen
Norway
5036
+47 99895219
ashmita.chaulagain@helse-bergen.no

Type(s)

Scientific

Contact name

Ms Ingvild Lyhmann

ORCID ID

<https://orcid.org/0000-0003-1376-1354>

Contact details

Center for Research and Education in Forensic Psychiatry
Haukeland University Hospital
Sandviksleitet 1
Bergen
Norway
5036
+47 98675744
ingvild.lyhmann@helse-bergen.no

Additional identifiers

Protocol serial number

288585

Study information

Scientific Title

Long-term effects of ADHD diagnosis

Study objectives

H1: There are causal effects of being diagnosed with ADHD (for patients on the margin of diagnosis) on a range of long-term outcomes: Emergency visits, comorbidity, education, contact with child welfare services, contact with adult psychiatric health services, employment and income, welfare dependency, crime, self-harm, suicide, and mortality. (Two-tailed hypotheses.)

H0: There are no causal effects of being diagnosed with ADHD (for patients on the margin of diagnosis) on these long-term outcomes.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 25/01/2018, Regional Committee for Medical and Health Research Ethics -South-East Norway Committee D (Gullhaugveien 1-3, 0484 Oslo, Norway; +47 22845511; post@helseforskning.etikkom.no), ref: 2017/2150/REK south-east D

Primary study design

Observational

Study design

Observational longitudinal cohort study

Study type(s)

Quality of life

Health condition(s) or problem(s) studied

Diagnosis of ADHD

Interventions

The study is a longitudinal cohort study, where the aim is to answer questions about the effects of ADHD diagnosis on long-term outcomes by studying patients that receive diagnosis due to provider preference.

The study sample consists of individuals between 5-18 years that were in contact (any reason) with child and adolescent psychiatric outpatient clinics in 2009-2011, with follow-up data to 2021. We examine several psychosocial outcomes that to some extent depends on age. For instance, welfare dependency will be measured from age 18, crime from 15 years of age, child welfare service is relevant until 18 and in some special cases until 23 years of age, and adult psychiatric health services is relevant from 18 years of age. Drop out from education is

dependent on track and school system. Mortality, emergency ward visit, the onset of new mental disorder is relevant regardless of age.

To identify the effect of ADHD diagnosis on long-term outcomes, we will use child and adolescent psychiatric outpatient clinics' catchment area variation in diagnosis as an instrumental variable (IV). IV is a common identification strategy in health-services research that exploits the variation induced in the treatment variable (here, ADHD diagnosis) by a plausibly exogenous (as-if random) event or process (i.e., the instrument). The role of the instrument is to "isolate" exogenous variation in the treatment, and hence to remove the endogenous (confounded) variation so that the analysis only uses the exogenous variation to estimate the treatment effect.

Our identification strategies exploit as-if random variation in providers' preference to diagnose patients with ADHD across Norwegian public sector child and adolescent psychiatric outpatient clinics.

Intervention Type

Other

Primary outcome(s)

We have nine primary outcomes (point 1a, 3a ... to 9b), and one overall primary outcome for the general prognosis, measured at age 21. The primary outcome for general prognosis is the avoidance of all primary adverse outcomes among those as listed below, measured as dichotomy until age 21. The secondary outcome is the number of adverse outcomes among those listed below. The specific primary and secondary outcomes for ADHD diagnosis, measured using patient records, are:

1. Emergency ward visits, measured anytime
 - 1.1. Primary: Emergency ward visit for severe injury
 - 1.2. Secondary: Any contact with emergency department. Analyses for type of and severity of injury. Measured for anytime.
2. Child welfare and protection services (Barnevernet), measured at any time during the period where services are available by law.
 - 2.1. Primary: Any intervention from the child welfare and protection services (in Norwegian: Barnevernet)
 - 2.2. Secondary: Type of contact, type of interventions (voluntary or involuntary), specifically for parental guideline and foster care.
3. Contact with adult psychiatric health services anytime from the age of 18
 - 3.1. Primary: Any contact
 - 3.2. Secondary: Type of contact
4. Onset of other mental disorder, measured anytime during follow-up
 - 4.1. Primary: Onset of other mental disorders > 2 years after initial contact with public sector child and adolescent psychiatric outpatient clinic in 2009-2011
 - 4.2. Secondary analyses for types of diagnosis
5. Educational attainment, measures to be included at age-appropriate time-points.
 - 5.1. Primary: Non-completion of high-school, 5 years after planned start of high-school.
 - 5.2. Secondary outcomes for average grades level at primary school, enrollment to high-school, delayed high-school completion, non-completion of high school, grades from high-school at completion, enrollment into higher education after high-school, dropout from higher education.
6. Welfare dependency measured at age-appropriate time-points.
 - 6.1. Primary: Award of work assessment allowance (WAA, in Norwegian AAP)

6.2. Secondary outcomes for receipt of any welfare benefits more than 1G over one year. Secondary outcomes for types of benefits (unemployment benefits, social benefits of 0.5G or more, sickness absence, work assessment allowance with more than 1 year duration, disability benefits for life), NEET (Not in Education, Employment or Training anytime from 15/16 onwards for 1 year or more).

7. Crime measures to be included at age-appropriate time-points

7.1. Primary: Any criminal charge (regardless of conviction)

7.2. Secondary: Type of crime

8. Self-harm, measured at any time during the follow up

8.1. Primary: Any type of self-harm

8.2. Secondary: Any type of secondary self-harm

9. Mortality, measured at any time during the follow up

9.1. Primary: Any mortality

9.2. Secondary: Cause of death in categories of suicide, injuries and other

Key secondary outcome(s)

As specified above

Completion date

31/12/2021

Eligibility

Key inclusion criteria

The patient sample is all persons in Norway between 5 and 18 years of age that were in contact with child and adolescent psychiatric outpatient clinics in 2009-2011 (i.e. birth cohorts 1991-2006). We will also include a control sample that is a random sample from the general population of equal size to the patient sample. The control sample is defined as individuals in Norway between 5-18 years not in contact with child and adolescent psychiatric outpatient clinics (regardless of reason) during the calendar years 2009-2011.

We have four different sample definitions:

1. All patients diagnosed with ADHD in 2009-2011, regardless of referral reason.
2. All patients referred to child and adolescent outpatient clinics from GP's with suspicion of ADHD in 2009-2011, regardless of whether the patient was diagnosed with ADHD.
3. All patients in contact with child and adolescent outpatient clinics for any reason in 2009-2011, regardless of whether the patient was diagnosed with ADHD.
4. All individuals in child and adolescent outpatient clinics catchment area in 2009-2011

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Child

Lower age limit

5 Years

Upper age limit

18 Years

Sex

All

Total final enrolment

178820

Key exclusion criteria

Does not meet inclusion criteria

Date of first enrolment

01/01/2009

Date of final enrolment

31/12/2011

Locations**Countries of recruitment**

Norway

Study participating centre

All public sector child and adolescent psychiatric outpatient clinics in Norway (BUP centres in Norway)

Norway

-

Study participating centre

Haukeland University Hospital

Center for Research and Education in Forensic Psychiatry

Sandviksleitet 1

Bergen

Norway

5036

Sponsor information**Organisation**

Haukeland University Hospital

ROR

Funder(s)

Funder type

Government

Funder Name

The Research Council of Norway

Funder Name

The Western Norway Regional Health Authority

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are not expected to be made available due personal data protection regulations (GDPR), restrictions in the ethics approval and restrictions by the registry owners, they may not be shared. The data are obtainable from the Norwegian Health Directorate, Norwegian Institute of Public Health and Statistics Norway. We can share statistical code.

IPD sharing plan summary

Not expected to be made available

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Protocol article	protocol	19/01/2021	22/01/2021	Yes	No
Study website	Study website	11/11/2025	11/11/2025	No	Yes