

Efficacy of indomethacin in in-vitro fertilisation treatment in the modified natural cycle

Submission date 07/03/2007	Recruitment status No longer recruiting	☐ Prospectively registered
		☐ Protocol
Registration date 07/03/2007	Overall study status Completed	☐ Statistical analysis plan
		☐ Results
Last Edited 26/08/2021	Condition category Pregnancy and Childbirth	☐ Individual participant data
		☐ Record updated in last year

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr M L Haadsma

Contact details

University Medical Centre Groningen
Department of Obstetrics and Gynaecology
Groningen
Netherlands
9700 RB
+31 (0)50 361 6161
m.l.haadsma@og.umcg.nl

Additional identifiers

Protocol serial number

NL656 (NTR907)

Study information

Scientific Title

Efficacy of indomethacin in in-vitro fertilisation treatment in the modified natural cycle

Study objectives

Indomethacin is known to be a strong inhibitor of ovulation in the spontaneous menstrual cycle. Therefore we assume that the use of indomethacin prior to follicle aspiration in In-Vitro Fertilisation-treatment in the Modified Natural Cycle (IVF-MNC) significantly decreases the number of patients with one or more premature ovulations compared to placebo.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approval received from the local medical ethics committee (Medisch Ethische Toetsingscommissie UMC Groningen) on the 24th June 2005 (ref: 2005/074).

Primary study design

Interventional

Study design

Randomised, placebo controlled, parallel group, double blinded trial

Study type(s)

Treatment

Health condition(s) or problem(s) studied

In-vitro fertilisation-treatment in the modified natural cycle, IVF in the natural cycle

Interventions

Use of indomethacin (Indocid®) 50 mg versus placebo during IVF-treatment in the modified natural cycle.

Dosage scheme: three times a day, starting on the day of ovulation triggering and ending on the morning of the follicle aspiration (total of seven capsules per cycle).

Intervention Type

Drug

Phase

Not Specified

Drug/device/biological/vaccine name(s)

Indomethacin (Indocid®)

Primary outcome(s)

Number/percentage of patients per study group that have one or more ovulations prior to follicle aspiration.

Key secondary outcome(s)

Number/percentage of patients per study group that achieve an on-going pregnancy (defined as an intact intra-uterine pregnancy at 12 weeks gestation)

Completion date

01/11/2007

Eligibility

Key inclusion criteria

1. Indication for IVF or Intracytoplasmic Sperm Injection (ICSI) treatment
2. Age 18 up to 37 years
3. Ovulatory cycle of 26 to 35 days

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 Years

Sex

Female

Key exclusion criteria

1. Prior IVF or ICSI treatment unless the last treatment was successful
2. Ovarian cysts disabling adequate sonographic assessment of the ovaries
3. Contraindications for indomethacin, such as asthma or prior gastro-intestinal ulcer

Date of first enrolment

05/12/2005

Date of final enrolment

01/11/2007

Locations

Countries of recruitment

Netherlands

Study participating centre

University Medical Centre Groningen

Groningen

Netherlands

9700 RB

Sponsor information

Organisation

University Medical Centre Groningen (UMCG) (The Netherlands)

ROR

<https://ror.org/03cv38k47>

Funder(s)**Funder type**

Hospital/treatment centre

Funder Name

University Medical Centre Groningen (UMCG) (The Netherlands)

Results and Publications**Individual participant data (IPD) sharing plan****IPD sharing plan summary**

Not provided at time of registration