

Study drug ELX-02 for patients with Alport syndrome

Submission date 09/03/2022	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 05/05/2022	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 15/10/2024	Condition category Genetic Diseases	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

The main aim of this study is to learn how well the study drug, ELX-02, works and how safe it is for patients with Alport syndrome caused by nonsense mutations. Alport syndrome caused by nonsense mutations is a genetic disorder that decreases kidney function and can also affect the ears and decrease hearing. In this form of Alport syndrome, the nonsense mutation prevents the body from producing the full-length Collagen IV protein that forms the filters in the kidney and has an important role in ear function. There is currently no treatment for Alport syndrome caused by nonsense mutations.

Who can participate?

Patients aged between 6 and 30 years with Alport syndrome caused by nonsense mutations

What does the study involve?

Patients will be in the study for about 26 weeks. This includes a screening period of up to 42 days, a treatment period of 60 days and a follow-up period of 90 days. Patients will receive a treatment of ELX-02 injected once a day for 8 weeks. After their last treatment, patients will be followed up for 12 weeks.

What are the possible benefits and risks of participating?

Participating in this research study will allow patients to play an active role in health care, help others by contributing the medical research and learn more about their condition.

Patients may or may not benefit from taking part in this research study. It is possible that their Alport syndrome symptoms will improve while they are receiving the study drug, however, there is no guarantee that patients will receive any benefits.

To date, more than 126 people have been exposed to ELX-02 at different doses. The most commonly reported side effects in studies of ELX-02 were injection site reactions, ear discomfort, ear pain, headaches and dizziness. Common injection site reactions are redness, warmth, pain or itching near the injection site. These reactions occurred in more than 80% of patients receiving ELX-02 and nearly 40% of patients in the placebo (dummy drug) group. Injection site reactions may appear immediately or be delayed by several days after the injection. Most reactions are mild and improve shortly after the injection is administered. Rare injection site reactions may include blistering and scab forming on the injection site several days

later. On very rare occasions, injection site reactions may be slow to improve, continue for an unknown period of time (several months), and may cause a scar on the skin to change colour around the injection site.

Blood sample collection may cause faintness and/or swelling, pain, redness, bruising, bleeding at the collection site, or infection (rarely) at the site where the needle is inserted. Temporary skin irritation may occur during an ECG from the patches or gel that is used. Patients may experience changes in hearing, speech, noise or ringing in the ears, or dizziness. Patients may experience a build-up of waste in their blood due to kidney injury. The most common complication with the kidney biopsy is blood in the urine, which usually stops within a few days, and pain at the biopsy site for a few hours. Rare cases include surgery to control the blood in the urine, kidney hematoma infection that requires antibiotic treatment or high blood pressure. The study may involve risks to the unborn child that are not yet known at this time, so patients and partners of patients should not become pregnant while participating in the study.

Where is the study run from?

Eloxx Pharmaceuticals Inc. (USA)

When is the study starting and how long is it expected to run for?

February 2022 to April 2024

Who is funding the study?

Eloxx Pharmaceuticals Inc. (USA)

Who is the main contact?

Prof. Detlef Bockenhauer

d.bockenhauer@ucl.ac.uk

Contact information

Type(s)

Principal investigator

Contact name

Prof Detlef Bockenhauer

Contact details

Great Ormond Street Hospital for Children

Great Ormond Street

London

United Kingdom

WC1N 3JH

+44 (0)207 4059200

d.bockenhauer@ucl.ac.uk

Type(s)

Scientific

Contact name

Dr Ali Hariri

Contact details

Eloxx Pharmaceuticals Inc
480 Arsenal Way, Suite 130
-Watertown
United States of America
MA 02472
+1 (0)203 464 9420
ali.hariri@eloxxpharma.com

Additional identifiers

ClinicalTrials.gov (NCT)
NCT05448755

Clinical Trials Information System (CTIS)
2022-000604-35

Integrated Research Application System (IRAS)
1005163

Central Portfolio Management System (CPMS)
52088

Protocol serial number
EL-014

Study information

Scientific Title

A Phase II open-label pilot study to evaluate the safety and efficacy of subcutaneously administered ELX-02 in patients with Alport syndrome with Col4A5 and Col4A3/4 nonsense mutation

Study objectives

Primary objective: The safety and tolerability of ELX-02 will be evaluated in this trial.

Secondary objectives: The readthrough effect of ELX-02 in the expression of Col IV will be evaluated in the kidney and ear. The expression of the Col IV in the kidney will be evaluated functionally by the changes in proteinuria and hematuria, two main early hallmarks of kidney disease in AS.

Histologically, the expression of Col IV will be evaluated by the expression of Col IV in the basement membrane. While the focus of the trial is to evaluate the efficacy in the kidney, and as such the patients are not selected based on their auditory phenotype. However, it is expected that most of the participants to have auditory phenotypes. The evaluation of the hearing of the participants is an exploratory endpoint.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approval 04/05/2022, East of England - Cambridge South Research Ethics Committee (Equinox House, City Link, Nottingham, NG2 4LA, UK; +44 (0)20 7104 8284; cambridgesouth.rec@hra.nhs.uk), ref: 22/EE/0064

Study design

Open-label pilot study

Primary study design

Interventional

Study type(s)

Treatment, Safety, Efficacy

Health condition(s) or problem(s) studied

Alport syndrome

Interventions

Patients will receive a treatment of ELX-02 0.75 mg/kg subcutaneously administered once a day for 8 weeks. After their last treatment, patients will be followed up for 12 weeks.

Intervention Type

Drug

Phase

Phase II

Drug/device/biological/vaccine name(s)

ELX-02

Primary outcome(s)

1. Incidence and characteristics of adverse events (AEs) collected using Common Terminology Criteria for Adverse Events (CTCAE) Version 5.0 from signing the informed consent form until the day 150/End of Study (EOS) visit
2. Injection site reactions classified and graded as per Division of AIDS (DAIDS) guidelines at the days 1, 15, 30, 45 and 60/End of Treatment (EOT) visits
3. Vital signs and physical examination performed at the initial, baseline, days 1, 15, 30, 45 and 60 /EOT visits
4. Audiometric testing (including high-frequency audiometry) and tinnitus and dizziness status assessed at the initial, baseline, days 30, 60/EOT, 90, 120 and 150/EOS visits
5. 12-lead ECG performed at the initial and day 60/EOT visits
6. Clinical laboratory evaluations (haematology, chemistry, urinalysis) performed at the initial, days 1, 15, 30, 45, 60/EOT, 90, 120 and 150/EOS visits

Key secondary outcome(s)

1. Proteinuria measured via the urine protein/creatinine ratio (UPCR) procedure at the initial, baseline, days 1, 15, 30, 60/EOT, 90, 120 and 150/EOS visits
2. Col IV protein expression measured using a renal biopsy collected at the initial and day 60 /EOT visits
3. Hematuria measured microscopically at the initial, baseline, days 1, 15, 30, 60/EOT, 90, 120 and 150/EOS visits

Completion date

04/04/2024

Eligibility

Key inclusion criteria

1. Evidence of signed and dated informed consent/assent document(s) indicating that the participant (and/or their parent/legal guardian) has been informed of all pertinent aspects of the trial
2. Understands, and is willing and able to comply with all scheduled visits, treatment plan, laboratory tests, and other study procedures
3. Male and female participants
4. Aged between 6 and 30 years
5. A confirmed diagnosis of X-linked or autosomal recessive Alport syndrome with a documented nonsense mutation of Col4A5 in a male or nonsense mutation of Col4A3 or Col4A4 (male or female)
6. The nonsense mutation should be UAG or UGA
7. eGFR >60 ml/min/1.73 m² (based on Chronic Kidney Disease Epidemiology Collaboration [CKD-EPI] for ages ≥18 years and Schwartz formula for participants <18 years)
8. Urinary protein based on two spot urine collections [urine protein/creatinine ratio (UPCR) ≥500 mg/g]
9. Stable regimen of ACEi/ARB for 4 weeks before screening (unless there is a contraindication)
10. Must be willing to abstain from strenuous exercise during the 48 h prior to study visits
11. Females of childbearing potential and males capable of fathering a child must meet the contraception requirements (outlined in protocol section 6.3)
12. Non-lactating females
13. Females on hormone replacement therapy (estrogen or progesterone) or contraceptive therapy must be stabilized on a product and dose for at least 30 days prior to Screening
14. Have not received systemic medications with the potential to impair renal function on a frequent basis (e.g., nonsteroidal anti-inflammatory drugs [NSAIDs]) or with ototoxic potential (e.g., quinine or salicylates), or any injectable aminoglycosides for a period of at least 14 days prior to dosing
15. Negative human immunodeficiency virus (HIV), hepatitis B surface antigen (HBsAg), and hepatitis C virus antigen (HCV Ag) serology tests at Screening
16. No history of alcohol or drug abuse within the 6 months prior to Screening
17. Body Mass Index (BMI) of 19.0 to 30.0 kg/m² (inclusive). Participants with a lower BMI may be entered into the study at the discretion of the investigator following consultation with the Sponsor

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Mixed

Lower age limit

6 years

Upper age limit

30 years

Sex

All

Total final enrolment

3

Key exclusion criteria

1. Participants who are investigational site staff members directly involved in the conduct of the study and their family members, site staff members otherwise supervised by the Investigator, or participants who are Eloxx Pharmaceuticals employees directly involved in the conduct of the study
2. Participation in clinical study including administration of any investigational drug or device in the last 30 days or 5 half-lives (whichever is longer) prior to investigational product dosing in the current study
3. Use of prohibited medications as defined in (protocol) section 6.1 within the specified windows
4. History of any comorbidity which in the opinion of the investigator might confound the study or pose an additional risk in administering the study drug to the participant
5. History of any organ transplantation
6. Mutation consistent with autosomal dominant Alport syndrome
7. Liver disease characterized by cirrhosis or portal hypertension. Participants with alanine aminotransferase (ALT), aspartate aminotransferase (AST), and/or a total bilirubin 3.0 times the upper limit of normal (ULN) will be excluded
8. History of congestive heart failure diagnosed clinically or with documented left ventricular ejection fraction (LVEF) $\leq 40\%$
9. Evidence or history of clinically relevant psychiatric condition
10. A positive urine drug screen (amphetamines, benzodiazepines, cocaine and opiates) at Screening
11. Screening supine 12-lead electrocardiogram (ECG) demonstrating any clinically significant findings as judged by the Investigator
12. Participants with any abnormalities in clinical laboratory tests at Screening, considered by the study Investigator as clinically relevant
13. Major surgery within 180 days (6 months) of Screening
14. History of dialysis
15. Known allergy to any aminoglycoside
16. Participants with any acute medical situation unresolved within 14 days of first dose that is considered of significance by the Investigator
17. Participants with >10 dB change in threshold between audiometric Initial test to baseline test for the frequencies from 0.5 -12.5 kHz
18. Dizziness Handicap Inventory (DHI) score at screening ≥ 16 for adults, and active dizziness reported for pediatric participants.

Date of first enrolment

31/10/2022

Date of final enrolment

30/03/2023

Locations

Countries of recruitment

United Kingdom

England

Australia

Study participating centre

Great Ormond Street Hospital for Children

Great Ormond Street

London

United Kingdom

WC1N 3JH

Study participating centre

Royal Free Hospital

Pond Street

London

United Kingdom

NW3 2QG

Study participating centre

Royal Children's Hospital

Dept of Nephrology

50 Flemington Road

Parkville

Victoria

Australia

3051

Study participating centre

Monash Medical Center

246 Clayton Road

Clayton

Victoria

Australia

3168

Sponsor information

Organisation

Eloxx Pharmaceuticals (United States)

ROR

<https://ror.org/0021myq38>

Funder(s)

Funder type

Industry

Funder Name

Eloxx Pharmaceuticals Inc.

Results and Publications

Individual participant data (IPD) sharing plan

The data-sharing plans for the current study are unknown and will be made available at a later date. All study data collected through the electronic case report form (eCRF) will be anonymized by a unique ID. The paper documents including personal data (Informed consent form and screening log) will be filed in the Investigator's File that will be locked and only the Investigational Team will have access.

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
HRA research summary			26/07/2023	No	No
Protocol file	version 2.1	21/03/2023	12/06/2023	No	No