

Obstetric analgesia: a comparison of patient controlled pethidine, remifentanyl and fentanyl in labour

Submission date 14/02/2006	Recruitment status No longer recruiting	☐ Prospectively registered
Registration date 14/02/2006	Overall study status Completed	☐ Protocol
Last Edited 04/11/2008	Condition category Pregnancy and Childbirth	☐ Statistical analysis plan
		☐ Results
		☐ Individual participant data
		☐ Record updated in last year

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr M R Douma

Contact details

Bronovo Hospital
Department of Obstetrics and Gynaecology
Bronovolaan 5
The Hague
Netherlands
2597 AX
+31 (0)61 427 6591
maritdouma@hotmail.com

Additional identifiers

Protocol serial number

NTR543

Study information

Scientific Title

Acronym

Baringspijn

Study objectives

The hypothesis of this study is that the new opioid remifentanil will provide less side-effects and better pain relief during labour, than the conventional opioids fentanyl and pethidine.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Received from local medical ethics committee

Primary study design

Interventional

Study design

Double-blind randomised active controlled parallel group clinical trial

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Labour pain

Interventions

The following drugs are used in a patient controlled method:

1. Remifentanil - 40 µg for 2 minutes, total dosage: 1200 µg/h
2. Pethidine - loading dose of 50 mg, then 5 mg for 10 minutes, total dosage: 200 mg
3. Fentanyl - loading dose of 50 µg, then 20 µg for 5 minutes, total dosage: 240 µg/h

Medication will be started in active labour and will be continued until complete dilation of the cervix is achieved. Baseline non-invasive measurements will be made, including maternal blood pressure, heart rate, respiratory rate and pulse oximetry. Measurements will be recorded every 30 minutes. Also, an observer sedation score will be recorded every 30 minutes.

Intervention Type

Drug

Phase

Not Specified

Drug/device/biological/vaccine name(s)

Pethidine, remifentanil, fentanyl

Primary outcome(s)

1. Quality of pain relief determined by Visual Analogue Scale, patient controlled analgesia (PCA) demands/rewards and the number of patients crossing over to epidural analgesia. Pain scores will be assessed every hour.

2. Patient satisfaction, assessed every hour
3. Foetal outcome as determined by Apgar, NACS and requirement for naloxone, taken after delivery
4. Presence of opioid substances in umbilical and maternal blood samples, taken after delivery

Key secondary outcome(s)

No secondary outcome measures

Completion date

01/09/2006

Eligibility

Key inclusion criteria

1. Aged at least 18 years old
2. Between 37 and 42 weeks of gestation
3. American Society of Anaesthesiologists (ASA) physical status I or II

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 Years

Sex

All

Key exclusion criteria

1. ASA physical status greater than or equal to III
2. Obesity (body mass index [BMI] equal or more than 40 kg/m²)
3. Substance abuse history
4. High risk patients: including pre-eclampsia (diastolic pressure equal or more than 100, proteinuria), hepatic insufficiency or renal failure, severe asthma, poorly controlled diabetes mellitus
5. Premature labour
6. Drug allergy; history of hypersensitivity to opioid substances

Date of first enrolment

10/08/2005

Date of final enrolment

01/09/2006

Locations

Countries of recruitment

Netherlands

Study participating centre

Bronovo Hospital

The Hague

Netherlands

2597 AX

Sponsor information

Organisation

Bronovo Hospital (The Netherlands)

ROR

<https://ror.org/03r781319>

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

Bronovo Hospital (The Netherlands) - research funds

Results and Publications

Individual participant data (IPD) sharing plan**IPD sharing plan summary**

Not provided at time of registration