

# Study of treatments for Dupuytren's contractures

<b>Submission date</b> 01/09/2020	<b>Recruitment status</b> No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
<b>Registration date</b> 18/09/2020	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
<b>Last Edited</b> 20/06/2024	<b>Condition category</b> Musculoskeletal Diseases	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

Dupuytren's contractures (DC) are scar-like tissue that form under the skin of the palm of the hand, causing one or more fingers to bend into the palm (<https://www.nhs.uk/conditions/dupuytren-contraction/>). They prevent them from straightening fully which causes difficulty with many everyday tasks like washing, dressing, or using a computer keyboard. It is common: in 2016 NHS England spent £36 million treating 16,700 patients with DC. DC are most commonly treated using one of two procedures and it is not known which is best. Needle fasciotomy (NF) involves cutting the contracture by moving a medical needle back and forth through it until it snaps, releasing the finger. It can be done in a clinic room and leaves no skin wound. Recovery takes 1-2 weeks. Limited fasciectomy (LF) involves cutting out the contractures and stitching the skin up. This is done in an operating theatre with the patient asleep or the arm "frozen". Recovery takes 4-12 weeks. About 8 in 10 treatments done currently are LF. The problem can come back (recur) after either treatment. Studies suggest that this happens sooner and more commonly with NF, but LF has a greater risk of causing other problems which restrict hand function, like finger stiffness and pain. NF is cheaper for the NHS, but any savings may be offset by an increased need for future treatments after recurrence. The aims of this study are to determine whether NF is not substantially worse than LF at preserving hand function and to find out which treatment offers the best value for money.

### Who can participate?

Patients aged 18 or over with DC in one or more untreated fingers.

### What does the study involve?

Participants will be randomly allocated by a computer to have treatment with either needle fasciotomy or limited fasciectomy. Participants will be followed up after 2, 3, 4 and 6 weeks, and 6, 12 and 24 months. The participant will be asked to complete questionnaires either by post or online for completion at 2, 3 and 4 weeks after the treatment asking about their ability to carry out normal daily activities with the hand, hand function and quality of life. The participant will be asked to complete these again 6 weeks after the treatment (a routine NHS clinic appointment) and at 6, 12 and 24 months at home by post or online. Participants will be asked to take measurements of their finger using a SLICK device at 6, 12 and 24 months. Participants will be shown how to do this and given an instruction manual at their initial clinic appointments.

Participants will also have access to an online video demonstration of how to do this. The researchers will also compare other measures of hand function, overall health, finger straightness, recurrence and costs. Some people will be contacted up to 2 years after they have had treatment to hear their thoughts on the treatment and about taking part in the study. The discussion will cover hand function before and after treatment, expectations and experiences of the treatment and recovery, and views on taking part in the study. The researchers have arranged to share their results with a research team comparing limited fasciectomy with a third treatment option, collagenase injections, in another study funded by NIHR. This will provide even more information on how best to treat DC.

**What are the possible benefits and risks of participating?**

There is no direct benefit of taking part in the study, but this study may help us to treat people with Dupuytren's contracture in the future. Participants will receive the same level of care whether they choose to participate in the study or not. Participants may not benefit personally from taking part in this study, but because of the contact with the research team, they will have more regular or frequent opportunity. Both treatments are a part of standard NHS care, so there is no extra risk involved in receiving them as part of the study. Taking part will mean spending some extra time to complete questionnaires but these can be completed online or by post. There are no physical risks if participants take part in the optional interviews. It is possible that participants talking about their feelings and other issues related to their diagnosis and treatment may feel overwhelmed. Participants are able to pause or finish the discussion with the researcher at any time.

**Where is the study run from?**

Nottingham University Hospitals NHS Trust and Nottingham Clinical Trials Unit (NCTU) (UK)

**When is the study starting and how long is it expected to run for?**

January 2020 to March 2026

**Who is funding the study?**

National Institute for Health Research (NIHR) (UK)

**Who is the main contact?**

Aisha Shafayat

Hand2@nottingham.ac.uk

## Contact information

**Type(s)**

Public

**Contact name**

Mrs Aisha Shafayat

**ORCID ID**

<https://orcid.org/0000-0003-0412-381X>

**Contact details**

Nottingham Clinical Trials Unit

Building 42, Room A17

University Park

University of Nottingham  
Nottingham  
United Kingdom  
NG7 2RD  
+44 (0)115 8231578  
hand2@nottingham.ac.uk

**Type(s)**

Scientific

**Contact name**

Prof Tim Davis

**ORCID ID**

<https://orcid.org/0000-0001-5729-6680>

**Contact details**

Nottingham University Hospital NHS Trust  
QMC Campus  
Nottingham  
United Kingdom  
NG7 2UH  
+44 (0)115 9249924 ext 64337  
tim.davis@nuh.nhs.uk

## Additional identifiers

**Integrated Research Application System (IRAS)**

282087

**Central Portfolio Management System (CPMS)**

45939

## Study information

**Scientific Title**

Hand-2: Needle fasciotomy versus limited fasciectomy for the treatment of Dupuytren's contractures of the fingers: a randomised, multi-centre non-inferiority trial

**Acronym**

Hand-2

**Study objectives**

To establish the clinical and cost-effectiveness of needle fasciotomy (NF) versus limited fasciectomy (LF) for treatment of DC in the NHS in terms of patient-reported hand function and symptoms and resource utilisation.

**Ethics approval required**

Old ethics approval format

## **Ethics approval(s)**

Approved 08/09/2020, London - Chelsea Research Ethics Committee (Health Research Authority, Skipton House, 80 London Road, London, SE1 6LH, UK; +44 (0)20 7972 2561; chelsea.rec@hra.nhs.uk), REC ref: 20/LO/0911

## **Study design**

Randomized; Interventional; Design type: Treatment, Surgery

## **Primary study design**

Interventional

## **Study type(s)**

Treatment

## **Health condition(s) or problem(s) studied**

Dupuytren's contractures

## **Interventions**

There are two different treatments for Dupuytren's contracture that are most commonly used in the NHS. However, there is no strong evidence about which of these treatments is best for patients or the NHS. The trial design for Hand-2 was guided by a randomised feasibility study (Hand-1 ISRCTN11164292) which demonstrated that Hand-2 can be successfully performed. It showed it is possible to recruit and retain participants to a trial of two very different treatment (needle fasciotomy (NF) and limited fasciectomy (LF)) for DC and allowed selection of appropriate outcome measures which patients consider relevant to this experience. It also demonstrated through qualitative interviews the acceptability of this trial to patients and health professionals.

Hand-2 is a multi-centre, two-arm, parallel, randomised, non-inferiority trial comparing the outcome of NF and LF among adults eligible for treatment of DC within the NHS. An internal pilot phase with an integrated QuinteT Recruitment Intervention (QRI) will be used to optimise the trial design. Potential participants will be identified by the local research team at sites from the GP referral letters and clinic lists. A short version of the information leaflet will be sent to potentially eligible patients explaining Dupuytren's contractures and the study before their clinic appointment.

This will enable us to assess how the study is explained to patients and how they are invited to take part. If they give permission, they will be asked to complete an audio-recording consent form. If they do not wish to be further approached for the study, they will have the option of opting out by letting any member of clinic staff know.

If after being told about the study by the surgeon and researcher, and reading a full patient information sheet, the patient wishes to take part, they will be asked to sign a consent form. This may take place on the same day as the clinic appointment, or the patient can consent on another day agreed with the researcher. They will be asked to complete a questionnaire and a researcher will take some measurements of the hand. Their surgical treatment will then be determined at random, and the patient will be informed verbally and in writing which of the following two treatments they will receive:

1. Needle fasciotomy – performed in a clinic room under local anaesthetic
2. Limited fasciectomy – performed in an operating theatre under regional or general anaesthetic.

Patients in the study will join the usual NHS waiting list for their treatment. Other than receiving their allocated treatment and completing one questionnaire about their current hand functionality, there are no other study procedures for patients to complete on the day of surgery. At 2, 3, and 4 weeks after treatment participants will be asked to complete questionnaires regarding their recovery from treatment. Six weeks after treatment participants will be asked to attend a clinic appointment that is part of their usual NHS care. At this appointment, we will ask participants to complete questionnaires and have further measurements taken of their hand. At 3, 6, 12 and 24 months after treatment participants will be asked to complete further questionnaires about their recovery, hand function and quality of life. We will use reminders to ensure data collection.

Reminders may be used. For the earlier timepoints, this may be a text message or e-mail on the day that the questionnaire is due. For the later time points (e.g. 12 and 24 months) this reminder will be sent approximately 2 weeks and 4 weeks after the questionnaire is due. The 24-month follow-up will capture early DC recurrences, which may be extended to 5 years and participants will be asked to consent to this.

Participant questionnaires contain standard questions that are used to assess how the hand looks, feels and works before and after treatment.

In addition at 6 weeks, 6, 12 and 24 months data will be collected about NHS resource use (visits to other health care professionals such as GPs, and any other treatments that participants may have used). Measurement of the angle of straightness of the affected finger(s) and grip strength will be taken at the clinic visit at 6 weeks. In addition, linear measurements of extension and flexion of the study finger will be obtained by the participant at home using the SLiCK device.

Evidence of patient experience and acceptability of treatment for DC in the long term is limited. Hand-2 participants will be invited to take part in qualitative interviews (by phone or face to face) to explore their experiences and the acceptability of treatment. Up to 30 semi-structured interviews will be conducted and some participants may be interviewed more than once to assess if views change over time.

Patients can agree to take part in none, all or some of the three components of this study, and consent will be taken for each component:

1. Audio recording of consultation(s) with the surgeon or other clinic staff
2. The feasibility trial, where surgical treatment is decided randomly
3. Qualitative interviews

Members of staff who are recruiting patients to the study may also be interviewed to assess their views on the trial and its conduct, including knowledge of the evidence and personal views about the interventions, how they explain the study to patients and perceived barriers to recruitment. This can help to highlight possible recruitment difficulties.

### **Intervention Type**

Procedure/Surgery

### **Primary outcome(s)**

Participant-reported assessment of hand function using the PEM questionnaire at 12 months post-treatment intervention

### **Key secondary outcome(s)**

1. Participant-reported hand function and hand health measured using the Hand Health Profile of the PEM at 3 weeks, 6 weeks, 3 months, 6 months and 24 months post-treatment intervention
2. Participant-reported assessment of location-specific health (the hand) using the Single Assessment Numeric Evaluation (SANE) tool and the Measure Yourself Medical Outcome Profile (MYMOP) tool at 2 weeks, 3 weeks, 4 weeks, 6 weeks, 3 months, 6 months and 24 months post-treatment intervention
3. Loss of finger extension using measuring tool (SLiCK device) at 6 weeks, and 6, 12 and 24 months post-treatment intervention
4. Adverse events and complications, recurrence of DC, and revisions or salvage surgery recorded in the Case Report Form and participant questionnaire responses up to 24 months post-treatment intervention
5. Health-related quality of life, health resource use, and cost-effectiveness assessed by a health economic analysis of health & social services costs at 2 weeks, 3 weeks 6 weeks, 6 months, 12 and 24 months post-treatment intervention

**Completion date**

31/03/2026

## Eligibility

**Key inclusion criteria**

1. Age  $\geq$  18 years
2. One or more fingers with a Dupuytren's contracture of  $>30$  with functional problems
3. No previous treatment for Dupuytren's contracture on affected finger
4. Well defined cord(s) and suitable for NF (needle fasciotomy) or LF (limited fasciectomy)
5. Able to comply with the requirements of the study up to 24 months post-intervention

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Adult

**Lower age limit**

18 years

**Sex**

All

**Total final enrolment**

406

**Key exclusion criteria**

1. Dupuytren's contracture of the distal interphalangeal joints (DIP) only
2. Planned dermofasciectomy or very limited fasciectomy (excision of  $<1$  cm cord segment)
3. Previously recruited into this study for treatment of either hand

**Date of first enrolment**

01/04/2021

**Date of final enrolment**

30/01/2024

## Locations

**Countries of recruitment**

United Kingdom

England

Scotland

**Study participating centre****Queens Medical Centre**

Derby Road

Nottingham

United Kingdom

NG7 2UH

**Study participating centre****Stoke Mandeville Hospital**

Mandeville Road

Aylesbury

United Kingdom

HP21 8AL

**Study participating centre****Aberdeen Royal Infirmary**

Foresterhill Road

Aberdeen

United Kingdom

AB25 2ZN

**Study participating centre****Southampton General Hospital**

Coxford Road

Southampton

United Kingdom

SO16 5YA

**Study participating centre**  
**Nuffield Orthopaedic Centre**  
Windmill Road  
Headington  
Oxford  
United Kingdom  
OX3 7HE

**Study participating centre**  
**Basingstoke and North Hampshire Hospital**  
Aldermaston Road  
Basingstoke  
United Kingdom  
RG24 9NA

**Study participating centre**  
**Royal Hampshire County Hospital**  
Romsey Road  
Winchester  
United Kingdom  
SO22 5DG

**Study participating centre**  
**Glasgow Royal Infirmary**  
84 Castle St  
Glasgow  
United Kingdom  
G4 0SF

**Study participating centre**  
**Whiston Hospital**  
Warrington Road  
Prescot  
United Kingdom  
L35 5DR

**Study participating centre**  
**Royal United Hospital Bath**  
Combe Park

Bath  
United Kingdom  
BA1 3NG

**Study participating centre**  
**Wrightington Hospital**  
Hall Lane  
Appley Bridge  
Wigan  
United Kingdom  
WN6 9EP

**Study participating centre**  
**North Devon District Hospital**  
Raleigh Heights  
Barnstaple  
United Kingdom  
EX31 4JB

**Study participating centre**  
**South Tyneside Hospital**  
Harton Lane  
South Shields  
United Kingdom  
NE34 0PL

**Study participating centre**  
**Sunderland Royal Hospital**  
Kayll Road  
Sunderland  
United Kingdom  
SR4 7TP

**Study participating centre**  
**Great Western Hospital**  
Marlborough Road  
Swindon  
United Kingdom  
SN3 6BB

**Study participating centre**  
**Leicester Royal Infirmary**  
Infirmary Square  
Leicester  
United Kingdom  
LE1 5WW

**Study participating centre**  
**Royal Blackburn Hospital**  
Haslingden Road  
Blackburn  
United Kingdom  
BB2 3HH

**Study participating centre**  
**Chelsea and Westminster Hospital NHS Foundation Trust**  
Chelsea & Westminster Hospital  
369 Fulham Road  
London  
United Kingdom  
SW10 9NH

## **Sponsor information**

**Organisation**  
Nottingham University Hospitals NHS Trust

**ROR**  
<https://ror.org/05y3qh794>

## **Funder(s)**

**Funder type**  
Government

**Funder Name**  
NIHR Evaluation, Trials and Studies Co-ordinating Centre (NETSCC); Grant Codes: NIHR127393

# Results and Publications

## Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are not expected to be made available.

## IPD sharing plan summary

Not expected to be made available

## Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Protocol article</a>		19/06/2024	20/06/2024	Yes	No
<a href="#">HRA research summary</a>			28/06/2023	No	No
<a href="#">Study website</a>	Study website	11/11/2025	11/11/2025	No	Yes