

Clinical evaluation of a urinary biomarker for the early detection of clear cell renal carcinoma

Submission date 22/01/2024	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 06/02/2024	Overall study status Ongoing	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 25/11/2025	Condition category Cancer	<input type="checkbox"/> Individual participant data <input checked="" type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Kidney cancer (KC) is the third most common type of tumor in the urinary system for both men and women and sadly, it's also one of the top causes of cancer-related deaths. Right now, there's no reliable test to catch KC early. This study aims to change that by looking into whether a substance called Perilipin-1 in your urine could be a good sign that you might have KC, no matter what stage the cancer is at. In simpler terms, this study is trying to find a urine test that could help catch KC earlier than it is right now.

Who can participate?

Patients with KC and healthy controls aged 18 years old and over

What does the study involve?

The present study is an analytical, case-control, prospective study that considers two groups: Cases: patients who have been diagnosed with KC at any stage who are going to undergo one of the following:

1. Surgical treatment (partial or radical nephrectomy).
2. Systemic treatment for KC with histological confirmation of renal cancer by percutaneous biopsy

Controls: healthy volunteers who undergo a preventive check-up where an image (ultrasound, CT, or MRI) excludes renal masses (benign or malignant).

Urine creatinine and PLIN-1 will be measured in urine samples at baseline (cases and controls) and six months after surgery (only for cases). The urine sample will be obtained from voided urine, except for the baseline sample of patients undergoing nephrectomy, which will be obtained by bladder catheterization on the day of surgery.

What are the possible benefits and risks of participating?

Participants will not receive any experimental therapy, and there will be no cost to participate in the study. Finding a urinary marker that allows population screening will permit early diagnosis of KC, which will imply a better oncological outcome.

No risks provided at the time of registration.

Where is the study run from?

Instituto de Ciencias e Innovación en Medicina (ICIM) (Santiago, Chile)
Centro de Medicina Regenerativa
Clinica Alemana (Santiago, Chile)

When is the study starting and how long is it expected to run for?

April 2023 to December 2026

Who is funding the study?

Clinica Alemana (Santiago, Chile)
Universidad del Desarrollo

Who is the main contact?

David Contador, PhD - dcontador@udd.cl
Rodrigo Acuña, PhD - rodrigoacuna@udd.cl
Hugo Otaola, PhD - hotaola@alemana.cl

Contact information

Type(s)

Public, Scientific, Principal investigator

Contact name

Dr Hugo Otaola Arca

ORCID ID

<https://orcid.org/0000-0003-4952-2283>

Contact details

Las Hualtatas 5099, depto 203, Vitacura
Santiago
Chile
7630352
+56 941089452
hotaola@alemana.cl

Type(s)

Principal investigator, Public, Scientific

Contact name

Dr David Contador Martínez

ORCID ID

<https://orcid.org/0000-0002-0383-214X>

Contact details

Avenida La Plaza 680, Las Condes
Santiago
Chile

7610315

-

dcontador@udd.cl

Type(s)

Principal investigator, Scientific, Public

Contact name

Dr Rodrigo Acuña Astudillo

ORCID ID

<https://orcid.org/0000-0002-2033-3107>

Contact details

Avenida La Plaza 680, Las Condes

Santiago

Chile

7610315

-

rodrigoacuna@udd.cl

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

Protocol serial number

ID 1201

Study information

Scientific Title

Evaluation of urinary perilipin-1 (PLIN-1) as a Diagnostic biomarker in patients with clear renal cancer

Acronym

TOREADOR

Study objectives

Urinary Perilipin-1 is a valid diagnostic biomarker of kidney cancer in all its stages.

Ethics approval required

Ethics approval required

Ethics approval(s)

approved 24/04/2023, Comité Etico Científico de la Facultad de Medicina Clínica Alemana de Santiago-Universidad Del Desarrollo (Avda. La Plaza 680, Las Condes, Santiago, 7610658, Chile; None provided; ceccasudd@udd.cl), ref: 2023-25

Study design

Analytical case-control prospective study

Primary study design

Observational

Study type(s)

Diagnostic, Screening

Health condition(s) or problem(s) studied

Screening in patients with kidney cancer

Interventions

This is an analytical, case-control, prospective study that involves two experimental groups:

a) Cases: patients diagnosed with kidney cancer who are going to undergo nephrectomy (partial or radical). A urine sample (30 mL) will be obtained from a Folley catheter at the beginning of the kidney surgery. Additionally, a voided urine sample (30 mL) will be collected six months after surgery.

b) Controls: healthy volunteers. A voided urine sample (30 mL) will be collected at the inclusion in the study.

Intervention Type

Other

Primary outcome(s)

1. Creatinine (mg/dL) in urine samples measured with the modified JAFFE method, using the CREATININE DMSO kit (Applied Clinical Chemistry. S.A, Amposta, Spain) at baseline (cases and controls) and at six months after surgery (only for cases)

2. Perilipin-1 (PLIN-1) (ng/dL) in urine samples measured with a Sandwich ELISA assay using Human PLIN1/Perilipin Sandwich ELISA kit at baseline (cases and controls) and at six months after surgery (only for cases)

Key secondary outcome(s)

There are no secondary outcome measures

Completion date

31/12/2026

Eligibility

Key inclusion criteria

Cases:

1. Patients \geq 18-yr, male or female
2. Diagnosed with kidney cancer at any stage
3. Who are willing to undergo one of the following:
 - 3.1. Surgical treatment (partial or radical nephrectomy).
 - 3.2. Systemic treatment for kidney cancer with histological confirmation of renal cancer by percutaneous biopsy

Controls:

1. Healthy volunteers, \geq 18-yr old, male or female

2. Who undergo a preventive check-up where an image (ultrasound, CT or MRI) excludes renal masses (benign or malignant)

Participant type(s)

Healthy volunteer, Patient

Healthy volunteers allowed

Yes

Age group

Mixed

Lower age limit

18 years

Upper age limit

150 years

Sex

All

Total final enrolment

0

Key exclusion criteria

Cases:

1. Personal history of kidney surgery (for benign or malignant disease).
2. Current or recent treatment (<6 months) with chemotherapy, immunotherapy, or immune checkpoint inhibitors.
3. Clinical diagnosis of kidney cancer without histological confirmation

Controls:

1. Study in progress for possible kidney cancer
2. Personal history of kidney surgery, both for benign and malignant pathology
3. Current or recent treatment (<6 months) with chemotherapy, immunotherapy, or immune checkpoint inhibitors

Date of first enrolment

18/05/2023

Date of final enrolment

31/12/2025

Locations

Countries of recruitment

Chile

Study participating centre

Centro de Medicina Regenerativa, Instituto de Ciencias e Innovación en Medicina (ICIM)
Avenida la Plaza 680, Las Condes
Santiago
Chile
7610315

Sponsor information

Organisation

Clínica Alemana

ROR

<https://ror.org/028ynny55>

Organisation

Universidad del Desarrollo

ROR

<https://ror.org/05y33vv83>

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

Clínica Alemana de Santiago

Alternative Name(s)

Clínica Alemana

Funding Body Type

Private sector organisation

Funding Body Subtype

Other non-profit organizations

Location

Spain

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study will be stored in a non-publicly available repository (www.redcap.alemana.cl)

IPD sharing plan summary

Stored in non-publicly available repository