

A randomised, double-blind, placebo-controlled, dose escalation study of single and multiple oral dose administration of BIIB014 in subjects with moderate to late stage parkinson's disease who are also receiving treatment with levodopa

Submission date 08/09/2006	Recruitment status No longer recruiting	☒ Prospectively registered
Registration date 13/09/2006	Overall study status Completed	☐ Protocol
Last Edited 25/04/2014	Condition category Nervous System Diseases	☐ Statistical analysis plan
		☐ Results
		☐ Individual participant data
		☐ Record updated in last year

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Prof David J Brooks

Contact details
MRC Clinical Sciences Centre
Faculty of Medicine
Imperial College
Hammersmith Hospital
Du Cane Road
London
United Kingdom
W12 0NN

Additional identifiers

Protocol serial number

Study information

Scientific Title

Study objectives

To establish a safe and tolerable BIIB014 dose range for future studies in subjects with moderate to late stage Parkinson's Disease (PD).

Ethics approval required

Old ethics approval format

Ethics approval(s)

Thames Valley Multi-Centre Research Ethics Committee (ref: 06/MRE12/67)

Ethics approval added as of 04/07/2007:

Nizam's Institute of Medical Sciences (India) (ref: EC/NIMS/702©/2007)

Primary study design

Interventional

Study design

Double-blind, placebo-controlled, multicentre, dose-escalation, single dose/washout/multiple dose study

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Moderate to late stage Parkinson's Disease (PD)

Interventions

Please note that this trial started in May 2007 and the anticipated trial end date has been extended to March 2008.

Group one: Intervention treatment - BIIB014 at either 5 mg, 5 mg/10 mg, 10 mg, 10 mg/30 mg, 30 mg, 30 mg/100 mg, 50 mg, 100 mg. If patients are randomised to a single dose group they will receive 26 days of treatment (72 hour washout after first dose). If patients are randomised to a two dose group they will receive 28 days of treatment (seven days at the lower dose and 21 days at the higher dose).

Group two: Control treatment - placebo and levodopa treatment. The control group is the standard of care. All doses are in capsule form to be taken once daily in the morning with food.

Intervention Type

Drug

Phase

Not Specified

Drug/device/biological/vaccine name(s)

BIIB014, levodopa

Primary outcome(s)

1. The number and proportion of subjects with adverse events
2. Assessment of clinical laboratory parameters
3. Assessment of vital signs
4. Assessment of ECG parameters

Key secondary outcome(s)

1. To explore the Pharmacokinetic (PK) drug interactions between BIIB014 and L-DOPA in subjects with moderate to late stage PD
2. To explore the PK of BIIB014 when administered as adjunct therapy to subjects with moderate to late stage PD
3. To explore the activity of BIIB014 when administered as adjunct therapy to subjects with moderate to late stage PD

Completion date

31/12/2007

Eligibility**Key inclusion criteria**

Inclusion criteria amended as of 18/06/2007:

1. Male or female subjects, aged 30 to 78 years old, inclusive
2. Must carry a diagnosis of idiopathic PD according to UK Parkinson's Disease Society Brain Bank Clinical Diagnostic Criteria made by a Movement Disorder Specialist, and be Hoehn & Yahr Stage II to IV (inclusive) when 'off'
4. Subjects must be on a stable dose of L-3,4-Dihydroxyphenylalanine (L-DOPA) / carbidopa or L-DOPA / benserazide for at least 4 weeks prior to enrollment
5. Some subjects must demonstrate a definite end of L-DOPA dose wearing off (at least two hours 'off' time per waking day) and must be able to keep accurate patient diaries of PD activity
6. Except for L-DOPA and certain allowed dopamine agonists, must not be receiving any other PD medication (Current treatment with certain dopamine agonists is allowed but must have been on a stable dose for at least 4 weeks prior to enrollment)

Inclusion criteria provided at time of registration:

1. Male or female subjects, aged 30 to 78 years old, inclusive
2. Must carry a diagnosis of idiopathic PD according to UK Parkinson's Disease Society Brain Bank Clinical Diagnostic Criteria made by a Movement Disorder Specialist, and be Hoehn & Yahr Stage II to IV (inclusive) when 'off'
4. Subjects must be on a stable dose of L-3,4-Dihydroxyphenylalanine (L-DOPA)/carbidopa or L-DOPA/benserazide for at least three weeks prior to enrollment
5. Must demonstrate an excellent motor response to L-DOPA, and have a definite end of L-DOPA dose wearing off (at least two hours 'off' time per waking day)
6. Subjects must not be receiving any other PD medications

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 Years

Sex

All

Key exclusion criteria

Exclusion criteria amended as of 18/06/2007:

1. Mini Mental State Examination (MMSE) score of <26
2. History or clinical features consistent with an atypical parkinsonian syndrome
3. Any significant non-PD central nervous system disorder
4. Any significant AXIS I psychiatric disease from the Diagnostic and Statistical Manual of Mental Disorders (DSM)
5. History of surgical intervention for PD
6. History of certain malignancies
7. History of severe allergic anaphylactic reactions to any drug
8. Clinically significant baseline Electrocardiogram (ECG)
9. Orthostatic hypotension
10. HbA1c >7.0%

Exclusion criteria provided at time of registration:

1. Mini Mental State Examination (MMSE) score of less than 27
2. History or clinical features consistent with an atypical parkinsonian syndrome
3. Any significant non-PD central nervous system disorder
4. Any significant AXIS I psychiatric disease from the Diagnostic and Statistical Manual of Mental Disorders (DSM)
5. History of surgical intervention for PD
6. History of malignancy
7. History of severe allergic anaphylactic reactions to any drug
8. Clinically significant baseline Electrocardiogram (ECG)
9. Orthostatic hypotension

Date of first enrolment

31/12/2006

Date of final enrolment

31/12/2007

Locations**Countries of recruitment**

United Kingdom

England

India

Study participating centre
MRC Clinical Sciences Centre
London
United Kingdom
W12 0NN

Sponsor information

Organisation
Biogen Idec (USA)

ROR
<https://ror.org/02jqkb192>

Funder(s)

Funder type
Industry

Funder Name
Biogen Idec (USA)

Alternative Name(s)

Funding Body Type
Private sector organisation

Funding Body Subtype
For-profit companies (industry)

Location
United States of America

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary
Not provided at time of registration