

StereoTactic radiotherapy for wet Age-Related macular degeneration (STAR)

Submission date 28/11/2014	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 13/12/2014	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 16/07/2025	Condition category Eye Diseases	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Age-related macular degeneration (AMD) is a painless eye condition that leads to loss of central vision. There are two main types of AMD, called dry AMD and wet AMD. Wet AMD is more serious and without treatment, vision can deteriorate within days. This study investigates the use of radiation to treat wet AMD, called stereotactic radiosurgery. The radiation is delivered using a robotically controlled device that projects overlapping beams of radiation onto the macula, the part of the eye that is affected by wet AMD. This study aims to determine if stereotactic radiosurgery can maintain vision and reduce the need for regular injections of ranibizumab (the standard anti-VEGF drug used to treat wet AMD).

Who can participate?:

Males and females aged 50 years or over with wet AMD requiring anti-VEGF treatment at the time of entry to the study.

What does the study involve?

Participants will be randomly allocated to receive either radiation (stereotactic radiotherapy) or simulated placebo (sham) treatment. They will be followed up regularly for two years, and then again at the end of three and four years for a safety visit. Participants will also receive injections of ranibizumab (Lucentis) into their eye if their wet AMD is active.

What are the possible benefits and risks of participating?

The main benefit of stereotactic radiotherapy is that it may reduce the number of eye injections that people require, or in some cases eliminate them entirely. Previous studies have suggested that the patients who will be eligible for the study would obtain a better vision outcome with radiotherapy (with injections as needed) than with injections alone. The main risk of radiation is that it can sometimes damage the healthy tissue in the macula and thereby damage vision. Subtle changes to the macula can sometimes be seen using specialised testing (fluorescein angiography) but in the majority of people this does not affect the vision.

Where is the study run from?

King's College Hospital (UK).

When is the study starting and how long is it expected to run for?
Recruitment to the study will begin in December 2014 and continue for approximately 41 months. Each participant will be in the study for 4 years.

Who is funding the study?
NIHR Efficacy and Mechanism Evaluation Programme (UK).

Who is the main contact?
Mr Tim Jackson
t.jackson1@nhs.net

Contact information

Type(s)
Scientific

Contact name
Dr Tim Jackson

Contact details
King's College London
Department of Ophthalmology
King's College Hospital
Denmark Hill
London
United Kingdom
SE5 9RS
+44 (0)20 3299 1297
t.jackson1@nhs.net

Type(s)
Public

Contact name
Mrs Riti Desai

Contact details
Ophthalmology Department
Research Office
1st Floor, Normanby Building
King's College Hospital
Denmark Hill
London
United Kingdom
SE5 9RS
+44 (0)20 3299 1297
ritidesai@nhs.net

Additional identifiers

ClinicalTrials.gov (NCT)

NCT02243878

Integrated Research Application System (IRAS)

86810

Study information

Scientific Title

StereoTactic radiotherapy for wet Age-Related macular degeneration (STAR): a randomised, double-masked, sham-controlled, clinical trial comparing low-voltage X-ray irradiation with as needed ranibizumab to as needed ranibizumab monotherapy

Acronym

STAR

Study objectives

The study's primary hypothesis is that the mean number of ranibizumab injections during the first 24 months after randomization will be less in the SRT group than in the sham group. The secondary hypothesis is that participants who undergo SRT will have a non-inferior visual outcome compared with those in the sham group.

Ethics approval required

Old ethics approval format

Ethics approval(s)

NRES Committee London - City & East, REC ref: 13/LO/1207

Primary study design

Interventional

Study design

Randomised double-masked sham-controlled pivotal clinical trial

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Retinal disease/age-related macular degeneration

Interventions

Participants in the treatment arm will receive 16 Gray stereotactic radiotherapy (SRT) with a concomitant injection of ranibizumab.

Intervention Type

Device

Phase

Not Applicable

Drug/device/biological/vaccine name(s)

Not provided at time of registration

Primary outcome(s)

The primary outcome is the mean number of ranibizumab injections over 24 months. SRT will be considered superior to sham if the mean number of injections in the SRT group is statistically less than (one-sided $p < 0.025$) the mean in the sham group.

Key secondary outcome(s)

SRT will be considered non-inferior to sham treatment if the lower bound of the 95% confidence interval for the difference in mean change in ETDRS VA at 24 months, between the SRT and sham groups, is no greater than 5 letters.

Completion date

30/06/2024

Eligibility**Key inclusion criteria**

Current participant inclusion criteria as of 01/08/2018:

1. Participants must have neovascular AMD in the study eye, for which they have received at least three prior intravitreal injections of either bevacizumab (Avastin), aflibercept (Eylea), ranibizumab (Lucentis) or pegaptanib (Macugen)
2. Participants must have received an anti-VEGF injection in the study eye within 3 months prior to enrolment
3. Participants must require treatment with anti-VEGF therapy at the time of enrolment, due to OCT evidence of subretinal fluid and/or cystoid macular oedema, and have a macular volume that is greater than a pre-defined threshold that varies for each different make of SD-OCT machine
4. Participants must be at least 50 years of age

Previous participant inclusion criteria:

1. Participants must have neovascular AMD in the study eye, for which they have received at least three prior intravitreal injections of either bevacizumab (Avastin), aflibercept (Eylea), ranibizumab (Lucentis) or pegaptanib (Macugen)
2. Participants must have received an anti-VEGF injection in the study eye within 3 months prior to enrolment
3. Participants must require treatment with anti-VEGF therapy at the time of enrolment, due to OCT evidence of subretinal fluid and/or cystoid macular oedema, and a macular volume that is greater than the 95th percentile of normal for the SD-OCT machines used in the investigational sites
4. Participants must be at least 50 years of age

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 Years

Sex

All

Total final enrolment

411

Key exclusion criteria

Current participant exclusion criteria as of 01/08/2018:

1. Disciform scarring that involves the fovea, in the study eye
2. Visual acuity worse than 6/96 (24 ETDRS letters) in the study eye
3. Lesion size greater than 4 mm in greatest linear dimension, or greater than 2 mm from the centre of the fovea to the furthest point on the lesion perimeter
4. An axial length of less than 20 mm, or greater than 26 mm, in the study eye
5. Contraindication or sensitivity to contact lens application, including recurrent corneal erosions, in the study eye
6. Type 1 or Type 2 diabetes mellitus
7. Retinopathy in the study eye
8. Prior or current therapies in the study eye for age-related macular degeneration, other than anti-VEGF agents, including submacular surgery, subfoveal thermal laser photocoagulation, photodynamic therapy (PDT), or transpupillary thermotherapy (TTT)
9. Presence of an intravitreal device in the study eye
10. Previous radiation therapy to the study eye, head or neck with the exception of radio-iodine treatment for hyperthyroidism, epimacular brachytherapy to the non-study eye, or Oraya SRT to the non-study eye
11. Inadequate pupillary dilation or significant media opacities in the study eye, including cataract, which may interfere with visual acuity testing, the clinical evaluation of the posterior segment, or fundus imaging
12. Study eyes with CNV due to causes other than AMD, including presumed ocular histoplasmosis syndrome (POH), angioid streaks, multifocal choroiditis, choroidal rupture, and pathological myopia (greater than 8 Dioptres spherical equivalent). Participants with retinal angiomatous proliferation (RAP) or idiopathic polypoidal choroidal vasculopathy (IPCV) are not excluded
13. Known allergy to intravenous fluorescein, ICG or intravitreal ranibizumab
14. Intraocular surgery or laser-assisted in situ keratomileusis (LASIK) in the study eye within 12 weeks prior to enrolment
15. Prior pars plana vitrectomy in the study eye
16. Current participation in another interventional clinical trial, or participation in such a clinical trial within the last six months
17. Unwilling, unable, or unlikely to return for scheduled follow-up for the duration of the trial
18. Women who are pregnant at the time of radiotherapy
19. Participants with an implantable cardioverter defibrillator (ICD) or pacemaker implant (or any implanted device) where the device labelling specifically contraindicates patients undergoing X-ray
20. Any other condition, which in the judgment of the investigator, would prevent the participant from granting informed consent or completing the study, such as dementia and mental illness (including generalized anxiety disorder and claustrophobia)

Previous participant exclusion criteria:

1. Disciform scarring that involves the fovea, in the study eye
2. Geographic atrophy that involves the fovea, or an area of geographic atrophy that is more than 500 microns in greatest diameter, immediately adjacent to the fovea, in the study eye
3. Visual acuity worse than 6/96 (24 ETDRS letters) in the study eye
4. Lesion size greater than 4 mm in greatest linear dimension, or greater than 2 mm from the centre of the fovea to the furthest point on the lesion perimeter
5. Distance from the center of the fovea to the nearest edge of the optic disc less than 3 mm in the study eye (this distance is confirmed by the Oraya SRT device software immediately prior to treatment)
6. An axial length of less than 20 mm, or greater than 26 mm, in the study eye
7. Contraindication or sensitivity to contact lens application, including recurrent corneal erosions, in the study eye
8. Type 1 or Type 2 diabetes mellitus
9. Retinopathy in the study eye
10. Prior or current therapies in the study eye for age-related macular degeneration, other than anti-VEGF agents, including submacular surgery, subfoveal thermal laser photocoagulation, photodynamic therapy (PDT), or transpupillary thermotherapy (TTT)
11. Presence of an intravitreal device in the study eye
12. Previous radiation therapy to the study eye, head or neck with the exception of radio-iodine treatment for hyperthyroidism, epimacular brachytherapy to the non-study eye, or Oraya SRT to the non-study eye
13. Inadequate pupillary dilation or significant media opacities in the study eye, including cataract, which may interfere with visual acuity testing, the clinical evaluation of the posterior segment, or fundus imaging
14. Likely to need cataract surgery in the study eye within two years of enrolment
15. Study eyes with CNV due to causes other than AMD, including presumed ocular histoplasmosis syndrome (POH), angioid streaks, multifocal choroiditis, choroidal rupture, and pathological myopia (greater than 8 Dioptres spherical equivalent). Participants with retinal angiomatous proliferation (RAP) or idiopathic polypoidal choroidal vasculopathy (IPCV) are not excluded
16. Known allergy to intravenous fluorescein, ICG or intravitreal ranibizumab
17. Intraocular surgery or laser-assisted in situ keratomileusis (LASIK) in the study eye within 12 weeks prior to enrolment
18. Prior pars plana vitrectomy in the study eye
19. Current participation in another interventional clinical trial, or participation in such a clinical trial within the last six months
20. Unwilling, unable, or unlikely to return for scheduled follow-up for the duration of the trial
21. Women who are pregnant at the time of radiotherapy
22. Participants with an implantable cardioverter defibrillator (ICD) or pacemaker implant (or any implanted device) where the device labelling specifically contraindicates patients undergoing X-ray
23. Any other condition, which in the judgment of the investigator, would prevent the participant from granting informed consent or completing the study, such as dementia and mental illness (including generalized anxiety disorder and claustrophobia)

Date of first enrolment

01/12/2014

Date of final enrolment

31/12/2019

Locations

Countries of recruitment

United Kingdom

England

Study participating centre

King's College Hospital

Ophthalmology Department

Normanby Building

Denmark Hill

London

United Kingdom

SE5 9RS

Study participating centre

University Hospitals Bristol NHS Trust

Bristol Eye Hospital

Lower Maudlin Street

Bristol

United Kingdom

BS1 2LX

Study participating centre

Frimley Park Hospital

Frimley Park Hospital

Portsmouth Road

Frimley

Camberley

United Kingdom

GU16 7UJ

Study participating centre

Maidstone and Tunbridge Wells NHS Trust

The Maidstone Hospital

Hermitage Lane

Maidstone

United Kingdom

ME16 9QQ

Study participating centre
The Royal Wolverhampton NHS Trust
New Cross Hospital
Wolverhampton Road
Heath Town
Wolverhampton
United Kingdom
WV10 0QP

Study participating centre
East Lancashire Hospitals NHS Trust
Royal Blackburn Hospital
Haslingden Road
Blackburn
United Kingdom
BB2 3HH

Study participating centre
The Sussex Eye Hospital
Eastern Road
Brighton
United Kingdom
BN2 5BF

Study participating centre
William Harvey Hospital
Kennington Road
Willesborough
Ashford
United Kingdom
TN24 0LZ

Study participating centre
Hillingdon Hospital
Hillingdon Hospital
Pield Heath Road
Uxbridge
United Kingdom
UB8 3NN

Study participating centre

Whipps Cross University Hospital NHS Trust

Whipps Cross Hospital
Whipps Cross Road
London
United Kingdom
E11 1NR

Study participating centre

Princess Alexandra Hospital

Hamstel Road
Harlow
United Kingdom
CM20 1QX

Study participating centre

Norfolk and Norwich University Hospitals NHS Foundation Trust

Colney Lane
Colney
Norwich
United Kingdom
NR4 7UY

Study participating centre

Essex County Hospital

Lexden Road
Colchester
United Kingdom
CO3 3NB

Study participating centre

Ophthalmology (calderdale Royal Hospital)

The Calderdale Royal Hospital
Huddersfield Road
Halifax
United Kingdom
HX3 0PW

Study participating centre

Stoke Mandeville Hospital

Mandeville Road
Aylesbury

United Kingdom
HP21 8AL

Study participating centre

James Paget University Hospitals NHS Foundation Trust
Lowestoft Road
Gorleston
Great Yarmouth
United Kingdom
NR31 6LA

Study participating centre

Royal Derby Hospital
Uttoxeter Road
Derby
United Kingdom
DE22 3NE

Study participating centre

Queens Hospital
Rom Valley Way
Romford
United Kingdom
RM7 0AG

Study participating centre

Birmingham Heartlands Hospital
Bordesley Green East
Bordesley Green
Birmingham
United Kingdom
B9 5SS

Study participating centre

Central Middlesex Hospital NHS Trust
Acton Lane
Park Royal
London
United Kingdom
NW10 7NS

Study participating centre
Salisbury District Hospital
Odstock Road
Salisbury
United Kingdom
SP2 8BJ

Study participating centre
Cambridge University Hospital
Hills Road
Cambridge
United Kingdom
CB2 0QQ

Study participating centre
St Helier Hospital
Wrythe Lane
Carshalton
United Kingdom
SM5 1AA

Study participating centre
Lincoln County Hospital
Greetwell Road
Lincoln
United Kingdom
LN2 5QY

Study participating centre
Leighton Hospital
Leighton
Crewe
United Kingdom
CW1 4QJ

Study participating centre
Yeovil District Hospital
Higher Kingston
Yeovil

United Kingdom
BA21 4AT

Study participating centre
Dorset County Hospital
Williams Avenue
Dorchester
United Kingdom
DT1 2JY

Study participating centre
Hinchingbrooke Hospital
Hinchingbrooke Park
Huntingdon
United Kingdom
PE29 6NT

Sponsor information

Organisation
King's College Hospital NHS Foundation Trust

ROR
<https://ror.org/01n0k5m85>

Organisation
King's College London

Funder(s)

Funder type
Government

Funder Name
National Institute for Health Research

Alternative Name(s)

National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

The data sharing plans for the current study are unknown and will be made available at a later date.

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		11/06/2024	16/07/2025	Yes	No
Protocol article	protocol	24/11/2016		Yes	No
HRA research summary			28/06/2023	No	No
Protocol file	version 1.9	25/01/2019	01/11/2023	No	No
Study website	Study website	11/11/2025	11/11/2025	No	Yes