

Does a propofol or sevoflurane based protocol has an effect on bowel motility during laparoscopic surgery.

Submission date 03/07/2015	Recruitment status No longer recruiting	☐ Prospectively registered ☐ Protocol
Registration date 19/08/2015	Overall study status Completed	☐ Statistical analysis plan ☒ Results
Last Edited 26/01/2016	Condition category Surgery	☐ Individual participant data

Plain English summary of protocol

Background and study aims:

During laparoscopic surgery, small bowel paralysis facilitates surgical conditions. This is especially important when complex surgery, with intestinal suturing or stapling, is performed. Both opioids and hypnotic agents might influence small bowel peristalsis. The effects of opioids on intestinal motility (intestinal contractions) are well studied and are known to result in intestinal paralysis. With reference to the effects of hypnotics on intestinal motility, data are scarce. Our group recently studied the effect of different volatile (inhaled) anaesthetics on intestinal motility during laparoscopic surgery requiring small bowel anastomosis (removal of a piece of bowel followed by the joining up of the remaining sections). There were statistically significant less peristaltic waves (involuntary muscle movements in the bowel) in the sevoflurane-based anaesthesia group compared to the desflurane-based anaesthesia group. Stimulation of the irritant Transient Receptor Potential channel A1 (TRPA1) was offered to explain the increased intestinal motility observed with desflurane compared to sevoflurane. Given the fact that propofol is a direct modulator of TRPA1, we hypothesized that propofol increases intestinal motility during anaesthesia compared to sevoflurane.

Who can participate?

All patients presenting for gastric bypass surgery at the AZ Groeninge Hospital

What does the study involve?

Patients are randomized in two study groups. One group receives a propofol based anesthesia and one group receives a sevoflurane based anesthesia. At a specific time point during surgery, intestinal motility is assessed by the surgeon and scrub nurse.

What are the possible benefits and risks of participating?

Benefits include an advancement in better understanding of the effects of anesthetics on intestinal motility. No risks are involved in participating in the study, besides the risk involved with anesthesia and surgery.

Where is the study run from?
AZ Groeninge Hospital, Kortrijk (Belgium)

When is the study starting and how long is it expected to run for?
April 2013 to December 2013

Who is funding the study?
AZ Groeninge Hospital (Belgium)

Who is the main contact?
Matthias Desmet

Contact information

Type(s)
Public

Contact name
Dr Matthias Desmet

Contact details
Loofstraat 43
Kortrijk
Belgium
8500

Additional identifiers

Protocol serial number
AZG 2013017

Study information

Scientific Title
The effect of propofol and sevofurane on intestinal motility during laparoscopic surgery: a single blind randomized controlled trial.

Study objectives
Propofol anaesthesia increases intestinal motility during laparoscopic gastric bypass surgery compared to sevoflurane based anaesthesia.

Ethics approval required
Old ethics approval format

Ethics approval(s)
AZ Groeninge Ethical Committee, 04/03/2013, ref: 13006

Study design
Single-centre prospective randomized controlled single-blind two-arm interventional study

Primary study design

Interventional

Study type(s)

Not Specified

Health condition(s) or problem(s) studied

Observation of intestinal motility during laparoscopic surgery

Interventions

2-arm study:

1. Group TIVA (Total Intravenous Anesthesia) will receive a propofol based anesthesia
2. Group Sevo will receive a sevoflurane based anesthesia

Intervention Type

Drug

Phase

Phase IV

Drug/device/biological/vaccine name(s)

1. Propofol
2. Sevoflurane

Primary outcome(s)

Intestinal motility measured by visual count of peristaltic waves

Key secondary outcome(s)

N/A

Completion date

01/06/2014

Eligibility**Key inclusion criteria**

All patients presenting for laparoscopic Roux en Y bariatric surgery at AZ Groeninge Hospital

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Key exclusion criteria

Redo surgery

Date of first enrolment

01/04/2013

Date of final enrolment

31/12/2013

Locations

Countries of recruitment

Belgium

Study participating centre

AZ Groeninge

Loofstraat 43

8500 Kortrijk

Kortrijk

Belgium

8500

Sponsor information

Organisation

Dpt Anesthesia, AZ Groeninge

ROR

<https://ror.org/01cz3wf89>

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

Department of Anesthesia, AZ Groeninge Hospital (Belgium)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary
Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/03/2016		Yes	No