

Manualisation and feasibility study of audiologist-delivered counselling for tinnitus

Submission date 22/03/2016	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 06/05/2016	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 01/11/2021	Condition category Ear, Nose and Throat	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Tinnitus is the experience of noises in the ears or head not related to external sound. It is incurable and is often associated with depression, anxiety, insomnia and reduced quality of life. Tinnitus is treated using a number of interventions. Sound therapy and patient education are widely available but counselling is only available in less than half of audiology departments and there is no agreed standard for what constitutes tinnitus counselling. There is substantial evidence for the clinical benefit of counselling for tinnitus delivered by clinical psychologists or psychiatrists, but there is a lack of evidence for the NHS model of tinnitus care where face-to-face counselling is delivered by audiology professionals. This remains an important unanswered question given that specialist tinnitus care is increasingly becoming the domain of the audiologist. The aim of this study is to assess the feasibility of an audiologist-delivered psychological intervention for tinnitus.

Who can participate?

Adult patients with tinnitus

What does the study involve?

Participants are randomly allocated to one of two groups. One group receives specialised care from an audiologist who has received training in the use of a manual to deliver a psychological intervention for tinnitus treatment. The other group receives usual care from an audiologist who has not trained this specialised training. The researchers evaluate the acceptability of the counselling, patient compliance with therapy, clinician compliance with the manual, willingness of patients to be randomly allocated, and whether there is sufficient interest and need for such a study. Patients and audiologists are interviewed about which elements of counselling are most /least beneficial and why.

What are the possible benefits and risks of participating?

As this stage it is not known whether one treatment is more effective than the other. The researchers cannot promise that the study will help the participants, but participation may help improve the way in which tinnitus is treated in audiology clinics. Psychological treatments are

known to result in improvement in psychological wellbeing for some patients, but a small minority of patients' condition declines from beginning to end of treatment. All participants are given information about a free psychological treatment service they can contact if they wish.

Where is the study run from?

1. Nottingham University Hospitals NHS Trust (UK)
2. Derby Hospitals NHS Foundation Trust (UK)
3. Sherwood Forest Hospitals NHS Foundation Trust (UK)

When is the study starting and how long is it expected to run for?

May 2016 to April 2017

Who is funding the study?

National Institute for Health Research (UK)

Who is the main contact?

1. Dr John Taylor
2. Dr Derek Hoare

Contact information

Type(s)

Public

Contact name

Dr John Taylor

Contact details

NIHR Nottingham Hearing Biomedical Research Unit
Ropewalk House
113 The Ropewalk
Nottingham
United Kingdom
NG1 5DU

Type(s)

Scientific

Contact name

Dr Derek Hoare

Contact details

NIHR Nottingham Hearing Biomedical Research Unit
Ropewalk House
113 The Ropewalk
Nottingham
United Kingdom
NG1 5DU

Additional identifiers

Protocol serial number

PB-PG-0613-31106

Study information

Scientific Title

Manualisation and feasibility study of an audiologist-delivered counselling for tinnitus

Acronym

The TinMan study

Study objectives

Chronic tinnitus is a common incurable condition often associated with depression, anxiety, insomnia and reduced quality of life. Within NHS audiology, tinnitus is treated using a number of recommended interventions but there is no standard protocol for deciding on a first line approach. Sound therapy and patient education are widely available but counselling is only available in less than half of audiology departments and there is no agreed standard for what constitutes tinnitus counselling. There is substantial evidence from systematic reviews for the clinical benefit of counselling for tinnitus delivered by clinical psychologists or psychiatrists, but no studies have sufficiently evidenced the NHS model of tinnitus care where face-to-face counselling is delivered by audiology professionals. This remains an important unanswered question given that specialist tinnitus care is increasingly becoming the domain of the audiologist.

This is a multi-centre feasibility trial of an audiologist-delivered manualised psychological intervention for tinnitus, across three audiology departments where patients are randomised to receive either usual care or care from an audiologist who has received training in the use of the manual.

Ethics approval required

Old ethics approval format

Ethics approval(s)

NHS Health Research Authority: North West - Preston Research Ethics Committee, 04/02/2016, REC ref: 16/NW/0047, IRAS Project ID: 172091

Study design

Multi-centre randomised feasibility trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Tinnitus

Interventions

Patients with tinnitus will be randomised to receive either:

1. Specialised care from an audiologist who has received training in the use of a manual to deliver a psychological intervention for tinnitus treatment
2. Usual care (control) from an audiologist who has not trained this specialised training

Intervention Type

Behavioural

Primary outcome(s)

Feasibility of a full-scale randomised controlled trial is the essential outcome of the feasibility trial. Feasibility would be confirmed by:

1. A conservative recruitment rate of 10% of all eligible patients, calculated using the total number of participation invitation letters sent out
2. Recruitment of 65% of target participants in the 5-month time-frame
3. Retention of 80% of participants with an equivalent proportion of primary outcome data collected
4. Continued compliance to the manualised tinnitus counselling

Key secondary outcome(s)

Secondary patient-reported outcome measures for the feasibility trial include a set of measures that might be typical in a tinnitus clinical trial:

1. Severity and impact of tinnitus, measured using the Tinnitus Functional Index (TFI)
2. General well-being, measured using Clinical Outcomes in Routine Evaluation — Outcome Measure (CORE-OM)
3. Health-related quality of life, measured using Health Utilities Index Mark 3 (HUI3)
4. Overall state of health, measured using EuroQol (EQ-5D)
5. Health resource usage, measured using Client Service Receipt Inventory (CSRI)
6. Quality of the therapeutic alliance, measured using the Working Alliance Inventory (WAI), including both patient and clinician forms

These will be completed at baseline, post-intervention and 6-month follow-up.

Audiologists who have delivered and patients who have received the manualised care will be interviewed about their experience of the intervention and qualitative analysis will be conducted to identify which components worked well, were not useful and why?

Completion date

04/01/2018

Eligibility

Key inclusion criteria

For inclusion in the feasibility trial patients must reflect the typical population that might be expected to take part in a Phase 2 clinical trial. All patients will:

1. Be enrolled at their first visit to audiology
2. Have a primary complaint of tinnitus
3. A clinically relevant need will be determined as a Tinnitus Functional Index (Meikle 2012) questionnaire score >24/100 which indicates tinnitus is at least a moderate problem for the patient
4. Be willing to partake in individual counselling if allocated to the specialist care arm of the study

5. Be willing to complete questionnaires at three time points
6. Take part in follow-up interviews about their experiences (if randomised to intervention)

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Key exclusion criteria

1. Patients whose tinnitus is of a medically treatable origin (i.e. not chronic subjective tinnitus)
2. Patients who are unable to communicate in English

Date of first enrolment

01/07/2016

Date of final enrolment

30/11/2016

Locations**Countries of recruitment**

United Kingdom

England

Study participating centre

Nottingham University Hospitals NHS Trust

United Kingdom

NG7 2UH

Study participating centre

Derby Hospitals NHS Foundation Trust

United Kingdom

DE22 3NE

Study participating centre

Sherwood Forest Hospitals NHS Foundation Trust
United Kingdom
NG17 4JL

Sponsor information

Organisation

Nottingham University Hospitals NHS Trust (UK)

ROR

<https://ror.org/05y3qh794>

Funder(s)

Funder type

Government

Funder Name

National Institute for Health Research

Alternative Name(s)

National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/12/2020	04/03/2021	Yes	No
Protocol article	protocol	15/05/2017		Yes	No
HRA research summary			28/06/2023	No	No
Other publications	Delphi study	01/03/2018	01/11/2021	Yes	No