

# Neuragen® for the relief of neuropathic pain: a randomised, double-blind, placebo controlled clinical trial

|  |  |  |
|--|--|--|
| <b>Submission date</b><br>09/06/2008   | <b>Recruitment status</b><br>No longer recruiting    | <input type="checkbox"/> Prospectively registered    |
| <b>Registration date</b><br>13/06/2008 | <b>Overall study status</b><br>Completed             | <input type="checkbox"/> Protocol                    |
| <b>Last Edited</b><br>25/05/2010       | <b>Condition category</b><br>Nervous System Diseases | <input type="checkbox"/> Statistical analysis plan   |
|  |  | <input checked="" type="checkbox"/> Results          |
|  |  | <input type="checkbox"/> Individual participant data |

**Plain English summary of protocol**  
Not provided at time of registration

## Contact information

**Type(s)**  
Scientific

**Contact name**  
Dr Li Li

**Contact details**  
Department of Kinesiology  
Louisiana State University  
Baton Rouge  
United States of America  
70803

## Additional identifiers

## Study information

**Scientific Title**

### Study objectives

Neuragen® reduces neuropathic pain more and longer than placebo.

**Ethics approval required**

Old ethics approval format

### **Ethics approval(s)**

Ethics approval received from the Institutional Review Board of Louisiana State University as an extension of IRB#2754, approved on 28th September 2007.

### **Study design**

Randomised, double-blind, placebo controlled clinical trial

### **Primary study design**

Interventional

### **Study type(s)**

Treatment

### **Health condition(s) or problem(s) studied**

Peripheral neuropathy

### **Interventions**

Topical application of Neuragen® versus placebo. The medicine was sprayed onto the subjects feet at the sole and on top of the feet. One time application with an 8-hour follow-up.

### **Intervention Type**

Drug

### **Phase**

Not Specified

### **Drug/device/biological/vaccine name(s)**

Neuragen®

### **Primary outcome(s)**

Foot sole pain on 11-point numerical pain scale. 0 - 10 visual analog scale was use to document pain level at 30 minutes before and after the administration of the medication, with 8-hour follow up every hour on the hour.

### **Key secondary outcome(s))**

Duration of pain reduction. 0 - 10 visual analogue scale was use to document pain level at 30 minutes before and after the administration of the medication, with 8-hour follow up every hour on the hour.

### **Completion date**

30/04/2008

## **Eligibility**

### **Key inclusion criteria**

1. Male and female, over 21 years
2. Diagnosed neuropathic pain for more three months
3. Pain level between 3 - 8 on a 0 - 10 visual pain scale
4. Does not have mental and communication impairments

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Adult

**Sex**

All

**Key exclusion criteria**

1. Pregnant
2. Have other types of pain
3. Skin condition
4. Central nerve impairment

**Date of first enrolment**

01/01/2008

**Date of final enrolment**

30/04/2008

**Locations****Countries of recruitment**

United States of America

**Study participating centre****Department of Kinesiology**

Baton Rouge

United States of America

70803

**Sponsor information****Organisation**

Origin BioMed, Inc. (Canada)

ROR

<https://ror.org/008mcnd42>

## Funder(s)

**Funder type**

Industry

**Funder Name**

Origin BioMed, Inc. (Canada)

## Results and Publications

Individual participant data (IPD) sharing plan

**IPD sharing plan summary**

Not provided at time of registration

### Study outputs

| Output type                     | Details | Date created | Date added | Peer reviewed? | Patient-facing? |
|---------------------------------|---------|--------------|------------|----------------|-----------------|
| <a href="#">Results article</a> | results | 20/05/2010   |            | Yes            | No              |