

Take home naloxone intervention in multicentre emergency settings

Submission date 09/02/2018	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered
		<input checked="" type="checkbox"/> Protocol
Registration date 16/02/2018	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
Last Edited 03/03/2026	Condition category Injury, Occupational Diseases, Poisoning	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

People who take opioid drugs such as heroin can overdose. The number of people who die this way is increasing, with tragic consequences for families, friends and communities. Naloxone is a drug that can reverse the effects of opioid overdose. Emergency ambulance staff and doctors in the Emergency Department regularly give drug users naloxone, but some people die before they reach emergency medical services. There are schemes in the UK and internationally where naloxone is given to drug users to give to others in an emergency. This is called 'Take Home Naloxone' (THN). It is not known whether THN saves lives. There are some concerns that it could encourage risk-taking behaviour, with the drug user feeling that they have a 'safety net' and can take a higher dose. The aim of this study is to see whether it is possible for ambulance paramedics and Emergency Department staff to give out THN kits to drug users they see, and to see if it is possible to collect data to find out whether it reduces deaths from overdose.

Who can participate?

Adult (aged 18 or over) opioid users at risk of overdose who are attended by emergency ambulance paramedic or who attend ED, and their accompanying friends, relatives or carers

What does the study involve?

This study is carried out in two areas where THN is given to patients who have overdosed or who are at risk of overdose, and two other areas where THN kits are not given out (treatment as usual). It is very difficult to follow up these patients because they don't respond to phone calls or may have no fixed address, therefore this study follows what happens to patients using the routine information which health services already collect about everyone they see. Information is collected about deaths, overdoses, emergency ambulance calls and emergency department attendances and admissions up to 1 year after the patients are seen. These figures are compared between the areas which give out THN and the areas which do not. Interviews are also carried out to find out about the experiences and views of patients receiving THN and staff who give out the kits.

What are the possible benefits and risks of participating?

If this study shows it is possible to give out THN kits through emergency services, and that data can be collected about the effects, a larger study will be carried out to find out whether THN

reduces deaths. THN is known to have the potential to save lives on an individual basis, but on an aggregate level the benefits and harms are unknown, which is the reason for this study. There are no participant incentives on offer due to the study design.

Where is the study run from?

1. Bristol Royal Infirmary (UK)
2. Hull Royal Infirmary (UK)
3. Northern General Hospital Sheffield (UK)
4. Wrexham Maelor Hospital (UK)

When is the study starting and how long is it expected to run for?

March 2018 to July 2022 (updated 19/07/2021, previously: November 2020)

Who is funding the study?

NIHR Health Technology Assessment Programme (UK)

Who is the main contact?

Jenna Jones, j.k.jones@swansea.ac.uk

Contact information

Type(s)

Public

Contact name

Ms Jenna Jones

ORCID ID

<https://orcid.org/0000-0002-1280-4941>

Contact details

Floor 2, ILS2
Swansea University
Swansea
United Kingdom
SA2 8PP
+44 (0)1792513424
j.k.jones@swansea.ac.uk

Type(s)

Scientific

Contact name

Prof Helen Snooks

Contact details

Floor 2, ILS2
Singleton Park Campus
Swansea
United Kingdom
SA28PP

Additional identifiers

Protocol serial number

HTA 16/91/04

Study information

Scientific Title

Take home naloxone Intervention Multicentre Emergency setting feasibility trial

Acronym

TIME

Study objectives

That it is feasible to conduct a fully powered randomised controlled trial to measure the clinical and cost effectiveness of take home naloxone distributed from emergency care settings using a cluster design and linked anonymised routine data.

Ethics approval required

Old ethics approval format

Ethics approval(s)

1. Approved 16/10/2018, NHS HRA Wales REC 4 (Health and Care Research Wales Support Centre, Castlebridge 4, 15-19 Cowbridge Road East, Cardiff, CF11 9AB; +44(0)7976 982591; tracy.biggs@wales.nhs.uk), ref: 18/WA/0337
2. Approved 07/01/2019, NHS HRA CRA (Skipton House, 80 London Road, London, SE1 6LH; +44 (0)20 7972 2557; hra.cag@nhs.net), ref: 18/CAG/0176

Study design

Feasibility study for a randomized controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Opioid overdose

Interventions

This is a feasibility study for an RCT of Take Home Naloxone (THN) in emergency settings clustered by emergency department (ED) catchment area and local ambulance service. Anonymised linked data outcomes will be captured for trial participants and their peers, and aggregate outcomes at general population level. The trialists will also collect qualitative data to examine implementation, patient safety and experience.

The four sites to be randomly allocated to study arms are:

Bristol: Bristol Royal Infirmary & associated ambulance service catchment area

Hull: Hull Royal Infirmary & associated ambulance service catchment area

Sheffield: Northern General Hospital Sheffield & associated ambulance service catchment area

Wrexham: Wrexham Maelor Hospital & associated ambulance service catchment area

It was agreed that the randomisation would be stratified to ensure that the two sites served by the Yorkshire Ambulance Service would be in different study arms; this reduced the randomisation options to four, as listed below:

Option A:

Intervention: Wrexham, Sheffield

Control: Bristol, Hull

Option B:

Intervention: Bristol, Hull

Control: Wrexham, Sheffield

Option C:

Intervention: Bristol, Sheffield

Control: Wrexham, Hull

Option D:

Intervention: Wrexham, Hull

Control: Bristol, Sheffield

Four identical opaque envelopes, each containing one option, were prepared, presented to a person not involved in their preparation, and one selected at random. The chosen envelope contained Option B, so that the proposed allocation is:

Intervention: Bristol, Hull

Control: Wrexham, Sheffield

The THN intervention will consist of training for participating paramedics and ED staff; a protocol for the supply of THN kits to those eligible for the intervention; a single use THN kit (ampoule of Naloxone Hydrochloride 400 micrograms/ml solution for intramuscular injection and syringe; and instructions on how to correctly administer the naloxone dose and provide basic life support in event of overdose). The training provided to recipients of THN kits by paramedic or ED staff will include preparation and administration of the naloxone, providing appropriate aftercare including calling 999, and resuscitation techniques. The trialists will consult with patient and public interest (PPI) representatives and a wider group of service users, addiction specialists, paramedics, and ED staff to assess acceptability, and identify advantages and disadvantages of our proposed THN kit. THN will be distributed in the intervention sites for a period of 12 months.

Control sites will operate treatment as usual (TAU).

The study will be carried out in the prehospital environment and at a receiving ED at each site. The prehospital environment will be the geographic catchment area for the receiving ED. Paramedics and clinical ED staff at study sites will be invited to take part.

The target populations eligible for the intervention (THN) will be: opioid users at high risk of fatal overdose who are attended by emergency ambulance paramedic or who attend ED; their accompanying friends, relatives or carers; and people at high risk of fatal overdose who make up a wider peer group of opioid users coming in to contact with emergency ambulance paramedics or ED.

The trialists propose to develop and test two related methods of defining these populations:

1. A discriminant function incorporating routinely recorded risk predictors used in risk indices for opioid-related events. These will be derived from linked opioid-related mortality (from ONS), ED and inpatient data. They have already obtained linked data to study the characteristics of decedents of opioid overdose in Wales during 2015, and they will extend this data set to cover those with ED and inpatient events associated with non-fatal opioid overdoses, and those with no such events. Although discriminant analysis is inherently multivariate in nature, they will also assess whether the discriminant function can be reduced to a single individual-level risk score; consider thresholds used in its definition; and evaluate its performance.
2. The discriminant analysis approach outlined will include recent ED and inpatient data related to opioid overdose and associated events, for which current coding is known to be of variable quality or low specificity. The trialists will therefore test whether the planned Emergency Care DataSet (ECDS) provides reliable data about attendances for opioid overdose and related problems that could be used to improve the ability to identify both the patients eligible for THN provision and the wider peer group.

Intervention Type

Mixed

Primary outcome(s)

Mortality (total and opioid overdose related), collected using anonymised data linkage at a single time point 12 months post intervention

Key secondary outcome(s)

Collected using anonymised data linkage at a single time point 12 months post intervention:

1. Emergency admissions
2. ITU admissions
3. ED attendances (total and opioid overdose related)
4. 999 attendances (total and opioid overdose related)
5. THN kits issued
6. NHS resource usage

Completion date

31/07/2022

Eligibility

Key inclusion criteria

Eligible for intervention:

Adult (18 years of age or over, male or female) opioid users at risk of overdose who are attended by emergency ambulance paramedic or who attend ED

Accompanying friends, relatives or carers (18 years of age or over, male or female) of opioid users at risk of overdose who are attended by emergency ambulance paramedic or who attend ED

Eligible for inclusion in outcome follow up and analyses:

High risk peers (18 years of age or over, male or female) - opioid users at high risk of fatal opioid overdose who make up a wider peer group who may or may not make contact with emergency services at study site during the study period

Healthy volunteers allowed

No

Age group

Mixed

Lower age limit

18 years

Upper age limit

100 years

Sex

All

Total final enrolment

0

Key exclusion criteria

1. Under 18 years of age
2. Actively dissent from involvement in study
3. Non-opioid overdose

Date of first enrolment

01/02/2019

Date of final enrolment

01/02/2020

Locations**Countries of recruitment**

United Kingdom

England

Wales

Study participating centre

Yorkshire Ambulance Service NHS Trust

Springhill

2 Brindley Way

Wakefield 41 Industrial Estate

Wakefield

England

WF2 0XQ

Study participating centre
South Western Ambulance Service NHS Foundation Trust
Abbey Court
Eagle Way
Sowton Industrial Estate
Exeter
England
EX2 7HY

Study participating centre
Welsh Ambulance Services NHS Trust
Unit 7
Ffordd Richard Davies
St Asaph Business Park
St. Asaph
Wales
LL17 0LJ

Study participating centre
Northern General Hospital
Northern General Hospital NHS Trust
C Floor, Huntsman Building
Herries Road
Sheffield
England
S5 7AU

Study participating centre
Hull Royal Infirmary
Anlaby Road
Hull
England
HU3 2JZ

Study participating centre
Bristol Royal Infirmary
Marlborough Street
Bristol
England
BS2 8HW

Study participating centre
Wrexham Maelor Hospital
Croesnewydd Road
Wrexham Technology Park
Wrexham
England
LL13 7TD

Sponsor information

Organisation

Swansea University

ROR

<https://ror.org/053fq8t95>

Funder(s)

Funder type

Government

Funder Name

Health Technology Assessment Programme

Alternative Name(s)

NIHR Health Technology Assessment Programme, Health Technology Assessment (HTA), HTA

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

Quantitative outcome data will be stored in the SAIL databank and accessed via the secure sail gateway (<https://saildatabank.com/>). Research team members (myself and Dr Alan Watkins the trial statistician) have access to the gateway via yubikey (<https://www.yubico.com/>). The trialists

are able to access the data but cannot take data outside of the gateway. Requests for exporting linked data in to the gateway are reviewed by internal and then external reviewers prior to being granted. Data will be linked and anonymised prior to export in to the gateway by NHS Digital and NWIS (for Welsh control site).

IPD sharing plan summary

Available on request, Stored in non-publicly available repository

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		29/08/2024	02/09/2024	Yes	No
Results article		31/10/2024	04/11/2024	Yes	No
Protocol article		09/07/2020	03/03/2026	Yes	No
HRA research summary			26/07/2023	No	No
Poster results		01/01/2019	01/08/2019	No	No