

The effect of starting dialysis earlier than usual in patients with damaged kidneys

Submission date 03/02/2019	Recruitment status No longer recruiting	☐ Prospectively registered ☒ Protocol
Registration date 18/02/2019	Overall study status Completed	☐ Statistical analysis plan ☒ Results
Last Edited 07/08/2023	Condition category Urological and Genital Diseases	☐ Individual participant data

Plain English summary of protocol

Background and study aims

More than 1 in 2 critically ill patients in the intensive care unit (ICU) develop acute kidney injury (AKI). A proportion of patients need dialysis treatment, also known as replacement therapy (RRT). The aim of RRT is to remove toxins and excess fluid which may have accumulated because of kidney failure. At present it is not clear whether it is best to start RRT only when patients have evidence of severe AKI or whether it is better to initiate RRT earlier, i.e. as soon as patients have signs of impaired kidney function.

Who can participate?

Patients aged over 18 years admitted to the intensive care unit with serious kidney problems.

What does the study involve?

RRT consists of having a catheter (a tube or 'line') in a vein in the neck or in the groin through which blood flows to a dialysis machine where it is cleansed and excess water is removed. The cleansed blood is then returned to the patient via the catheter.

What are the possible benefits and risks of participating?

RRT can be lifesaving but has side effects, possibly including bleeding and bloodstream infections. Importantly, if at any stage the treating clinician feels that RRT is mandatory or alternatively should not be started, their judgement will override. Also, there will be no change in any clinical management and no extra blood samples will be taken.

Where is the study run from?

Guy's & St Thomas Hospital, Westminster Bridge Road, London, SE1 9RT, UK.

When is the study starting and how long is it expected to run for?

February 2018 to March 2020.

Who is funding the study?

The National Institute for Health Research (United Kingdom) Health Technology Assessment programme.

Who is the main contact?

Mrs. Marlies Ostermann, Marlies.Ostermann@gstt.nhs.uk

Contact information

Type(s)

Scientific

Contact name

Mrs Marlies Ostermann

Contact details

Guy's & St Thomas Hospital

Department of Critical Care

Westminster Bridge Road

London

United Kingdom

SE1 9RT

+44 (0)2071883038

Marlies.Ostermann@gstt.nhs.uk

Additional identifiers

ClinicalTrials.gov (NCT)

NCT02568722

Integrated Research Application System (IRAS)

191390

Study information

Scientific Title

STandard versus Accelerated initiation of Renal Replacement Therapy in Acute Kidney Injury (STARRT-AKI): a multi-centre, randomized, controlled trial

Acronym

STARRT-AKI

Study objectives

In critically ill patients with severe acute kidney injury (AKI), randomization to accelerated initiation of renal replacement therapy (RRT), compared to a conservative strategy consistent with standard care, leads to:

1. improved survival at 90 days; and
2. better recovery of kidney function, defined as independence from dialysis at 90 days.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 22/05/2017, Camberwell St Giles REC (Institute of Psychiatry, Seminar Room 6, IOP Main Building, SE5 8AF, +44 (0) 207 1048044, NRESCCommittee.London-CamberwellStGiles@nhs.net), ref:

Primary study design

Interventional

Study design

Randomised controlled trial

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Acute kidney injury

Interventions

Patients in the ICU with early AKI will be randomised to RRT using standard criteria versus accelerated (earlier) RRT. Patients randomised to the accelerated arm will be started on RRT within 12 hours of eligibility. In patients randomised to standard initiation, RRT will be discouraged until serum potassium >5.9 mmol/L, pH <7.21 or serum bicarbonate <13 mmol/L, respiratory failure due to fluid overload, or persistent AKI stage 2 or 3 for >72 hours.

Intervention Type

Procedure/Surgery

Primary outcome(s)

Mortality measured at 90 days

Key secondary outcome(s)

Recovery of kidney function defined as independence from dialysis at 90 days

Completion date

31/05/2021

Eligibility**Key inclusion criteria**

1. Age ≥ 18 years
2. Admission to an intensive care unit (ICU)
3. Evidence of kidney dysfunction [serum creatinine ≥ 100 $\mu\text{mol/L}$ in women and ≥ 130 $\mu\text{mol/L}$ in men]
4. Evidence of severe AKI

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 Years

Sex

All

Total final enrolment

2927

Key exclusion criteria

1. Lack of commitment to provide RRT as part of limitation of ongoing life support.
2. Presence of a drug overdose that necessitates initiation of RRT.
3. Any RRT within the previous 2 months.
4. Kidney transplant within the past 365 days.
5. Known pre-hospitalization advanced chronic kidney disease, defined by an estimated glomerular filtration rate $<20 \text{ mL/min/1.73 m}^2$ in a patient who is not on chronic dialysis.
6. Presence or clinical suspicion of renal obstruction, rapidly progressive glomerulonephritis, vasculitis, thrombotic microangiopathy (eg, thrombotic thrombocytopenic purpura, hemolytic uremic syndrome, malignant hypertension, scleroderma renal crisis) or acute interstitial nephritis.
7. Likelihood that an absolute indication for RRT will arise in the subsequent 24 hours based on the most recent blood work for the following parameters: serum K >5.5
8. Likelihood that an absolute indication for RRT will arise in the subsequent 24 hours based on the most recent blood work for the following parameters: serum bicarbonate $<15 \text{ mmol/L}$

Date of first enrolment

01/02/2018

Date of final enrolment

31/05/2020

Locations**Countries of recruitment**

United Kingdom

England

Canada

Study participating centre

Guy's & St Thomas Hospital

Westminster Bridge Road

London

United Kingdom

SE1 9RT

Sponsor information

Organisation

University of Toronto

ROR

<https://ror.org/03dbr7087>

Funder(s)

Funder type

Government

Funder Name

Health Technology Assessment Programme

Alternative Name(s)

NIHR Health Technology Assessment Programme, Health Technology Assessment (HTA), HTA

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

Not provided at time of registration

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		16/07/2020	18/11/2022	Yes	No
Protocol file	version 3.3	15/12/2020	07/08/2023	No	No