

Effect of immunitum supplementation on prevention of recurrent infectious episodes in children with moderate malnutrition: a randomised, double-blind, placebo-controlled trial

Submission date 06/05/2017	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 04/07/2017	Overall study status Completed	<input type="checkbox"/> Protocol
Last Edited 01/04/2019	Condition category Infections and Infestations	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims:

Improving child survival remains a significant challenge in low-income countries. The leading causes of death among children younger than five years of age are still preventable conditions including malaria, sudden chest infections and diarrheal diseases. In addition, malnutrition is still very common in low-income countries. It accounts for nearly half of the burden of under-five child death mainly through increasing vulnerability to infections. The aim of this study is to evaluate the effect of immunitum (an immune system booster micronutrient supplement) in preventing repeated infectious episodes in children with moderate malnutrition.

Who can participate?

Children who are aged 12 to 59 months, have moderate malnutrition and a history of recurrent chest and airways, and digestive system infections

What does the study involve?

Participants undergo a medical examination to determine their eligibility. They are then randomly assigned to one of two groups. Participants in the first group receive a capsule of immunitum to take by mouth twice daily. Those in the second group receive a capsule of a placebo (dummy drug) to take by mouth twice daily. Participants are followed up monthly for 12 months. The follow up include clinical examination, assessment of growth (weight and height) and general vitality. Blood and stool samples are taken at the start of the study and then during the sixth and twelfth months of the follow up.

What are the possible benefits and risks of participating?

Participants may benefit from the expected improvement in immune system and decrease in the number and severity of infectious episodes common in children with malnutrition. The risk to participants is minimal to none. There are no expected side effects as a result of taking the

supplement. However, participants are checked at each follow up and, in between, they are encouraged to report any unusual signs/symptoms.

Where is the study run from?

1. Gakurazo Health Center (Rwanda)
2. Kamabuye Health Center (Rwanda)
3. Mayange Health Center (Rwanda)

When is study starting and how long is it expected to run for?

March 2012 to October 2013

Who is the main contact?

Dr Lisine Tuyisenge
tuyislisine@gmail.com

Contact information

Type(s)

Scientific

Contact name

Dr Lisine Tuyisenge

Contact details

University Teaching Hospital of Kigali (CHUK)
KN 4 Ave
Kigali
Rwanda
655
+250 788 411 764
tuyislisine@gmail.com

Type(s)

Scientific

Contact name

Prof Walter Wahli

Contact details

Chemin du Baillon 12
Échichens
Switzerland
1112

Type(s)

Scientific

Contact name

Dr Celestin Hategeka

ORCID ID

<https://orcid.org/0000-0001-7808-4652>

Contact details

University of British Columbia
2206 East Mall
Vancouver
Canada
V6T1Z3
+1 778 990 7941
celestine.hategeka@alumni.ubc.ca

Additional identifiers

Protocol serial number

RPA0001

Study information

Scientific Title

Effect of immunitum supplementation on prevention of recurrent infectious episodes in children (aged 12 - 59 months) with moderate malnutrition: a randomised, double-blind, placebo-controlled trial

Study objectives

Daily supplementation of immunitum reduces infectious episodes in children with moderate malnutrition.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Rwanda National Ethic Committee (RNEC), ref: 158/RNEC/2011

Primary study design

Interventional

Study design

Randomised double-blind placebo-controlled trial

Study type(s)

Prevention

Health condition(s) or problem(s) studied

Recurrent infectious diseases in children with malnutrition:

1. ENT sphere infections
2. Broncho-pulmonary infections
3. Gastrointestinal tract infections
4. Malaria

Interventions

Participants are randomly allocated to one of two groups using a 1:1 randomisation at the time of enrolment to allocate five participants in each arm for every round of ten enrollments. The study is double-blind, with the study and control products being indistinguishable in shape, size, colour and taste. They are pre-labeled for the entire study by a third-party person who does not play a role in the study implementation.

Group 1: Participants in this group receive an oral capsule of immunitum twice daily.

Group 2: Participants receive the placebo oral capsule which is taken twice daily.

Participants in both groups receive the treatment for 12 months. Clinical follow up and study visits are the same for both groups, with the study product given at the time of enrollment and randomisation and then participants are seen monthly for a clinical follow-up, nutritional status and treatment compliance until month 12.

Intervention Type

Supplement

Primary outcome(s)

1. Incidence of infectious diseases (e.g., ENT sphere infections, broncho-pulmonary infections, gastrointestinal tract infections, and malaria) common in children (12-59 months) with moderate malnutrition as defined by WHO is assessed by a trained general medical doctor with experience in pediatric case management in district hospitals (supervised daily and closely by an experienced pediatrician) Each enrolled participant was discussed together at her arrival at health center, on regular appointments, and every time a family called saying that the child has a health problem. In addition, when recruited children arrived to health centers, they were seen by a trained nurse working regularly in PCIME or IMCI (Integrated Management of Childhood Illness), who records the case in a dedicated register and reported the case immediately on phone when urgent or complicated situations. Otherwise the case was reported to the study team during the following monthly visit.

2. Severity and length (in days) of each infectious episode is measured by the medical doctor who is assigned to see participants from the beginning of the disease and follow participant till the end of episode of the disease. Symptoms notified at the beginning are verified every day by a community health worker who see the participant twice a day for giving the immunitum /placebo and asks questions about the signs and symptoms of the disease at each visit then records in the trial notebook which is presented to the medical doctor at the regular visit at the health center or for severe cases at the district hospital, when the child is hospitalized.

Key secondary outcome(s)

1. Weight and height are measured using standard anthropomorphic techniques at health center during each monthly visit for 12 months

2. General vitality is assessed using the general vitality questionnaire that measures apathy, sleep, feeding and fatigue; at health center during each monthly visit for 12 months

3. Tolerance and compliance is assessed by the assigned community health worker at home, and then by the study team at health center during each monthly visit for 12 months

4. Proteinemia is measured using spectrophotometry, transferrin is measured by turbidimetry, CRP is measured by ELISA, blood sugar is measured by spectrometry, hemoglobin is measured by spectrometry, erythrocyte sedimentation rate, RBC, WBC, thick blood smear, HIV, stool (for parasites and culture when child has diarrhea, one of the stool sample taken at 12 months will be analysed for microbiome composition) are measured using sample analysis by specific lab techniques for each lab exam and following well defined lab protocols at baseline, six and 12

months

5. Levels of vitamins A, B1, B2, B9, B12, C, D, E, PP are measured using high-performance liquid chromatography at baseline, six and 12 months

6. Levels of iodine will be determined by mass spectrometry, sodium, potassium and chlorine by ion-selective electrode (ISE), selenium by atomic absorption spectroscopy, zinc by atomic absorption spectrophotometry, and iron by spectrophotometric assay at baseline, six and 12 months

7. Genomic analyses will be done by single nucleotide polymorphism and proteomics by electrophoresis at baseline, six and 12 months

Completion date

18/10/2013

Eligibility

Key inclusion criteria

1. Children aged 12 to 59 months
2. Moderate malnutrition as per WHO criteria
3. History of recurrent infectious episodes of upper and lower respiratory tract and gastrointestinal tract

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Child

Lower age limit

12 Months

Upper age limit

59 Months

Sex

All

Key exclusion criteria

1. Children aged less than 12 months and 60 months and over
2. Suffering from severe malnutrition (MUAC<115 mm et W/A \leq - 3 Z Score et H/A \leq - 3 Z Score)
3. Suffering from a progressive disease (Tuberculosis, HIV)

Date of first enrolment

24/09/2012

Date of final enrolment

18/10/2012

Locations

Countries of recruitment

Rwanda

Study participating centre

Mayange Health Center

Bugesera

Rwanda

c/o 01

Study participating centre

Kamabuye Health Center

Bugesera

Rwanda

c/o 01

Study participating centre

Gakurazo Health Center

Bugesera

Rwanda

c/o 01

Sponsor information

Organisation

Foundation Actigenomics

Funder(s)

Funder type

Industry

Funder Name

Actigenomics SA

Funder Name

Africa Alive

Funder Name

Sucafina

Funder Name

Nutrigenomics

Funder Name

Renta 4

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study is not expected to be made available due to access being required by the Rwanda National Ethics Committee (RNEC) and by the principal investigator Dr. Lisine Tuyisenge tuyisline@gmail.com.

IPD sharing plan summary

Not expected to be made available

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Participant information sheet	version V2	03/07/2017	01/04/2019	No	Yes