ISRCTN13849110 https://doi.org/10.1186/ISRCTN13849110

Can we safely reduce the number of days of radiotherapy needed to treat people with breast cancer who need boost treatment?

Submission date 19/02/2025	**Recruitment status** Recruiting	[X] Prospectively registered [_] Protocol [_] Statistical analysis plan [_] Results [_] Individual participant data [_] Record updated in last year
Registration date 03/03/2025	**Overall study status** Ongoing	
Last Edited 03/03/2025	**Condition category** Cancer	

Plain English summary of protocol

Background and study aims
Each year, 37,000 people in the UK with breast cancer receive radiotherapy, which uses radiation to kill cancer cells. Most people can now be treated over 5 days, but about 10,000 still need up to 23 days of treatment because they require an extra dose called a boost. This study aims to find out if the boost can be given within a 5-day radiotherapy course, comparing two different boost doses over 5 days with the standard 15-day boost dose.

Who can participate?
People with breast cancer who need a boost dose as part of their radiotherapy treatment can participate. Participants will be invited from 50 UK radiotherapy centres.

What does the study involve?
Participants will be randomly placed into one of three groups by a computer:
One group will receive the boost dose over 15 days.
Two groups will receive the boost dose over 5 days, with different boost doses for each group.
Participants will provide information about side effects, changes to the treated breast, extreme tiredness, and quality of life for five years.
What are the possible benefits and risks of participating? The study aims to show that the 5-day treatment is as effective as the 15-day treatment in preventing cancer from returning, with the same or fewer side effects and faster recovery. Risks include potential side effects from radiotherapy and the possibility that the shorter treatment may not be as effective.

Where is the study run from?
Institute of Cancer Research - Clinical Trials and Statistics Unit (UK).

When is the study starting and how long is it expected to run for?
April 2025 to September 2033

Who is funding the study?
The study is funded by the National Institute for Health and Care Research - Health Technology Assessment (NIHR-HTA) programme (UK).

Who is the main contact?
Institute of Cancer Research - Clinical Trials and Statistics Unit
fastforwardboost-icrctsu@icr.ac.uk

Contact information

Type(s)
Scientific

Contact name
Prof Judith Bliss

Contact details
ICR-CTSU,
Sir Richard Doll Building,
The Institute of Cancer Research,
Cotswold Road
Sutton
United Kingdom
SM2 5NG
+44 208 722 4104
Fastforwardboost-icrctsu@icr.ac.uk

Type(s)
Principal investigator

Contact name
Dr Anna Kirby

ORCID ID
https://orcid.org/0000-0002-5528-1669

Contact details
Royal Marsden Hospital
Downs Road
Sutton
United Kingdom
SM2 5PT
+44 2087224104
anna.kirby@rmh.nhs.uk

Type(s)
Public

Contact name
Mr Mark Sydenham

ORCID ID
https://orcid.org/0000-0002-9157-9710

Contact details
ICR-CTSU,
Sir Richard Doll Building,
The Institute of Cancer Research,
Cotswold Road
Sutton
United Kingdom
SM2 5NG
+44 208 722 4104
fastforwardboost-icrctsu@icr.ac.uk

Additional identifiers

Integrated Research Application System (IRAS)
341881

Central Portfolio Management System (CPMS)
57885

National Institute for Health and Care Research (NIHR)
157800

Protocol serial number
ICR_CTSU/2024/10088

Study information

Scientific Title
A randomised clinical trial testing a 1-week schedule of curative simultaneous integrated boost radiotherapy against a standard 3-week schedule in patients with early breast cancer

Acronym
FAST-Forward Boost

Study objectives
Local recurrence rates at 5-years will be no higher with appropriately dosed 1-week Simultaneous Integrated Boost radiotherapy (SIB RT) than with 3-week SIB radiotherapy and that this can be achieved without an increase in normal tissue side-effects

Ethics approval required
Ethics approval required

Ethics approval(s)
Approved 14/01/2025, London - South East REC (2 Redman Place, Stratford, London, E20 1JQ, United Kingdom; +44 207 104 8222; londonsoutheast.rec@hra.nhs.uk), ref: 24/LO/0910

Primary study design

Interventional

Study design
Phase III multi-centre randomized controlled trial

Study type(s)
Treatment

Health condition(s) or problem(s) studied
Breast cancer

Interventions
Patients will be treated using standard radiotherapy to the breast +/- nodes with a SIB to the tumour bed and randomised on a 1:1:1 basis to one of the following schedules:
• Standard radiotherapy to the breast +/- nodes using a schedule of 40Gy/15Fr over 3 weeks with a 48Gy/15Fr simultaneous integrated boost (SIB) of the tumour bed (Control Group)
• 26Gy/5Fr over 1 week with a 31Gy/5Fr SIB (Test Group 1)
• 26Gy/5Fr over 1 week with a 30Gy/5Fr SIB (Test Group 2)

Radiotherapy treatment
All patients will attend the radiotherapy department for a planning CT scan so their radiotherapy treatment can be planned.
Once the treatment has been planned. The patient will start their radiotherapy treatment. Treatment is usually given daily on week days. Those in the control group will have their radiotherapy treatment in 15 treatments, the test groups will have their radiotherapy in 5 treatments.

Questionnaires
For those taking part in the early side-effects sub-study, questionnaires about side-effects will need to be completed by the patient prior to radiotherapy, weekly for 7 weeks from start of radiotherapy and at 3 months.
All patients will be asked to complete questionnaire booklets pre treatment and at weeks 1, 3 and 5 from the start of radiotherapy then at 3 months, 1, 2, 3, 4 and 5 years.

Follow up
For those taking part in the early side-effects sub-study:
Control group: will also be reviewed at weeks 1, 3, 5 and 7 from the start of radiotherapy
Test Groups: will also be reviewed at weeks 1, 3, 5 and 7 from the start of radiotherapy
All patients will be reviewed by the clinical team at 3 months, 1 years, 3 years and at 5 years.

Intervention Type
Procedure/Surgery

Primary outcome(s)
Ipsilateral breast tumour recurrence (IBTR) measured using patient records at 5 years

Key secondary outcome(s)
1. Patient-reported acute radiotherapy adverse effects, with a focus on skin, breast, and oesophageal effects, are measured using PRO-CTCAE and trial-specific questionnaires at baseline, weekly for 7 weeks from the start of radiotherapy, and at 3 months
2. Patient-reported late effects on quality of life, with a focus on breast symptoms and shoulder

/arm functioning, are measured using EORTC QLQ BR-23 and trial-specific questionnaires at baseline, weeks 1, 3, and 5 from the start of radiotherapy, and at 3 months, 1, 2, 3, 4, and 5 years
3. Patient-reported fatigue is measured using EORTC QLQ FA-12 at baseline, weeks 1, 3, and 5 from the start of radiotherapy, and at 3 months, 1, 2, 3, 4, and 5 years
4. Health-related quality of life is measured using EORTC QLQ C-30 and EQ5D-5L at baseline, weeks 1, 3, and 5 from the start of radiotherapy, and at 3 months, 1, 2, 3, 4, and 5 years
5. Body image is measured using the Body Image Scale (BIS) at baseline, weeks 1, 3, and 5 from the start of radiotherapy, and at 3 months, 1, 2, 3, 4, and 5 years
6. Clinician-reported acute radiotherapy adverse effects, with a focus on skin, oesophageal, and lung toxicity, are measured using CTCAE v5.0 and RTOG at baseline, weekly for 7 weeks from the start of radiotherapy, and at 3 months
7. Clinician-reported breast oedema is measured using trial-specific tools at baseline, weekly for 7 weeks from the start of radiotherapy, and at 3 months
8. Clinician-reported late radiotherapy adverse effects, with a focus on normal tissue effects and cosmesis, are measured using tools developed in previous breast radiotherapy trials and the Harvard-Harris scale at baseline, 3 months, 1, 3, and 5 years
9. Clinician-reported lung toxicity is measured using RTOG at baseline, 3 months, 1, 3, and 5 years
10. Recurrence-free survival, breast cancer-related survival, and overall survival are measured using NHS routinely collected data at baseline, 3 months, 1, 3, and 5 years
11. Cost-effectiveness is measured using health economic analysis incorporating data on patient-reported health resource use, out-of-pocket expenses, and health status (EQ5D-5L) at baseline, 3 months, 1, 3, and 5 years

Completion date
30/09/2033

Eligibility

Key inclusion criteria
1. Age >=18 years
2. Histologically confirmed breast cancer (T1-T3, N0-3, M0) (multifocal disease is allowed) requiring a tumour bed boost plus whole breast radiotherapy +/- radiotherapy to nodes* (axilla +/- internal mammary chain) or DCIS (Tis, N0-3, M0) requiring a tumour bed boost according to local centre policy
3. Treated with breast conservation surgery
4. Complete microscopic resection (invasive cancer and/or DCIS clear of ink on radial margins or, if at margin, surgeon confirms no further breast tissue to excise)
5. Patient can provide informed consent
* Axilla levels as per MDM recommendation
NB Patients with synchronous bilateral breast cancer can be included as long as the disease on at least one side fulfils the inclusion criteria above. Where the patient has synchronous bilateral disease and needs a tumour bed boost on both sides, both sides will need to fulfil the inclusion criteria for the patient to be eligible for the trial

Participant type(s)
Patient

Healthy volunteers allowed
No

Age group

Adult

Lower age limit
18 Years

Sex
Female

Key exclusion criteria
1. Treated with ipsilateral mastectomy
2. Previous radiotherapy to ipsilateral chest area that precludes delivery of a radical dose of adjuvant radiotherapy to the breast with tumour bed boost. NB for any scenarios where there is overlap with previous radiotherapy approval must be sought from the FAST-Forward Boost trial team prior to randomisation
3. Presence of metastatic disease
4. Unavailable for any trial related follow-up
5. History of malignancy except non-melanomatous skin cancer, CIS cervix, previously unirradiated precancerous changes in the breast (including ductal carcinoma in-situ and lobular carcinoma in-situ), and non-breast malignancy if curative intent and at least 5 years disease free
6. Pregnant and/or currently breast feeding
7. Participating in the PARABLE trial

Date of first enrolment
01/04/2025

Date of final enrolment
30/09/2028

Locations

Countries of recruitment
United Kingdom

England

Northern Ireland

Scotland

Wales

Ireland

Study participating centre
Royal Marsden Hospital
Royal Marsden Hospital
Downs Road

Sutton
United Kingdom
SM2 5PT

Study participating centre
The Royal Marsden Hospital (london)
Fulham Road
London
United Kingdom
SW3 6JJ

Study participating centre
The Christie
550 Wilmslow Road
Withington
Manchester
United Kingdom
M20 4BX

Study participating centre
Addenbrookes
Addenbrookes Hospital
Hills Road
Cambridge
United Kingdom
CB2 0QQ

Study participating centre
Beatson West of Scotland Cancer Centre
1053 Great Western Road
Glasgow
United Kingdom
G12 0YN

Study participating centre
Belfast City Hospital
51 Lisburn Rd
Belfast
United Kingdom
BT9 7AB

Study participating centre
Bristol Haematology & Oncology Centre
Horfield Road
Bristol
United Kingdom
BS2 8ED

Study participating centre
Charing Cross Hospital
Fulham Palace Road
London
United Kingdom
W6 8RF

Study participating centre
Cheltenham General Hospital
Sandford Road
Cheltenham
United Kingdom
GL53 7AN

Study participating centre
Churchill Hospital
Churchill Hospital
Old Road
Headington
Oxford
United Kingdom
OX3 7LE

Study participating centre
Clatterbridge Cancer Centre
Clatterbridge Hospital
Clatterbridge Road
Wirral
United Kingdom
CH63 4JY

Study participating centre

Derriford Hospital
Derriford Road
Crownhill
Plymouth
United Kingdom
PL6 8DH

Study participating centre
Colchester General Hospital
Turner Road
Colchester
United Kingdom
CO4 5JL

Study participating centre
Guys Hospital
Guys Hospital
Great Maze Pond
London
United Kingdom
SE1 9RT

Study participating centre
Ipswich Hospital
Heath Road
Ipswich
United Kingdom
IP4 5PD

Study participating centre
South Tees Hospitals NHS Foundation Trust
James Cook University Hospital
Marton Road
Middlesbrough
United Kingdom
TS4 3BW

Study participating centre
Leicester Royal Infirmary
Infirmary Square
Leicester

United Kingdom
LE1 5WW

Study participating centre
United Lincolnshire Teaching Hospitals NHS Trust
Lincoln County Hospital
Greetwell Road
Lincoln
United Kingdom
LN2 5QY

Study participating centre
Mount Vernon Hospital
Rickmansworth Road
Northwood
United Kingdom
HA6 2RN

Study participating centre
Musgrove Park Hospital (taunton)
Musgrove Park Hospital
Taunton
United Kingdom
TA1 5DA

Study participating centre
Norfolk and Norwich University Hospitals NHS Foundation Trust
Colney Lane
Colney
Norwich
United Kingdom
NR4 7UY

Study participating centre
Glan Clwyd General Hospital NHS Trust
Ysbyty Glan Clwyd
Sarn Lane
Bodelwyddan
United Kingdom
LL18 5UJ

Study participating centre
Nottingham University Hospitals NHS Trust - City Campus
Nottingham City Hospital
Hucknall Road
Nottingham
United Kingdom
NG5 1PB

Study participating centre
Peterborough City Hospital
Edith Cavell Campus
Bretton Gate
Bretton
Peterborough
United Kingdom
PE3 9GZ

Study participating centre
Queen Alexandras Hospital
Southwick Hill Road
Cosham
Portsmouth
United Kingdom
PO6 3LY

Study participating centre
Barking, Havering and Redbridge University Hospitals NHS Trust
Queens Hospital
Rom Valley Way
Romford
United Kingdom
RM7 0AG

Study participating centre
Royal Berkshire Hospital
London Road
Reading
United Kingdom
RG1 5AN

Study participating centre
Royal Devon University Healthcare NHS Foundation Trust
Royal Devon University NHS Ft
Barrack Road
Exeter
United Kingdom
EX2 5DW

Study participating centre
Royal Free London NHS Foundation Trust
Royal Free Hospital
Pond Street
London
United Kingdom
NW3 2QG

Study participating centre
Lancashire Teaching Hospitals NHS Foundation Trust
Royal Preston Hospital
Sharoe Green Lane
Preston
United Kingdom
PR2 9HT

Study participating centre
Royal Surrey County Hospital Guildford
Egerton Road
Guildford
United Kingdom
GU2 7XX

Study participating centre
Brighton and Sussex University Hospitals NHS Trust
Royal Sussex County Hospital
Eastern Road
Brighton
United Kingdom
BN2 5BE

Study participating centre

Singleton Hospital
Sketty Lane
Sketty
Swansea
United Kingdom
SA2 8QA

Study participating centre
University Hospital Southampton NHS Foundation Trust
Southampton General Hospital
Tremona Road
Southampton
United Kingdom
SO16 6YD

Study participating centre
Barts and the London NHS Trust
The Royal London Hospital
Whitechapel
London
United Kingdom
E1 1BB

Study participating centre
Leeds Teaching Hospitals NHS Trust
St. James's University Hospital
Beckett Street
Leeds
United Kingdom
LS9 7TF

Study participating centre
University Hospitals Coventry and Warwickshire NHS Trust
Walsgrave General Hospital
Clifford Bridge Road
Coventry
United Kingdom
CV2 2DX

Study participating centre

Velindre Cancer Centre
Velindre Road
Cardiff
United Kingdom
CF14 2TL

Study participating centre
Western General Hospital
Crewe Road South
Edinburgh
Lothian
United Kingdom
EH4 2XU

Study participating centre
Weston Park Hospital NHS Trust
Whitham Road
Sheffield
United Kingdom
S10 2SJ

Study participating centre
Worcestershire Acute Hospitals NHS Trust
Worcestershire Royal Hospital
Charles Hastings Way
Worcester
United Kingdom
WR5 1DD

Sponsor information

Organisation
Institute of Cancer Research

ROR
https://ror.org/00dpztj76

Funder(s)

Funder type

Government

Funder Name
National Institute for Health and Care Research

Alternative Name(s)
National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

Funding Body Type
Government organisation

Funding Body Subtype
National government

Location
United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan
Not provided at time of registration

IPD sharing plan summary
Data sharing statement to be made available at a later date