

A single-centre, cross-over trial evaluating the metabolic effects of a ketone ester food supplement in intensive care patients

Submission date 24/10/2023	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 27/10/2023	Overall study status Ongoing	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 16/12/2025	Condition category Other	<input type="checkbox"/> Individual participant data <input checked="" type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

The majority of long-term critically ill patients develop muscle weakness and muscle wasting during their stay in intensive care, which hampers recovery and leads to increased complications and a greater risk of death. In animal experiments administering a nutritional supplement, namely ketone esters, can improve muscle weakness during disease. The next step in this research is to see if this supplement can be used to prevent or improve muscle weakness in critically ill patients. The first step is to determine a well-tolerated and effective dose of the dietary supplement (ketone esters) in critically ill patients.

Who can participate?

Adults over the age of 18 years who are admitted to the intensive care unit in UZ Leuven

What does the study involve?

This pilot study will test three doses (25 g, 50 g and 75 g) of ketone esters in adult critically ill patients. Participants are randomly allocated to receive the product or a placebo through a nasogastric feeding tube on 2 consecutive days. Frequent blood samples are scheduled to determine at which dose there is an effective increase in ketones in the blood without any adverse side effects.

What are the possible benefits and risks of participating?

There are no known interactions of ketone esters with drugs. Still, it is possible that nausea or abdominal pain, for example, may occur if the dietary supplement is not well tolerated.

Where is the study run from?

University Hospitals Leuven (Belgium)

When is the study starting and how long is it expected to run for?

January 2023 to December 2027

Who is funding the study?

1. European Research Council (ERC)
2. Flemish Government (Belgium)

Who is the main contact?

Prof. Dr. Greet Van den Berghe, greet.vandenbergh@kuleuven.be

Contact information

Type(s)

Principal investigator

Contact name

Prof Greet Van den Berghe

ORCID ID

<https://orcid.org/0000-0002-5320-1362>

Contact details

Herestraat 49
Leuven
Belgium
3000
+32 (0)16 344021
greet.vandenbergh@kuleuven.be

Type(s)

Scientific

Contact name

Prof Lies Langouche

ORCID ID

<https://orcid.org/0000-0002-8564-6809>

Contact details

Herestraat 49
Leuven
Belgium
3000
+32 (0)16 33 05 24
lies.langouche@kuleuven.be

Type(s)

Public

Contact name

Mrs Liese Mebis

ORCID ID

<https://orcid.org/0000-0002-6941-3044>

Contact details

Herestraat
Leuven
Belgium
3000
+32 (0)16 343125
liese.mebis@uzleuven.be

Type(s)

Scientific

Contact name

Prof Jan Gunst

ORCID ID

<https://orcid.org/0000-0003-2470-6393>

Contact details

Herestraat 49
Leuven
Belgium
3000
+32 (0)16 344021
jan.gunst@uzleuven.be

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

Protocol serial number

S67928

Study information

Scientific Title

A single-centre, placebo-controlled, cross-over randomized controlled trial evaluating the metabolic effects of a ketone ester food supplement in intensive care patients: the KETOCARE RCT

Acronym

KETOCARE RCT

Study objectives

To study the feasibility, safety and tolerability of enteral ketone ester supplementation to increase circulating ketone levels in adult critically ill patients.

Ethics approval required

Ethics approval required

Ethics approval(s)

approved 12/09/2023, Ethics Committee Research UZ/KU Leuven (Herestraat 49, Leuven, 3000, Belgium; +32 (0)16 34 86 00; ec@uzleuven.be), ref: S67928

Study design

Single-center placebo-controlled cross-over randomized controlled trial

Primary study design

Interventional

Study type(s)

Efficacy, Safety

Health condition(s) or problem(s) studied

Adult critically ill patients

Interventions

A blocked randomization scheme with blocks of 4 patients will be used in this study. The code for randomization blinding is stored in the randomization tool, developed in FilemakerPro with restricted access to the database administrators. Randomization will be carried out by dedicated research staff members using a role-based user access.

With a cross-over design, patients will receive in random order on 2 consecutive days ketone ester (3HHB) and a placebo (5% glucose). The ketone esters will be supplied by TDeltaS Ltd (UK).

Patients will receive the investigational product between ICU days 4 and 10. The investigational product (3HHB) or placebo will be delivered through the enteral feeding tube via repeated small boluses. The researchers will first perform this trial with a 25 g dose of 3HHB divided over 6 boluses of 3,9 ml. Next, they will assess the primary and secondary endpoints in this cohort of patients. If the dose of 25 g is found safe but ineffective for increasing plasma 3-hydroxybutyrate at least with 0.22 mmol/l, a second cohort of patients will be tested with the same cross-over study design but with a higher 3HHB dose (50 g of 3HHB, divided over 6 boluses of 7,8 ml). If this dose is again safe but ineffective, a third cohort of patients will be tested with the same cross-over study design with an increased dose of 75 g of 3HHB, divided over 6 doses of 11.7ml. For each cohort of patients tested (cohort 1 to test 25 g, cohort 2 to test 50 g and cohort 3 to test 75 g), 16 patients will be included.

Intervention Type

Supplement

Primary outcome(s)

Increase in plasma 3-hydroxybutyrate by at least 0.22 mmol/l following administration of enteral 3HHB, measured by LC-MS/MS

Key secondary outcome(s)

1. Plasma levels of 3HHB and metabolites 3-hydroxybutyrate and 1,3-butanediol measured over time during (6-hour period) and after the intervention (12-hour period), measured by LC-MS/MS
2. Urine levels of 3HHB and metabolites 3-hydroxybutyrate and 1,3-butanediol over time during (6-hour period) and after the intervention (12-hour period), measured by LC-MS/MS

3. Plasma levels of cholesterol (HDL, LDL, total), triglycerides and free fatty acids over time during (6-hour period) and after the intervention (12-hour period), measured by commercial kits
4. Blood glucose concentrations during the 48 h study period, measured by blood-gas-analyzer
5. Incidence of severe (<40 mg/dl) hypoglycemia during the intervention window and until 12 h after stopping study infusion (3HHB or placebo), measured with a blood gas analyser in the ICU
6. Incidence of ketoacidosis during the intervention window and until 12 h after stopping the study infusion, measured with a ketone stick test on arterial blood in the ICU
7. Intolerance for the supplement (including abdominal discomfort or vomiting) during the intervention up until 48 hours after the last intervention, as reported by the patient or observed by the attending (study) nurse

Completion date

31/12/2027

Eligibility

Key inclusion criteria

1. Voluntary written informed consent of the participant or their legally authorized representative has been obtained
2. Age \geq 18 years
3. Patient expected to stay at the ICU for at least 5 days
4. The presence of a nasogastric feeding tube

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Mixed

Lower age limit

18 years

Upper age limit

99 years

Sex

All

Total final enrolment

36

Key exclusion criteria

1. Therapy restriction code
2. Patients refusing blood transfusion upon ICU admission will be considered as having a therapy restriction upon admission and will not be included
3. Expected to die within 48 hours after screening (= moribund patients)
4. No arterial and central venous line, or expected to have one of these lines removed before the

end of the study period (= not critically ill enough to be representative for the future target population).

5. Contraindication for enteral feeding

6. Readmission to the ICU after previous inclusion in the RCT

7. Inborn metabolic disease

8. Receiving ketogenic diet in ICU

9. Underweight (BMI<20) or admitted with complications due to anorexia nervosa

10. Known to be pregnant or lactating

11. ICU admission with diabetic ketoacidosis or hyperosmolar hyperglycemic state

12. Acute or chronic liver failure

13. High glucose need to prevent spontaneous hypoglycemia

14. Metabolic acidosis (pH <7.30 and bicarbonate <18 mmol/l)

Date of first enrolment

08/01/2024

Date of final enrolment

13/06/2024

Locations

Countries of recruitment

Belgium

Study participating centre

UZ Leuven

Herestraat 49

Leuven

Belgium

3000

Sponsor information

Organisation

Universitair Ziekenhuis Leuven

ROR

<https://ror.org/0424bsv16>

Funder(s)

Funder type

Research council

Funder Name

European Research Council

Alternative Name(s)

The European Research Council, ERC

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location**Funder Name**

Vlaamse regering (Methusalem Program)

Alternative Name(s)

Flanders, Flemish Government, Flandre, Flandern, Vlaanderen

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

Belgium

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study will be stored in a publicly available repository (name: RDR; <https://rdr.kuleuven.be/>).

IPD sharing plan summary

Stored in publicly available repository

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Protocol file	version 1.3	17/04/2024	24/04/2024	No	No