

# Testing a novel, self-contained chest drainage system (THOPAZ+) for post-operative cardiac drainage

<b>Submission date</b> 29/05/2019	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 07/06/2019	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
<b>Last Edited</b> 27/08/2019	<b>Condition category</b> Surgery	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

Most cardiac procedures still require the use of cardiopulmonary bypass (CPB) and a sternotomy. One of the complications after such procedures is bleeding, since patients have to be fully anti-coagulated during CPB and therefore require post-operative chest drainage.

**Aim:** A new, self-contained, electronic, continuous pump-driven chest drainage system is compared for the first time in an RCT to a traditional wall-suction system in cardiac surgery.

### Who can participate?

Adult elective cardiac patients undergoing CABG and/or valve surgery.

### What does the study involve?

Patients undergoing cardiac surgery require post-operative chest-drainage in order to evacuate blood and clot formation. This is achieved with one or several chest-tubes which are hooked to a conventional canister system drained by vacuum. The aim of our study was to assess a new, self-contained, battery-operated, digital drainage system (THOPAZ+) and to compare it to a conventional system. Our prospective, randomized clinical trial included 120 patients undergoing elective routine cardiac surgery. They were randomized between the group with conventional and the group with THOPAZ+ drainage.

### What are the possible benefits and risks of participating?

The benefits of the new system are a continuous drainage from the operating room until the chest-tube removal, the possibility to mobilize the patient and the presence of alarms on the THOPAZ+ system and a shorter drainage duration which may have a positive impact on the outcome of the patient.

### Where is the study run from?

Department of Cardiac Surgery, University Hospital, Verona, Italy.

### When is the study starting and how long is it expected to run for?

September 2015 to June 2016

Who is funding the study?  
Medela AG, Switzerland

Who is the main contact?  
Dr Luca Barozzi,  
luca.barozzi@ospedaleuniverona.it

## Contact information

**Type(s)**  
Scientific

**Contact name**  
Dr Luca Barozzi

**ORCID ID**  
<https://orcid.org/0000-0001-5297-0058>

**Contact details**  
Cardiac Surgery  
University Hospital of Verona  
Piazzale le Stefani 1  
Verona  
Italy  
37126  
0039 347 2739 436  
luca.barozzi@ospedaleuniverona.it

## Additional identifiers

**Clinical Trials Information System (CTIS)**  
Nil known

**Protocol serial number**  
584CESC

## Study information

**Scientific Title**  
Randomized controlled trial of a novel, self-contained chest drainage system in cardiac surgery

**Acronym**  
ThopCard2

**Study objectives**  
Improvement of chest drainage: we have investigated a new chest drainage system THOPAZ+ (Medela AG, Switzerland) to improve and simplify the management of chest drainage after cardiac surgical procedures and which gives the patient a small, mobile, user-friendly, continuous suction from the operating table to the ICU/ward.

## **Ethics approval required**

Old ethics approval format

## **Ethics approval(s)**

Approved 28/07/2015, Ethics Committee of the University of Verona (Borgo Trento, P. le A. Stefani, 1-37126 Verona, Italy; 045 8123236; comitatoetico.veronarovigo@ospedaleuniverona.it), ref: 584CESC.

## **Study design**

Interventional single centre study

## **Primary study design**

Interventional

## **Study type(s)**

Treatment

## **Health condition(s) or problem(s) studied**

Cardiac surgery

## **Interventions**

This study compared the conventional chest drainage system (Argyle Aqua-Seal, Covidien, USA) with the new THOPAZ+ digital self-contained system from the end of cardiac surgery until the chest tube removal.

Randomization: Patients were allocated to either the usual suction system or the Thopaz+ pump using block randomization, with blocks of 4 or 6 (block size is also random). Group allocation was put in sealed envelopes, which were opened after surgery, just before the OR nurse opened the tubing system. Thus, the surgical team was blinded to the suction system during the surgery.

All the data were collected prospectively during the study by two surgical residents and supervised by Dr. Luca Barozzi.

## **Intervention Type**

Device

## **Phase**

Not Applicable

## **Primary outcome(s)**

Adequacy of chest drainage with THOPAZ+ measured on the digital read-out of the THOPAZ+ (in ml of drainage fluid) at the following time points: end of OR, before and after transport to ICU, on ICU (hourly), POD1, and at chest-tube removal.

## **Key secondary outcome(s)**

Patient safety and Satisfaction Assessment of the hospital personnel.

1. The patient safety was part of the routine ICU care of the patient and special attention was given to the functioning of the THOPAZ+ chest drainage system, e.g. fluid collection, air leak.
2. The satisfaction assessment of the hospital personnel was performed with a short, internet-based form was filled by the users according to their specialty, e.g. cardiac surgeons,

anesthetists, nurses to measure the following characteristics of the device on a visual analog scale (from 0 to 10).

- Ease of use
- No/reduced milking
- Mobility of patients
- Noise reduction
- Gain of time
- Security for patients
- How useful were the alarms?

**Completion date**

20/06/2016

## **Eligibility**

**Key inclusion criteria**

1. Adult aged 18 - 80 years
2. Scheduled to undergo elective, first-time: CABG, valve surgeries and combinations with cardio-pulmonary bypass (CPB)

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Adult

**Lower age limit**

18 years

**Sex**

All

**Key exclusion criteria**

1. Unstable angina
2. Emergency procedures
3. Off-pump surgery
4. Re-operation
5. Anti-coagulation or anti-platelet therapy (except aspirin cardio) until surgery

**Date of first enrolment**

17/09/2015

**Date of final enrolment**

14/06/2016

## **Locations**

## Countries of recruitment

Italy

## Study participating centre

### Department of Cardiac Surgery

University Hospital

Piazzale le Stefani 1

Verona

Italy

37126

## Sponsor information

### Organisation

Medela AG

### ROR

<https://ror.org/02cedh792>

## Funder(s)

### Funder type

Industry

### Funder Name

Medela AG

## Results and Publications

### Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are available from the corresponding author on special request from Dr. Luca Barozzi, [luca.barozzi@ospedaleuniverona](mailto:luca.barozzi@ospedaleuniverona). it with the following access criteria: detailed description for which purpose the anonymized data, or part of the data, will be used and only after having received permission from the Ethical Commission of the University of Verona.

### IPD sharing plan summary

Data sharing statement to be made available at a later date