

Increased brain activation after painful stimulation of the forearm muscles in patients with fibromyalgia syndrome

Submission date 02/09/2015	Recruitment status No longer recruiting	☐ Prospectively registered
		☐ Protocol
Registration date 24/09/2015	Overall study status Completed	☐ Statistical analysis plan
		☒ Results
Last Edited 22/10/2015	Condition category Musculoskeletal Diseases	☐ Individual participant data

Plain English summary of protocol

Background and study aims

Fibromyalgia syndrome (FMS) is a common long-term condition, which causes widespread muscle and joint pain all over the body. The exact cause of FMS is unknown, but it is thought that a variety of physical, mental and emotional factors are responsible. One theory is that FMS is to do with an abnormal increase of chemicals in the brain (neurotransmitters), which causes people to feel pain more intensely and are more sensitive to pain. Many studies link fibromyalgia with depression, which is also related to abnormal levels of neurotransmitters in the brain, however little is known about how these two conditions might influence each other. The aim of this study is to find out whether the cortex of the brain (which is responsible for consciousness) is more active in people with FMS than in people with depression with no pain and healthy controls.

Who can participate?

Adults with fibromyalgia syndrome, depression with no pain, and healthy age matched controls.

What does the study involve?

Participants receive two different forms of stimulation. In the first stimulation, painful pressure is applied to the forearm of the patient, and in the second, a word-based memory test (verbal fluency test) is performed. Whilst these stimulations are happening, participants undergo a special type of brain imaging which shows which areas of the brain are active, by looking at the amount of blood flow (functional near-infrared spectroscopy). All participants are also asked to complete questionnaires to measure their levels of pain and emotion.

What are the possible benefits and risks of participating?

Participants receive no direct benefits from the study as it is an observational study. There are no risks of participating in the study.

Where is the study run from?

University of Würzburg and University of Tübingen (Germany)

When is the study starting and how long is it expected to run for?
January 2007 to December 2012

Who is funding the study?
University of Würzburg (Germany)

Who is the main contact?
Dr Nurcan Üçeyler
ueceyler_n@ukw.de

Contact information

Type(s)
Scientific

Contact name
Dr Nurcan Üçeyler

ORCID ID
<https://orcid.org/0000-0001-6973-6428>

Contact details
Department of Neurology
University of Würzburg
Josef-Schneider-Str. 11
Würzburg
Germany
97080
+49 931 201 24617
ueceyler_n@ukw.de

Additional identifiers

Study information

Scientific Title
Increased cortical activation upon painful stimulation in fibromyalgia syndrome

Study objectives
Cortical activation upon painful stimulation is increased in patients with fibromyalgia syndrome compared to patients with depression and no pain and to healthy controls.

Ethics approval required
Old ethics approval format

Ethics approval(s)
Würzburg Medical School Ethics Committee, 27/01/2009, ref: 12/09

Primary study design

Observational

Study design

Non-interventional single-center study observational study.

Study type(s)

Other

Health condition(s) or problem(s) studied

Fibromyalgia

Interventions

Twenty five patients with fibromyalgia syndrome, ten patients with unipolar major depression (MD) without pain, and thirty five healthy controls are recruited for the study. Patients undergo functional near-infrared spectroscopy (fNIRS) whilst being subjected to two stimulations:

1. Painful pressure stimulation at the dorsal forearm
2. Verbal fluency test (VFT) to assess cognitive function using memory recall

All patients underwent neurological examination and all subjects were investigated with questionnaires (pain, depression, FMS, empathy).

Intervention Type

Primary outcome(s)

Muscular pressure pain: fNIRS measurements were performed during the application of painful pressure on the muscle bulk of the finger extensors of the right side using a calibrated algometer. The stimulation conditions were as follows: pressure application for two seconds; pause for ten seconds between two stimuli; total of 40 stimuli, i.e. measurement at baseline and up to 8 minutes after first stimulation.

Key secondary outcome(s)

Verbal fluency test (VFT): The VFT paradigm consisted of three conditions. Subjects were asked to produce as many different nouns as possible a) starting with a certain letter (A, F, and S), or b) belonging to the same category (animals, fruits, and flowers) or c) to name the days of the week as a control condition. Each condition lasted for 30 sec followed by 30 sec rest. Subjects worked on nine blocks in total (3 x letters, 3 x categories, 3 x week days), i.e. measurement at baseline and up to 9 minutes after start.

Completion date

31/12/2012

Eligibility

Key inclusion criteria

1. Aged 18 years or over
2. Patients with:
 - 2.1. Fibromyalgia syndrome
 - 2.2. Unipolar major depression (MD) without pain
 - 2.3. Healthy controls

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 Years

Sex

All

Key exclusion criteria

1. Other possible differential diagnoses (e.g. rheumatoid arthritis; post-surgery pain)
2. Current or prior cerebral disease (e.g. stroke, cerebral hemorrhage, head trauma)
3. Any clinically relevant psychiatric disorder (examined by systematic psychiatric interview)

Date of first enrolment

01/01/2007

Date of final enrolment

31/12/2011

Locations**Countries of recruitment**

Germany

Study participating centre**University of Würzburg**

Department of Neurology

Sanderring 2

Würzburg

Germany

97080

Study participating centre**University of Würzburg**

Department of Psychiatry

Sanderring 2

Würzburg

Germany

97080

Study participating centre
University of Tübingen
Department of Psychiatry
Geschwister-Scholl-Platz
Tübingen
Germany
72070

Sponsor information

Organisation
University of Würzburg

ROR
<https://ror.org/03pvr2g57>

Funder(s)

Funder type
University/education

Funder Name
University of Würzburg (EFIC-Grünenthal Grant and intramural funds)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary
Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	20/10/2015		Yes	No