

A study to compare the efficacy of intra-articular injection of methylprednisolone versus triamcinolone hexacetonide in patients with knee osteoarthritis

Submission date 16/09/2013	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 25/09/2013	Overall study status Completed	<input type="checkbox"/> Protocol
Last Edited 10/06/2016	Condition category Musculoskeletal Diseases	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Steroids can be used in the treatment of patients with knee osteoarthritis (arthritis of the knee), but the best agent is still unknown. This study aimed to compare the effect of two drugs (triamcinolone hexacetonide and methylprednisolone acetate) on the condition.

Who can participate?

Men and women who are at least 40 years old, with painful knee osteoarthritis took part in this study.

What does the study involve?

Participants were randomly allocated to receive a single knee injection of either triamcinolone hexacetonide or methylprednisolone acetate. After 24 weeks participants attended a follow-up visit to assess the effectiveness of the treatment with questionnaires.

What are the possible benefits and risks of participating?

Possible benefits were pain relief and physical function improvement. Risks were related to the injection and included pain, swelling, bleeding and infection in the knee.

Where is the study run from?

The study was run from Hospital Heliópolis, São Paulo, Brazil

When is the study starting and how long is it expected to run for?

The study started in December 2010 and ended in May 2013

Who is funding the study?

The investigators funded the study

Who is the main contact?

Dr Andrea Lomonte
avannucci@bol.com.br

Contact information

Type(s)

Scientific

Contact name

Dr Andrea Lomonte

Contact details

Rua Maria Bucalem Haddad
123, ap 52, bloco B - Vila Firmiano Pinto
Sao Paulo
Brazil
04125-010

Additional identifiers

Protocol serial number

N/A

Study information

Scientific Title

A randomized, double-blind study to evaluate the efficacy of intra-articular injection of methylprednisolone versus triamcinolone hexacetonide in patients with knee osteoarthritis

Study objectives

Intra-articular injections of methylprednisolone acetate and triamcinolone hexacetonide have similar efficacy at reducing knee pain and improving physical function both in the short as in the long-term in patients with symptomatic knee osteoarthritis.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Research Ethics Committee of the Hospital Heliopolis [Comitê de Ética em Pesquisa do Hospital Heliópolis], 14/10/2008

Study design

24 weeks prospective randomized two arms double-blinded study

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Knee osteoarthritis

Interventions

1. Intra-articular triamcinolone hexacetonide 40mg single dose
2. Intra-articular methylprednisolone acetate 40mg single dose

Total duration of the study: 24 weeks

Intervention was one day (single injection)

Follow-up: 24 weeks

Intervention Type

Drug

Phase

Not Applicable

Drug/device/biological/vaccine name(s)

Methylprednisolone, triamcinolone hexacetonide

Primary outcome(s)

Mean change from baseline in patients assessment of pain in the targeted knee by VAS at week 4

Key secondary outcome(s)

1. Mean changes from baseline in patients assessment of pain by VAS at 12 and 24 weeks
2. Mean changes from baseline in patients and physicians global assessment of disease activity by VAS
3. Improvement in patients global assessment of disease by Likert scale
4. Mean changes from baseline in total Western Ontario and McMaster Universities (WOMAC) questionnaire and its subscores of pain, stiffness and physical function
5. Mean changes from baseline in total Lequesne algofunctional index
6. Outcome Measures in Rheumatology Clinical Trials and Osteoarthritis Research Society International (OMERACT-OARSI) response criteria
7. Analysis of predictors of response

All secondary outcomes were measured at 4, 12 and 24 weeks.

Completion date

17/05/2013

Eligibility**Key inclusion criteria**

1. Diagnosis of knee osteoarthritis (OA) according to the American College of Rheumatology criteria
2. Kellgren-Lawrence radiographic grade II or III of the knee
3. Visual analogue scale (VAS) for knee pain of at least 40mm (maximum 100mm)

4. Both genders, aged 40 years or older
5. Failure to control OA symptoms with previous or current analgesics and/ or non steroidal antiinflammatory drugs

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Key exclusion criteria

1. Other rheumatic or inflammatory condition
2. Symptomatic disease of the lower limbs (other than knee OA)
3. Serious and/or uncontrolled concomitant medical illness
4. Body mass index (BMI) greater or equal to 35kg/m²
5. IA injection of corticosteroid or hyaluronic acid in the previous 6 months
6. knee replacement in the targeted joint
7. Local or systemic infection
8. Pregnancy
9. Skin lesions in the IA injection site
10. Physical therapy for the knee
11. Known hypersensitivity to corticosteroids or lidocaine; and the use of anticoagulants

Date of first enrolment

18/12/2010

Date of final enrolment

17/05/2013

Locations**Countries of recruitment**

Brazil

Study participating centre

Rua Maria Bucalem Haddad

Sao Paulo

Brazil

04125-010

Sponsor information

Organisation

Hospital Heliópolis (Brazil)

ROR

<https://ror.org/01bkdwe73>

Funder(s)

Funder type

Other

Funder Name

Investigator initiated and funded (Brazil)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/09/2015		Yes	No