

# Examining antibiotics for ulcerated skin cancer surgical excision

<b>Submission date</b> 20/06/2025	<b>Recruitment status</b> Recruiting	<input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
<b>Registration date</b> 08/08/2025	<b>Overall study status</b> Ongoing	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
<b>Last Edited</b> 11/03/2026	<b>Condition category</b> Infections and Infestations	<input type="checkbox"/> Individual participant data <input checked="" type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

Skin cancer is the most common type of cancer. Every year in the UK, around 200,000 people have their skin cancer removed surgically. Some people develop skin cancers that break through the skin surface, causing a wound on the skin (called an ulcerated skin cancer), which are six times more likely to develop a wound infection after surgery. Doctors often prescribe antibiotics at the time of surgery to prevent wound infections, but it is unknown whether antibiotics reduce the risk of getting an infection. Using more antibiotics than are needed may lead to patients having unnecessary side effects and lead to the bacteria causing the infection becoming resistant to antibiotics, which then work less well in the future. The study looks at whether antibiotics should be prescribed to patients at the time of surgically removing their ulcerated skin cancer to reduce their risk of wound infection. Participants will be given a one-off dose of antibiotic or no antibiotic (dummy pill called placebo). The study will see how many in each group develop wound infections. This will help to decide whether antibiotics should be given to patients before skin surgery.

### Who can participate?

Patients with ulcerated skin cancers from participating UK NHS hospitals.

### What does the study involve?

Before surgery, participants will be randomly allocated by a computer to have a one-off dose of a common antibiotic or no antibiotics (placebo). Everyone will receive the same wound care advice that they normally get as part of their NHS treatment. Participants will be asked to contact us if they are worried about their wound, and they will receive a photo booklet to help identify potential infections. They will be contacted between days 5-10, 15-20 and 30 days after surgery to ask about their wound, signs of infection and side effects. Participants will be seen in the hospital if a wound infection is suspected, and all will receive additional treatment if required. They will also be asked how long it took to return to normal activity, and the impact on daily activities.

### What are the possible benefits and risks of participating?

By taking part in this trial, regardless of which treatment is received by participants, they will be helping to determine if antibiotics given just before surgery can reduce the number of wound

infections seen in patients who have this type of surgery. Participants will be followed up more closely than they would be if they were not taking part in the trial.

The antibiotic, flucloxacillin, has been used for a long time for treating skin and wound infections and has a low risk of side effects that will cause no harm to pregnant/ breastfeeding participants or male participants' partners. However, some common side effects include feeling sick, bloating, indigestion and diarrhoea. Participants will be closely monitored as part of their usual care. If participants do experience side effects, they will be encouraged to let their local recruitment site know, or if urgent, contact their GP or emergency department. Oral flucloxacillin is currently in use in the UK NHS to reduce SSI risk following skin cancer surgery. A multi-speciality clinician survey (n=129) conducted in 2021 reported that peri-operative antibiotics were 'always'/ 'often'/ 'sometimes' prescribed for excision of ulcerated skin cancers by 49% of respondents and 'rarely/ never' by 51%, demonstrating clinical equipoise. This demonstrates that both the use and non-use of the proposed intervention in the EXCISE trial are considered the current standard of practice for skin cancer surgery in the NHS. To minimise risk, potential participants with documented poor renal function (creatinine clearance < 10ml/min) or with a previous allergic reaction to penicillin will be excluded from participation. In addition to the eligibility criteria, other risk mitigation strategies include trial oversight from the Trial Management Group (TMG), Independent Data Monitoring Committee (IDMC) and Trial Steering Committee (TSC).

Participants may also be randomised to receive a placebo. This is unlikely to cause any harm as it does not contain any of the active ingredients, although it is possible that a participant could react with the excipient.

If they are diagnosed with a wound infection, they will be asked to attend their local recruiting site for review. If they are diagnosed with a surgical site infection, they will have swabs taken of the wound for antimicrobial resistance analysis in addition to those taken for standard care. These wound swabs may cause some slight discomfort. Participants will be offered compensation for travel to their local recruiting site if required to ensure that the participant is not disadvantaged.

Completion of patient diaries, taking wound photos and answering questionnaires about their health is not anticipated to cause any pain, discomfort, distress or intrusion, although it will take up participants' time.

Additionally, a selection of participants, those who declined to participate and health care professionals will be asked to take part in a semi-structured interview about their experience of taking part in the trial. It is not likely to involve any particular risks, although participating in such a discussion may bring back memories of a difficult and distressing time.

Where is the study run from?  
University Hospital of Wales, UK

When is the study starting and how long is it expected to run for?  
March 2026 to December 2027

Who is funding the study?  
The National Institute for Health and Care Research (NIHR), UK

Who is the main contact?  
Mr Burhan Ben Karatas excise-ctr@cardiff.ac.uk

## Contact information

### Type(s)

Public, Scientific

### Contact name

Mr Burhan Ben Karatas

### Contact details

Centre for Trials Research, Cardiff University, 7th Floor, Neuadd Meirionnydd, Heath Park  
Cardiff

United Kingdom

CF14 4YS

-

excise-ctr@cardiff.ac.uk

### Type(s)

Principal investigator

### Contact name

Dr Rachel Abbott

### Contact details

Welsh Institute of Dermatology, Glamorgan House, University Hospital of Wales, Heath Park  
Cardiff

United Kingdom

CF14 4XW

-

rachel.abbott@wales.nhs.uk

## Additional identifiers

### Integrated Research Application System (IRAS)

1009425

### Central Portfolio Management System (CPMS)

60084

### Protocol serial number

24/APR/8825E

## Study information

### Scientific Title

EXamining antibiotics for ulCerated skin cancer Surgical Excision (EXCISE): a pragmatic, double-blinded clinical and cost effectiveness randomised controlled trial.

### Acronym

EXCISE

## Study objectives

This study wants to know whether antibiotics should be prescribed to patients at the time of surgically removing their ulcerated skin cancer to reduce their risk of wound infection. Participants will be given a one-off dose of antibiotic or no antibiotic (dummy pill called placebo). The study will see how many in each group develop wound infections. This will help us to decide whether antibiotics should be given to patients before skin surgery.

- The study will see what the side effects are for taking a single dose of oral flucloxacillin.
- The study will measure the antibiotic resistance that occurs in infected wounds by taking a swab of wounds from patients who have an infection.
- To look at the costs and benefits of the antibiotic treatment
- To look at participants' and clinicians' views on using antibiotics to prevent wound infection.
- To look at whether participants are willing and able to send wound photos.
- To validate the Wound healing questionnaire to try and decide if a patient has a wound infection without needing to do this in person.

## Ethics approval required

Old ethics approval format

## Ethics approval(s)

Approved 08/08/2025, Wales REC 3 (Health and Care Research Wales, Castlebridge 4, Cardiff, CF11 9AB, UK; +44 (0)2922 941107, +44 (0)2922 940954, +44 (0)2922 940963; E. Wales. REC3@wales.nhs.uk), ref: 25/WA/0196

## Study design

Pragmatic double-blind randomized controlled parallel-group trial

## Primary study design

Interventional

## Study type(s)

Efficacy, Safety

## Health condition(s) or problem(s) studied

Antibiotic treatment for the prevention of surgical site infection in participants receiving ulcerated skin cancer excision

## Interventions

To investigate the efficacy of a single pre-operative 1g dose of oral flucloxacillin in preventing SSI within 30 days in adults undergoing surgical excision of an ulcerated skin cancer under local anaesthetic with planned wound closure.

Participants will receive either Intervention: pre-operative oral flucloxacillin 1g (500mg x 2) in addition to standard clinical care, or Comparator: no antibiotic (placebo) and standard clinical care.

Outcomes will be assessed at baseline, 5-10 days, 15-20 days, 30 days and at 3 months post randomisation. Data will be collected by the research nurses either in person or via telephone, email or post, or review of patient records, and entered in electronic format on the EXCISE database. Patients will be contacted at follow-up points, and if (up until 30 days) there is concern

of a wound infection, they will be asked to return to their recruiting site for a face-to-face assessment of their wound.

Participants will be randomised between the two groups (190 participants per group) with a 1:1 allocation ratio. Randomisation will occur electronically prior to surgery. Allocations will be minimised (with a random element) by SSI risk associated with reconstruction method (primary closure, skin graft or local skin flap) and anatomical site (below knee or other anatomical site) and stratified by the recruitment site. A computerised web-based remote randomisation system will be used. The Trial Manager will also be notified that a participant has been randomised via an automated email alert mechanism to the EXCISE email inbox. This system will be maintained by the Cardiff Centre for Trials Research (CTR) and accessed by the local investigator or site staff delegated to do so, following consent and completion of baseline assessments. In the event the online randomisation system is unavailable at the site, or the site has problems accessing the online website, participants will not be disadvantaged by having their surgery delayed and will be excluded from the trial.

A qualitative study will be conducted with trial participants (n=40), Health Care professionals (n=20) and those who have declined to take part in the trial (n=8). Assessment of the fidelity of possible influences on treatment outcomes in the design and implementation of surgical randomised controlled trials, factors affecting trial compliance, wound management and reporting SSI, the acceptability of treatment received, including views on antibiotics, barriers and facilitators and the wound burden.

## **Intervention Type**

Drug

## **Phase**

Phase IV

## **Drug/device/biological/vaccine name(s)**

Flucloxacillin [Flucloxacillin]

## **Primary outcome(s)**

The proportion of participants with surgical site infection within 30 days post-randomisation. Participants will be contacted by a research nurse at days 5-10, 15-20 and day 30 (+/- 7 Days) to ask about their wound and signs of infection. If the participant or research nurse has any concerns, they will be asked to come to the participating centre to confirm surgical site infection using CDC criteria.

## **Key secondary outcome(s)**

Secondary outcome measures

1. The number of participants with adverse events within 30 days of surgery, defined using the Medical Dictionary for Regulatory Activities
2. Antibiotic resistance to clinically relevant antibiotics, and measured in isolates from participants' wounds at diagnosis of SSI and after 7 days if no response to treatment. This will be assessed by a central microbiology laboratory (Public Health Wales)
3. Health-related QoL measured using the EQ-5D-5L questionnaire at baseline, 30 days and 3 months
4. Time to return to normal activity/work measured using 'Time to return to normal activity/work' case report form (CRF) at 30 days and 3 months
5. Resource use (related to wound complications, including SSI) and cost of hospital visits/stays,

measured using a 'resource use measure' CRF at 30 days and 3 months

6. Qualitative process evaluation assessing implementation and acceptability of oral antibiotics, including facilitators and barriers, measured using qualitative interviews at 30 days

7. Feasibility of 'Selfi-wound' photos measured using the number of participants who were able to successfully take and transmit an image of their wound via the online system at days 5-10

8. Validation of WHQ remote use for SSI assessment measured using a Study within A Trial (SWAT) at 30 days

Tertiary outcome

Qualitative process evaluation exploring participant wound burden measured using qualitative interviews at 30 days

### **Completion date**

31/12/2027

## **Eligibility**

### **Key inclusion criteria**

1. Adult patients ( $\geq 16$  years) with a clinically ulcerated suspected skin cancer (at any body site) listed for excision under local anaesthetic with planned wound closure by any secondary care speciality.

2. First time in the EXCISE trial.

3. Ability to provide Informed Consent (by participant or through a participant's personal legal representative).

### **Participant type(s)**

Patient

### **Healthy volunteers allowed**

No

### **Age group**

Mixed

### **Lower age limit**

16 years

### **Upper age limit**

100 years

### **Sex**

All

### **Total final enrolment**

0

### **Key exclusion criteria**

1. Clinical evidence of skin cancer infection at baseline using CDC criteria.

2. Skin tumour removal planned with curettage, Mohs micrographic surgery/ margin-controlled excision or shave excision.

3. Wound left for delayed reconstruction or secondary intention healing or closed with dermal substitute.
4. Concurrent oral antibiotic treatment (<24 hours after last dose).
5. Documented poor renal function (creatinine clearance < 10ml/min).
6. Previous allergic reaction to penicillin.

**Date of first enrolment**

10/03/2026

**Date of final enrolment**

01/04/2027

## **Locations**

**Countries of recruitment**

United Kingdom

England

Northern Ireland

Scotland

Wales

**Study participating centre**

**University Hospital of Wales**

Heath Park

Cardiff

Wales

CF14 4XW

**Study participating centre**

**Walsall Manor Hospital**

Moat Road

Walsall

England

WS2 9PS

**Study participating centre**

**Castle Hill Hospital**

Castle Road

Cottingham

England

HU16 5JQ

**Study participating centre**

**Churchill Hospital**

Old Road  
Headington  
Oxford  
England  
OX3 7LE

**Study participating centre**

**James Cook University Hospital**

Marton Road  
Middlesbrough  
England  
TS4 3BW

**Study participating centre**

**Royal Victoria Hospital**

Department of Dermatology, Belfast HSCNI Trust  
Belfast  
Northern Ireland  
BT12 6BA

**Study participating centre**

**Stoke Mandeville Hospital**

Mandeville Road  
Aylesbury  
England  
HP21 8AL

**Study participating centre**

**Morrison Hospital**

Heol Maes Eglwys  
Cwmrhydyceirw  
Swansea  
Wales  
SA6 6NL

**Study participating centre**

**Northampton General Hospital**

Cliftonville  
Northampton  
England  
NN1 5BD

**Study participating centre**

**Brighton General Hospital**

Elm Grove  
Brighton  
England  
BN2 3EW

**Study participating centre**

**Royal United Hospital**

Combe Park  
Bath  
England  
BA1 3NG

**Study participating centre**

**Victoria Hospital, NHS Fife**

Department of Oral and Maxillofacial Surgery, Hayfield Road  
Kirkaldy  
Scotland  
KY2 5AH

**Study participating centre**

**Norfolk and Norwich University Hospitals NHS Foundation Trust**

Colney Lane  
Colney  
Norwich  
England  
NR4 7UY

**Study participating centre**

**West Suffolk NHS Foundation Trust**

West Suffolk Hospital  
Hardwick Lane

Bury St. Edmunds  
England  
IP33 2QZ

## Sponsor information

### Organisation

Cardiff and Vale University Health Board

### ROR

<https://ror.org/0489f6q08>

## Funder(s)

### Funder type

Government

### Funder Name

National Institute for Health and Care Research

### Alternative Name(s)

National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

### Funding Body Type

Government organisation

### Funding Body Subtype

National government

### Location

United Kingdom

## Results and Publications

### Individual participant data (IPD) sharing plan

The data are held by the Centre for Trials Research, Cardiff, UK. Applications to share the deidentified and anonymised trial data with other investigators for use in future research will be considered, subject to review of the aims and scientific methods of the application, and any contractual obligations required by organisations involved in the study. Requests for data

sharing or collaboration should be made by completing an application form available via the CTR website at <https://www.cardiff.ac.uk/centre-for-trials-research/collaborate-with-us/data-requests>.

## IPD sharing plan summary

Available on request

### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Protocol file</a>	version 1.1	20/10/2025	03/12/2025	No	No
<a href="#">Study website</a>		11/11/2025	11/11/2025	No	Yes