

An Open Randomised Trial of Zoladex versus CMF as Adjuvant Therapy in the Management of Node Positive Stage II Breast Cancer in Pre/Peri-Menopausal Women Aged 50 Years or Less

Submission date 19/08/2002	Recruitment status No longer recruiting	☐ Prospectively registered ☐ Protocol
Registration date 19/08/2002	Overall study status Completed	☐ Statistical analysis plan ☒ Results
Last Edited 13/11/2012	Condition category Cancer	☐ Individual participant data

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr - -

Contact details

UKCCCR Register Co-ordinator
MRC Clinical Trials Unit
222 Euston Road
London
United Kingdom
NW1 2DA

Additional identifiers

Protocol serial number

ZEN2802

Study information

Scientific Title

Study objectives

Not provided at time of registration

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Primary study design

Interventional

Study design

Randomised controlled trial

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Breast

Interventions

Patients are randomised to one of two treatment arms:

1. Arm A: Zoladex, 3.6 mg given subcutaneously every 28 days for 2 years or until a treatment endpoint is reached.
2. Arm B: Chemotherapy with cyclophosphamide, methotrexate and 5-fluorouracil (CMF) repeated every 28 days for six cycles.

Intervention Type

Other

Phase

Not Specified

Primary outcome(s)

Not provided at time of registration

Key secondary outcome(s)

Not provided at time of registration

Completion date

31/12/1996

Eligibility**Key inclusion criteria**

1. Pre- or peri-menopausal
2. Node positive stage II breast cancer, at entry to study, consisting of:

- 2.1. Histologically proven operable invasive breast cancer
- 2.2. Pathologically involved lymph nodes
- c. No evidence of metastatic disease
- 3. No previous systemic therapy for breast cancer
- 4. Adequate hepatic, renal and haematological function
- 5. No bilateral oophorectomy or radiotherapy to the ovaries
- 6. No concurrent or previous malignancy, except squamous or basal cell carcinoma of the skin or carcinoma in situ of the cervix, adequately cone biopsied

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

Female

Key exclusion criteria

Not provided at time of registration

Date of first enrolment

01/01/1995

Date of final enrolment

31/12/1996

Locations**Countries of recruitment**

United Kingdom

England

Study participating centre

UKCCCR Register Co-ordinator

London

United Kingdom

NW1 2DA

Sponsor information

Organisation

AstraZeneca Clinical Research Group (UK)

ROR

<https://ror.org/04r9x1a08>

Funder(s)

Funder type

Industry

Funder Name

AstraZeneca Pharmaceuticals

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	15/12/2002		Yes	No