

The effect of modulating gut microflora on clinical outcome in elective surgical patients.

Submission date 30/09/2005	Recruitment status No longer recruiting	☐ Prospectively registered ☐ Protocol
Registration date 30/09/2005	Overall study status Completed	☐ Statistical analysis plan ☒ Results
Last Edited 27/08/2008	Condition category Signs and Symptoms	☐ Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Mr Bala S Reddy

Contact details
Dept of General Surgery
Scarborough General Hospital
Woodlands Drive
Scarborough
United Kingdom
YO12 6QL
+44 (0)1723 368111
drbalasreddy@yahoo.co.uk

Additional identifiers

Protocol serial number
N0285150582

Study information

Scientific Title

Study objectives

The aim of this study is to investigate whether or not modulation of gastrointestinal microflora might impact upon septic morbidity in surgical patients?

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Primary study design

Interventional

Study design

Randomised controlled trial

Study type(s)

Not Specified

Health condition(s) or problem(s) studied

Signs and Symptoms: Sepsis

Interventions

Patients will be recruited from the elective colorectal admissions at Scarborough Hospital and will be randomised by a series of sealed envelopes into one of the four groups when they are seen in the pre-assessment clinic.

Group 1 - Control group.

Group 2 - Will receive Neomycin prior to surgery plus bowel preparation.

Group 3 - Neomycin, bowel prep, plus synbiotics. Group 4 - Neomycin, symbiotics but no bowel preparation.

Intervention Type

Other

Phase

Not Specified

Primary outcome(s)

Gut microflora:

1. Nasogastric aspirate
2. Fecal samples.

Gut barrier function:

1. Bacterial Translocation
2. Intestinal Permeability.

Inflammatory response:

1. C reactive protein (CRP)
2. Interleukin-6 (IL-6)
3. AntiEndotoxin core antibody (IgM Endo CAB).

Septic morbidity:

1. Wound infection
2. Intra-abdominal abscess
3. Chest Infection
4. UTI.

Key secondary outcome(s)

Not provided at time of registration

Completion date

01/10/2005

Eligibility

Key inclusion criteria

Using bacterial translocation as the primary end point for sample size calculation, if the sample size in each of the 4 groups is 20, $\alpha=0.05$ (2-tailed) a chi-square test will have 80% power to detect a difference in proportions.

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Not Specified

Sex

Not Specified

Key exclusion criteria

Failure to obtain informed consent, patients with co-existing infections, patients on antibiotics in the previous 2 weeks before recruitment and patients with severe hepatic or renal failure.

Date of first enrolment

19/07/2004

Date of final enrolment

01/10/2005

Locations

Countries of recruitment

United Kingdom

England

Study participating centre
Dept of General Surgery
Scarborough
United Kingdom
YO12 6QL

Sponsor information

Organisation
Department of Health

Funder(s)

Funder type
Government

Funder Name
North Yorkshire Alliance R&D Unit (UK)

Funder Name
Scarborough & NE Yorkshire Healthcare NHS Trust (UK)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary
Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/05/2007		Yes	No